

# Deafness Genes in Israel: Implications for Diagnostics in the Clinic

ZIPPORA BROWNSTEIN AND KAREN B. AVRAHAM

*Department of Human Molecular Genetics and Biochemistry, Sackler School of Medicine, Tel Aviv University, Tel Aviv, 69978 Israel*

**ABSTRACT:** The identification of the molecular basis of deafness in the last decade has made a remarkable impact on genetic counseling and diagnostics for the hearing impaired population. Since the discovery of the most prevalent form of deafness associated with mutations in the *GJB2* (connexin 26) gene, many other genes have been found worldwide, with a subset of these, including unique mutations, in Israel. Here, we review the current status of deafness genes in Israel and report one known mutation in a syndromic form of deafness, Usher syndrome, described in the Jewish Israeli population for the first time. In the future, the identification of specific mutations may be relevant for specific types of treatment. (*Pediatr Res* 66: 128–134, 2009)

## Hereditary Hearing Loss

Patients with hearing loss (HL) and parents of hearing impaired children often wish to discover the cause of this most common sensory characteristic, one in which both medical geneticists and otolaryngologists are involved. The presence of a particular mutation may aid in making decisions regarding treatments and interventions, as well as prognosis for the future. Nonsyndromic deafness affects approximately 1 in 1000 newborns and 4% of people aged younger than 45 y (1). This number rises dramatically in regions where consanguinity is an acceptable social custom and is encouraged (2,3). Progressive HL affects an even greater proportion of the population. Approximately one quarter of persons develop HL by the age of 65 (1) and ~50% of persons by age 80 (4). HL is estimated to be inherited or has a genetic basis in approximately 60% of cases (5). Genetic HL includes ~30% syndromic HL (SHL), associated with other clinical findings such as blindness and neurologic defects, and 70% non-SHL (NSHL), where HL is an isolated characteristic (6).

Because in diverse populations different genes are involved, genetic evaluation of each population should be based on an empirical study of that population. In this article, we review the genes involved in HL in Jewish Israeli samples, and we use the empirical data to develop a genetic testing algorithm for use in the Israeli population. We have updated the number of patients screened for connexin 26 mutations since this data

were published in 2000 (7). We also report a mutation in the harmonin gene in Usher syndrome (USH), not previously detected in the Jewish population in Israel. Although the particular list of genes may be different in other populations, the general scheme of the algorithm is relevant to other populations as well.

SHL includes nearly 400 forms of deafness combined with other clinical findings [(6); Hereditary Hearing Loss Homepage, available at <http://webh01.ua.ac.be/hhh/> and Online Mendelian Inheritance in Man, available at <http://www.ncbi.nlm.nih.gov/sites/entrez?db=omim>]. The severity of HL varies between and within syndromes, and some of the syndromes are rare. However, HL is most significant in several well-characterized syndromes. In the deaf Jewish Israeli sample, mutations were detected in two genes involved in USH. USH is defined by congenital NSHL, retinitis pigmentosa (RP), and vestibular dysfunction, and it accounts for more than 50% of the deaf-blind population (8), approaching a prevalence of 1 in 10,000 between the ages of 30 and 49 (9).

Nonsyndromic deafness is represented by extreme genetic heterogeneity, as up to 130 loci have been mapped and 45 nuclear genes have been identified to date, including 27 *DFNB* (autosomal recessive) genes, 22 *DFNA* (autosomal dominant) genes (7 genes are involved in both dominant and recessive forms), 1 X-linked gene, and 2 mitochondrial genes (<http://webh01.ua.ac.be/hhh/>). The most common form of NSHL is inherited in an autosomal recessive mode, which accounts for up to 80% of cases (10). Autosomal dominant inherited NSHL is involved in 20% of cases and X-linked in 1%. Mitochondrial HL has been described in approximately 1 in 500 of NSHL in Europeans (11,12) and is higher in Spain (5,13). Autosomal recessive NSHL is mostly due to cochlear defects (sensorineural deafness) (14), and in the majority of cases, it is congenital profound, whereas autosomal dominant HL is mostly postlingual and progressive (10).

## Deafness Genes in the Jewish Israeli Population

To achieve our goal in defining the number of genes associated with deafness in the Jewish Israeli population in Israel, we worked with most genetic centers in Israel. The project was approved by the National Helsinki Committee for

---

Received January 29, 2009; accepted March 13, 2009.

Correspondence: Zippora N. Brownstein, Ph.D., Department of Human Molecular Genetics and Biochemistry, Sackler School of Medicine, Tel Aviv University, Tel Aviv 69978, Israel; e-mail: [brownste@post.tau.ac.il](mailto:brownste@post.tau.ac.il)

Supported by the Grants NIH R01 DC005641, the European Commission FP6 Integrated Project EUROHEAR LSHG-CT-20054-512063, and the Hirschfeld Foundation.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site ([www.pedresearch.org](http://www.pedresearch.org)).

---

**Abbreviations:** EVA, enlarged vestibular aqueduct; HL, hearing loss; NSHL, nonsyndromic hearing loss; RP, retinitis pigmentosa; USH, Usher syndrome

Human Genetic Research of the Israel Ministry of Health in a multicenter study, including the Haim Sheba Medical Center, Rabin Medical Center, Hadassah Medical Center, Sourasky Medical Center, and the Wolfson Medical Center. Blood samples were obtained after informed consent from each individual and, in case of individuals under 18, from their parents.

Overall, 230 unrelated deaf Israeli Jews were investigated for deafness-causing genes. All subjects were tested for *GJB2* mutations by sequencing the coding exon of the gene. The other genes involved in deafness in this population were screened in all subjects in whom *GJB2* mutations were excluded. Some of the subjects belonged to large families with many deaf family members. Several such families were ascertained and genome scans were conducted, leading to mapping or cloning of the genes involved in deafness. Most of the resulting data were published previously and references are provided while describing the genes involved. A harmonin mutation, previously undetected in the Israel Jewish population, was identified in the course of this study (see later).

Anonymous DNA samples from hearing Jews of all ethnic groups were obtained from the Sheba Medical Center and from the Rabin Medical Center and used as controls. In the Jewish Israeli population, as in many populations in the world, the most common gene involved in HL is the *GJB2* gene encoding the connexin 26 (Cx26) protein (7). We found that mutations only in *GJB2* are associated with 27% of congenital HL in our Israeli sample (Table 1). Thus far, mutations in four additional genes are known to lead to NSHL in the Jewish Israeli population, among which is *GJB6*, encoding connexin 30 (Cx30) (15). Both the connexin encoding genes, *GJB2* and *GJB6*, are involved in 31% of NSHL in the Jewish Israeli population studied. Two more genes, *MYO3A* (encoding myosin IIIA) (16) and *SLC26A4* (encoding pendrin) (17), are involved in autosomal recessive deafness, and one gene, *POU4F3* (18), is associated with autosomal dominant HL. In addition, one dominant gene has been mapped for otosclerosis, *OTSC4* (19). However, according to our findings, the mutations in these six genes are involved in only approxi-

mately one third of the genetic deafness cases in the portion of the Israeli population studied. For recessive NSHL, a minimum estimate of eight to nine genes was calculated in the general deaf Israeli population (20), with a range of 7–22 loci within and between intraethnic and interethnic matings, respectively, indicating that in different ethnic groups, different loci may exist. We describe four recessive genes causing HL in less than half of our sample, suggesting that there is still more than half way to go to identify all the genes involved in deafness in the Jewish Israeli population.

### Autosomal Recessive Genes *GJB2* and *GJB6*

*GJB2* (NM\_004004.4) was the first autosomal recessive gene found to cause NSHL, discovered in 1997. This was especially compelling because it has turned out to be the largest single contributor to prelingual NSHL. The *GJB2* gene maps to chromosome 13q11-q12 and contains two exons of which only one is coding. *GJB2* encodes Cx26, a gap junction protein of 226 amino acids. Cx26 belongs to a family of more than 20 members that share a common structure of four transmembrane segments. Most cell types express more than one connexin species, which may form homomeric or heteromeric connexons. In the auditory system, intercellular channels are formed predominantly by Cx26 but also by Cx30, Cx31, and Cx43 (21,22). Cx26 seems to be involved in maintaining a high-extracellular K concentration in the endolymph by facilitating the circulation of K<sup>+</sup> ions (23,24).

*GJB6* (NM\_001110219.1) has five exons, of which only one is coding. The Cx30 protein of 261 amino acids, encoded by this gene, shares 76% identity with human Cx26 (25). *GJB6*, as *GJB2*, maps to 13q11-q12 and colocalizes with *GJB2* in the same inner ear structures: in the supporting cells of the organ of Corti, in the stria vascularis, and in the spiral ligament (22). Connexons composed of Cx26 can bind connexons composed of Cx30 to form heterotypic gap junction channels (26).

**Table 1.** *GJB2* (Cx26) and *GJB6* (Cx30) mutations in the Jewish Israeli population (70 *GJB2*/*GJB6* out of 230 NSHL probands)

|  | Mutations                                | Number of individuals | Ethnic group                       | Frequency among <i>GJB2</i> / <i>GJB6</i> deafness (%) | Frequency among NSHL (%) |
|--|--|-----------------------|------------------------------------|--|--------------------------|
| Homozygotes for <i>GJB2</i> mutations                          | c.35delG                                 | 16                    | All                                | 22.9   | 7.0                      |
|  | c.167delT                                | 21                    | Ashkenazi                          | 30.0   | 9.1                      |
|  | c.51del12insA                            | 3                     | Buchari                            | 4.3  | 1.3                      |
| Compound heterozygotes for <i>GJB2</i> mutations               | c.35delG/c.167delT                       | 14                    | Ashkenazi, Ashkenazi-other, Syrian | 20.0   | 6.1                      |
|  | c.35delG/p.L90P (c.269 T <sup>®</sup> C) | 3                     | Iraqi-other                        | 4.3  | 1.3                      |
|  | p.L90P/IVS1+1(G->A)                      | 1                     | Iraqi                              | 1.4  | 0.5                      |
|  | c.51del12insA/p.W24X                     | 1                     | Buchari                            | 1.4  | 0.5                      |
|  | c.35delG/p.R32C                          | 1                     | Ashkenazi                          | 1.4  | 0.5                      |
|  | p.V37I/c.35delG                          | 1                     | Ashkenazi                          | 1.4  | 0.5                      |
| <b>Deafness due to <i>GJB2</i> mutations</b>                   |  | <b>61</b>             |                                    | <b>87.1</b>  | <b>26.8</b>              |
| Homozygotes for <i>GJB6</i> mutations                          | del( <i>GJB6</i> -D13S1830)              | 1                     | Ashkenazi                          | 1.4  | 0.5                      |
| Double heterozygotes: <i>GJB2</i> / <i>GJB6</i>                | del( <i>GJB6</i> -D13S1830)/c.35delG     | 3                     | Ashkenazi                          | 4.3  | 1.3                      |
|  | del( <i>GJB6</i> -D13S1830)/c.167delT    | 5                     | Ashkenazi                          | 7.1  | 2.2                      |
| <b>Total deafness due to <i>GJB2</i>/<i>GJB6</i> mutations</b> |  | <b>70</b>             |                                    | <b>99.9</b>  | <b>30.8</b>              |

More than 100 mutations in *GJB2* (the Connexin-Deafness Homepage, <http://davinci.crg.es/deafness/>) are responsible for up to 50% of severe to profound prelingual recessive deafness in several worldwide populations (27,28). In Israel, in 2000, *GJB2* was reported to be involved in 38.7% of the general deaf Israeli population (7) and for 70.4% of the deaf Ashkenazi Jews (29). In 2001, Lerer *et al.* (29) identified a *GJB6* deletion in the Ashkenazi Jewish population, found on one allele in conjunction with a *GJB2* mutation. The same *GJB6* deletion of 342 kb was reported in Spain (31). The deletion did not directly affect the coding region of *GJB2*, but truncated the adjacent *GJB6* gene, which encodes Cx30 (30–32), and it has been shown in skin that this deletion abolishes *GJB2* expression, probably by deleting a regulatory element (33). The *GJB6* deletion was found to accompany in *trans*, the only *GJB2* mutant allele in heterozygous affected subjects (double heterozygosity), and it was also found in homozygosity in several cases worldwide, including in 1 individual in Israel (15,31,32). Double heterozygotes for *GJB2* and *GJB6* mutations manifest the same phenotypes as homozygotes for *GJB2* as well as homozygotes for *GJB6* (15,30,31). This deletion, named del(*GJB6*-D13S1830), was the accompanying mutation in 50% of the heterozygotes with only one *GJB2* mutant allele in different world populations (31) and in 36% (8 of 22) of the Jewish Cx26 heterozygotes in our sample.

In our study, out of 230 Israeli probands screened for mutations in *GJB2* and *GJB6*, 61 (27%) were homozygotes or compound heterozygotes for 8 mutations in *GJB2* and 22 were heterozygotes. Eight of 22 *GJB2* heterozygotes (36%) were double heterozygotes *GJB2*/*GJB6*, and one was homozygous for the del(*GJB6*-D13S1830) mutation. Combining the *GJB2* and *GJB6* mutations, the estimated prevalence of NSHL involving connexins is 31%, lower than the 38% reported previously (7), but still remaining the main known cause for congenital deafness among the Jews in Israel. Table 1 summarizes the *GJB2* and *GJB6* mutations detected in the Jewish Israeli sample.

The most common mutations in the Israeli population are the biallelic truncating mutations, c.35delG and c.167delT, and therefore, the degree of HL of most individuals with connexin-related HL ranges from severe to profound. All the deafness in which del(*GJB6*-D13S1830) is involved is severe

to profound as well. The few cases in which the nontruncating mutations p.V37I and p.L90P are associated, mild-to-moderate HL is observed (Table 2). All our findings are compatible with those of other populations, as was presented by a worldwide multicenter study comprising 1531 genetically and audiometrically documented individuals with connexin autosomal recessive HL (34).

Distinct mutations were associated with certain ethnic groups and were not detected in Jewish deaf individuals originating elsewhere, *i.e.* c.167delT was found mostly among Ashkenazi Jews and in several cases of deaf individuals originating in Syria; the del(*GJB6*-D13S1830) mutation was detected only among Ashkenazi Jews; the c.51del12insA frame shift mutation was found to be unique to Bukharan deaf Jews; whereas the p.L90P missense mutation was identified only in deaf individuals originating in Iraq. In many genetic clinical laboratories in Israel, only the c.35delG and the c.167delT mutations are screened by restriction enzyme assays. Based on our findings, it is clear that at least in some ethnic groups other mutations exist. Because the *GJB2* gene has only one coding exon and the other mutations detected in Israel are also found frequently outside of the Jewish community, the complete coding region of the *GJB2* gene should be sequenced as a routine, and considering ethnicity of the deaf individual, additional tests for the splice site mutation and the *GJB6* deletion should be performed.

#### Autosomal Recessive Gene *MYO3A*

Myosin IIIA (*MYO3A*; NM\_017433.4) maps to chromosome 10p11.1 and contains 35 exons (<http://genome.ucsc.edu>). The 1616 amino acid protein encoded by this gene belongs to the myosin superfamily. Myosins are actin-dependent motor proteins and are categorized into conventional myosins (class II) and unconventional myosins (classes I and III through XV) based on their variable C-terminal cargo-binding domains. Class III myosins, expressed in photoreceptors, are hybrid motor-signaling molecules, with an N-terminal kinase domain, highly conserved head and neck domains, and a class III-specific tail domain. Expression of this gene is highly restricted, with the strongest expression in retina and cochlea (16).

**Table 2.** Severity of the *GJB2* and *GJB6* mutations in a Jewish Israeli sample

| Severity*       | Mutation          |                     |                    |                 |                  |               |                             |                      |                 |                 |                  |  |
|-----------------|-------------------|---------------------|--------------------|-----------------|------------------|---------------|-----------------------------|----------------------|-----------------|-----------------|------------------|--|
|                 | 35delG/<br>35delG | 167delT/<br>167delT | 35delG/<br>167delT | 35delG/<br>Δ30† | 167delT/<br>Δ30† | Δ30‡/<br>Δ30‡ | 51del12InsA/<br>51del12InsA | 51del12InsA/<br>W24X | 35delG/<br>L90P | 35delG/<br>R32C | 167delT/<br>V37I |  |
| Profound        | 11                | 10                  | 9                  | 2               | 4                | 1             | 1                           | 1                    |                 | 1               |                  |  |
| Severe profound | 1                 | 1                   |                    | 1               | 1                |               |                             |                      |                 |                 |                  |  |
| Severe          |                   | 5                   | 1                  |                 |                  |               |                             |                      |                 |                 |                  |  |
| Moderate severe | 1                 |                     | 2                  |                 |                  |               | 1                           |                      |                 |                 |                  |  |
| Moderate        |                   |                     | 1                  |                 |                  |               | 1                           |                      | 2               |                 |                  |  |
| Mild moderate   |                   | 1                   |                    |                 |                  |               |                             |                      |                 |                 | 1                |  |
| Mild            |                   |                     |                    |                 |                  |               |                             |                      | 1               |                 |                  |  |
| Total           | 13                | 17                  | 13                 | 3               | 5                | 1             | 3                           | 1                    | 3               | 1               | 1                |  |

Subjects are classified based on audiogram data on the better performing ear.

\*Profound HL, >95dB; severe HL, 71–95 dB; moderate HL, 41–70 dB; Mild HL 21–40 dB.

†*GJB6*-D13S1830.

Three different autosomal recessive, loss-of-function mutations in *MYO3A* are involved in progressive NSHL in an extended Israeli family, originating in Mosul, Iraq (16). The HL begins in the second decade, first affects the high frequencies, and by age 50 is severe in high and middle frequencies and moderate at low frequencies. The mutations detected include a nonsense mutation c.3126T->G, p.Y1042X causing protein truncation at the junction of the head and neck domains of the myosin IIIA protein; a mutation in the splice acceptor of intron 17, 1777(-12)G->A, leading to deletion of exon 18 and protein truncation at codon 668 in the myosin head domain; and a mutation in the splice acceptor of intron 8, 732(-2)A->G, resulting in an unstable message, as revealed by the absence of message from this allele in carriers of the mutation. Of 18 affected individuals in this family, 7 are homozygous and 11 are compound heterozygous for pairs of mutant alleles. A genotype-phenotype correlation was observed among the deaf members of the family. Between the ages of 25 and 50, hearing across all frequencies was significantly poorer, and the rate of progress was higher among the individuals homozygous for the nonsense mutation comparing to the compound heterozygotes for the nonsense mutation and either of the splice mutations.

Among 172 Iraqi Jewish hearing controls, four (2.3%) carried the nonsense mutation; none carried either splice mutation. None of the three mutations were observed among 96 Ashkenazi Jewish hearing controls (Table S1, Supplemental Digital Content 1, <http://links.lww.com/PDR/A43>).

As for the phenotypic-clinical aspect, the *MYO3A* gene is unique in that 1) mutations in this gene are involved in an uncharacteristic autosomal recessive HL, which unlike the other recessive deafness is late onset and progressive, and 2) three different *MYO3A* mutations underlies deafness in the same family. The existence of multiple *MYO3A* mutations in the same kindred also suggests that other mutations in this gene may be involved in recessive late onset HL.

#### Autosomal Recessive Gene *SLC26A4*

More than 100 mutations in the *PDS* (*SLC26A4*) gene are involved in both Pendred syndrome (PS) and NSHL *DFNB4* associated with temporal bone abnormalities (35). The *SLC26A4* gene (NM\_000441.1), located on the long arm of chromosome 7, contains 1 noncoding and 20 coding exons and encodes pendrin, a protein of 780 amino acids, expressed in various tissues, including the inner ear, thyroid, and kidney.

The HL caused by mutations in *SLC26A4* is mostly congenital, nonsyndromic high frequency, but sometimes postlingual progressive SNHL and associated with malformations of the temporal bone. The temporal bone abnormalities vary, ranging from isolated enlarged vestibular aqueduct (EVA) to Mondini dysplasia, an abnormality in which the normal cochlear spiral of 2.5 turns is replaced by a hypoplastic coil of 1.5 turns. Albert *et al.* (35) estimated that *SLC26A4* mutations are involved in 4% of NSHL, which emphasizes the importance of the molecular confirmation of the *PDS* gene in the diagnosis of deafness and particularly when temporal bone abnormality conditions are involved.

The c.1458\_1459insT mutation of the *SLC26A4* gene was detected in a Jewish Iranian proband with NSHL and EVA after excluding mutations in *Cx26* (17). The deaf individual had progressive postlingual HL that began at age 6 and progressed to profound deafness. In his mid twenties, he underwent cochlear implantation. In his presurgery CT, EVA was observed and therefore the *SLC26A4* gene was sequenced. The c.1458\_1459insT mutation was detected on both alleles. The mutation causes a frameshift at position 487 in exon 13 (of 21), leading to a stop codon at position 526 in exon 14 of the *SLC26A4* gene, p.Ile487TyrfsX39.

An additional 203 deaf probands, in which mutations in the connexins were excluded, were screened for the c.1458\_1459insT mutation (17). Three individuals with Mondini dysplasia were included in this population. All 203 probands were wild type for this nucleotide. In total, the probability of the c.1458\_1459insT mutation of the *PDS* gene, among the Jewish deaf population in Israel, is 0.5% (1 of 204). Normal hearing controls (310) were screened, including 21 of Iranian ethnic origin, and all 310 were found to be WT, *i.e.* no carriers of the mutation were detected (Table S1, Supplemental Digital Content 1, <http://links.lww.com/PDR/A43>).

#### USH Gene *PCDH15*

Three clinical subtypes, USH1–3, are described based on the characteristics of deafness, vestibular involvement, and RP. Defects in protocadherin 15 (*PCDH15*) cause USH1F (36,37). USH1 is the most common and most severe and characterized by congenital profound deafness, onset of RP in the beginning of the second decade of life, and constant vestibular dysfunction.

*PCDH15* (NM\_033056.3) is located on chromosome 10q11.2–q21. The *PCDH15* gene contains 33 exons, one of which is noncoding, and it is predicted to encode a protein of 1955 amino acids. Mice carrying a mutation in *Pcdh15* show disorganization in the placement of the stereocilia. It was suggested that *PCDH15* plays a role in regulation of planar polarity in the sensory neuroepithelium of the inner ear. Members of the cadherin superfamily are also required in the eye and inner ear for maintenance of normal function (38). Recently, *PCDH15* has been shown to be a part of the tip link complex, together with cadherin 23 (39).

The R245X mutation is caused by a C to T transition at position 733 in exon 8 of the *PCDH15* gene, and it leads to the substitution of an arginine by a stop codon in the extracellular, second (of 11) cadherin domain. The R245X mutation accounts for more than half (58.3%) of USH in the Ashkenazi Jewish population. This mutation was detected only among Ashkenazi Jews and was not detected in other Jewish ethnic groups (40). In our study, Ashkenazi Jewish individuals ( $n = 59$ ) diagnosed with NSHL were screened for the R245X mutation (41). Twenty of the probands were younger than 10 y, and 39 were older than 10 y. This age distinction was chosen because RP is typically diagnosed after the age of 10 (42). In the first group of children under 10, 2 of 20 (10%) were homozygous for R245X. In the older group, no homozy-

gotes were detected; but 1 of 39 (2.6%) was heterozygous for the R245X mutation.

To determine the carrier rate, 505 DNA samples of hearing Ashkenazi Jews were screened for the R245X mutation. Five out of 505 (0.99%) were heterozygous for the mutation (Table S1, Supplemental Digital Content 1, <http://links.lww.com/PDR/A43>). The R245X mutation was not detected among 293 non-Ashkenazi Jews screened elsewhere (40), and the mutation is assumed to be a founder effect in the Ashkenazi ethnic group.

These findings led to the inevitable consequence that Ashkenazi children diagnosed with NSHL should be evaluated for the R245X mutation to determine whether they will develop RP before clinical signs of this retinal defect, detectable by fundoscopy and ERG. Habilitation can then begin before loss of vision.

### USH Gene *USH1C* (Harmonin)

Mutations in harmonin are associated with *USH1C*, a subtype of *USH1*, and are characterized by congenital deafness, RP, and vestibular dysfunction. The *USH1C* gene (NM\_153676.2) maps to chromosome 11p15.1 and contains 27 exons, encoding a protein of 899 amino acids. Mutations in *USH1C*, as well as mutations in myosin VIIa (*MYO7A*), *CDH23*, *PCDH15*, or sans lead to *USH1* in humans and hair bundle disorganization in mice. It was suggested that harmonin anchors the tip link upper end (likely to be made of cadherin 23) to the stereocilia actin core, hence, playing a key role in hair bundle cohesion and proper orientation of the stereocilia; it is also involved in stereociliary elongation (43).

The c.238–239InsC mutation causes a frameshift at position 78 in exon 3 (of 27), leading to a stop codon at position 148 in exon 5 of the *USH1C* gene, p.Tyr78TyrfsX70 (44). Previously identified only outside of Israel (44,45), this mutation was recently detected in Israel in two brothers with congenital deafness, born to unrelated Ashkenazi-Jewish parents. We describe this data here for the first time. The two children, 5- and 1-y-old, were misdiagnosed as having NSHL. Homozygosity mapping was conducted using the GeneChip Human Mapping 250 K Nsp Array of Affymetrix, revealing a 13.63-Mb homozygous region on chromosome 11p15.4-p15.1. Homozygosity was further confirmed by five microsatellite markers, which spanned the region (including the commercially available D11S4149, D11S4116, and D11S902; Orly Elpeleg, personal communication, Hadassah Hebrew University Medical Center, Jerusalem, Israel, 2008). Of the 174 genes in this region, the harmonin gene at chromosome 11p15.1-p14, associated with *USH1C* and with nonsyndromic recessive deafness (DFNB18), was a good candidate. The mutation c.238-239InsC was detected by sequencing this gene. Fundoscopy confirmed onset of RP in the older brother.

Exon 3 of the harmonin gene was sequenced in addition to 20 *USH* unrelated patients and none of them harbored the mutation. Normal hearing controls were screened (384), including 271 of Ashkenazi origin and 113 of non-Ashkenazi ethnic groups. No carriers of the mutation were detected (Table S1, Supplemental Digital Content 1, <http://links.lww.com/PDR/A43>).

Because harmonin, as *PCDH15*, is a subtype of *USH1*, in which the onset of RP is in the beginning of the second decade of life, children are often misdiagnosed as having NSHL

instead of *USH*. Therefore, children with no family history of deafness and with no *GJB2/GJB6* mutations should be evaluated for the c.238-239InsC mutation.

### Autosomal Dominant Gene *POU4F3*

*POU4F3* is a member of the POU family of transcription factors, with expression mostly restricted to hair cells of the inner ear (46,47), and it is essential for inner ear hair cell maintenance. The 2-exon *POU4F3* gene (NM\_002700.1) maps to chromosome 5q31 and encodes a protein of 338 amino acids. In hearing impaired members of one large Israeli Jewish family originating in Lybia, an 8 bp deletion was identified, starting at nucleotide 884 (884del8) of exon 2 of the *POU4F3* gene (18). The predicted result of this deletion is a frameshift beginning at the second nucleotide of codon 295 and a premature translation stop at position 302, p.Ile295ThrfsX7. The 884del8 mutation in the *POU4F3* gene is associated with late onset, progressive, NSHL. This deletion leads to a frameshift, predicted to cause a stop codon to be formed prematurely in the first helix of the POU-homeodomain. Mutant *POU4F3* loses most of its transcriptional activity and most of its ability to bind to DNA. Moreover, although wild-type *POU4F3* is found exclusively in the nucleus, it was demonstrated that the mutant protein is localized both to the nucleus and the cytoplasm, where transcriptional activity cannot be carried out (48).

Affected members suffer from progressive high-tone, autosomal dominant SNHL, with onset between ages 18 and 30 y. The hearing impairment becomes more severe with time, eventually causing significant HL across the spectrum at all frequencies (49). Two novel missense mutations were recently discovered, in the Dutch population, with a similar phenotype to the Israeli family (50).

Twenty-seven autosomal dominant deaf screened were negative for the mutation, resulting in the prevalence of 3.5% (1 of 28) of the mutation in the *POU4F3* gene among autosomal dominant HL in our study. To determine the carrier rate, 114 unrelated individuals of various North African and Middle Eastern Jewish ancestries were tested for the mutation. None of the 228 control chromosomes tested carried the 8 bp deletion in *POU4F3* found in the Jewish Lybian family.

### A Mapped Locus for Otosclerosis *OTSC4*

Otosclerosis is a common bone disorder with a prevalence of 0.2 to 1% among adults, leading to a progressive hearing impairment that begins as a conductive HL, and might develop into a mixed to SNHL, with an autosomal dominant mode of inheritance with reduced penetrance. The age of onset for otosclerosis is usually 20–40 y and in most cases both ears are involved, but HL is often asymmetric (51). A definitive diagnosis of otosclerosis can be made only by surgery, but existence of family history of the disease and some other clinical characterizations are considered sufficient to diagnose otosclerosis in a family where only some members were surgically confirmed (52–54). Both environmental and genetic factors have been implicated in otosclerosis. A 2:1 female-to-male predominance suggests hormonal involvement as well (55).

We mapped a new otosclerosis locus, *OTSC4*, to chromosome 16q21-23.2 in a Yemenite Israeli family with otosclerosis (19). The hearing impairment in the family ranged from conductive to mixed- to SNHL, from unilateral to bilateral HL, with variable severity, age of onset, and progression. The considerable inter-subject and interaural variability within the same family might be because of the different stages of the activity of the otosclerotic lesion, which are thought to be directly correlated to the type and severity of HL and audiogram configuration.

### Implications for Patients With HL


Except for the *GJB2/GJB6* genes involved in one third of the nonsyndromic deaf population of our study, all the other deafness genes are of low prevalence, frequently detected in one family only (Table S1, Supplemental Digital Content 1, <http://links.lww.com/PDR/A43>). There is no doubt that *GJB2* should be sequenced first in each hearing impaired individual, followed by screening for the large deletion truncating *GJB6* in all the Ashkenazi deaf in which *GJB2* was ruled out. Today, it is tedious and not cost-effective to screen the known genes in Israel routinely after mutations in *GJB2/GJB6* are ruled out. Therefore, we suggest a diagnostic algorithm (Fig. 1) for Jewish individuals presenting with HL, depending on the mode of inheritance, ethnic origin, and clinical/audiological manifestation, and taking into account the most cost-efficient method to address the molecular diagnoses available today. Furthermore, other genes that have been implicated in HL in other parts of the world but not yet found in Israel should be considered when the relevant phenotype is discovered. For example, inherited in a dominant form for nonsyndromic deafness, *WFS1* mutations are associated with low frequency HL (56). Patients with dominant *COCH* mutations have a characteristic vestibular dysfunction that is similar to Meniere's disease and is associated with HL that begins in the third decade, with high frequencies affected first (57). If aminoglycosidase damage is suspected, the mitochondrial 12S rRNA mutation should be examined, particularly if the HL appears to

follow a maternal form of inheritance (58). *TECTA* mutations may be responsible when prelingual, nonprogressive, midfrequency hearing impairment is observed (59). Recessive auditory neuropathy, characterized by the absence of detectable responses to auditory evoked potentials (AEP) and the presence of bilateral TEOAE, may be associated with mutations in the *OTOF* gene (60,61). The identification of genes in the world is a dynamic process, with new genes and mutations being found monthly. Therefore, the literature and the Hereditary Hearing Loss Homepage should be scrutinized routinely to determine whether there are additional genes that should be examined in a patient presenting with suspected genetic hearing impairment. In the future, DNA arrays may be designed that will make screening multiple genes/variants more cost efficient. Furthermore, as we gain knowledge of a complete repertoire of genes involved in hearing, and technology advances, the approach to future treatment and intervention will change accordingly. In particular, there is promise for gene and cell therapy (62,63) with this increased knowledge and advanced tools, and the future holds great promise for the hearing impaired population.

**Acknowledgments.** We thank all the families who contributed to the studies described in this review, without whom this work could not have taken place. Their participation in the study enabled many more families to learn about their genetic basis of HL. We also thank our collaborators over the years that have been involved in patient ascertainment and gene identification and contributed an essential component to the success of this work. We especially thank Prof. Moshe Frydman, a devoted clinician and researcher, whose guidance and dedication has carried us throughout this project.

### REFERENCES

- Nadol JB Jr 1993 Hearing loss. *N Engl J Med* 329:1092-1102
- Khabori MA, Patton MA 2008 Consanguinity and deafness in Omani children. *Int J Audiol* 47:30-33
- Sajjad M, Khattak AA, Bunn JE, Mackenzie I 2008 Causes of childhood deafness in Pukhtoonkhwa Province of Pakistan and the role of consanguinity. *J Laryngol Otol* 122:1057-1063
- Morton NE 1991 Genetic epidemiology of hearing impairment. *Ann N Y Acad Sci* 630:16-31
- Nance WE 2003 The genetics of deafness. *Ment Retard Dev Disabil Res Rev* 9:109-119
- Gorlin RJ, Toriello HV, Cohen MM 1995 Hereditary Hearing Loss and its Syndromes. Oxford University Press, Oxford
- Sobe T, Vreugde S, Shahin H, Berlin M, Davis N, Kanaan M, Yaron Y, Orr-Urtreger A, Frydman M, Shohat M, Avraham KB 2000 The prevalence and expression of inherited connexin 26 mutations associated with nonsyndromic hearing loss in the Israeli population. *Hum Genet* 106:50-57
- Boughman JA, Vernon M, Shaver KA 1983 Usher syndrome: definition and estimate of prevalence from two high-risk populations. *J Chronic Dis* 36:595-603
- Hope CI, Bunday S, Proops D, Fielder AR 1997 Usher syndrome in the city of Birmingham—prevalence and clinical classification. *Br J Ophthalmol* 81:46-53
- Van Camp G, Willems PJ, Smith RJ 1997 Nonsyndromic hearing impairment: unparalleled heterogeneity. *Am J Hum Genet* 60:758-764
- Bitner-Glindzicz M, Pembrey M, Duncan A, Heron J, Ring SM, Hall A, Rahman S 2009 Prevalence of mitochondrial 1555A->G mutation in European children. *N Engl J Med* 360:640-642
- Vandebona H, Mitchell P, Manwaring N, Griffiths K, Gopinath B, Wang JJ, Sue CM 2009 Prevalence of mitochondrial 1555A->G mutation in adults of European descent. *N Engl J Med* 360:642-644
- Estivill X, Govea N, Barcelo E, Badenas C, Romero E, Moral L, Scozzari R, D'Urbano L, Zeviani M, Torroni A 1998 Familial progressive sensorineural deafness is mainly due to the mtDNA A1555G mutation and is enhanced by treatment of aminoglycosides. *Am J Hum Genet* 62:27-35
- Petersen MB, Willems PJ 2006 Non-syndromic, autosomal-recessive deafness. *Clin Genet* 69:371-392
- Del Castillo I, Moreno-Pelayo MA, Del Castillo FJ, Brownstein Z, Marlin S, Adina Q, Cockburn DJ, Pandya A, Siemerling KR, Chamberlin GP, Ballana E, Wuyts W,



**Figure 1.** A flow chart demonstrating a diagnostic algorithm for hearing impaired individuals in Israel. The clinical evaluation includes a full medical history and physical exam, as well as ophthalmologic and audiologic evaluations. This chart is a model for other populations, demonstrating the relevance of diagnosing the molecular basis of HL and its application to treatment of patients.

- Maciel-Guerra AT, Alvarez A, Villamar M, Shohat M, Abeliovich D, Dahl HH, Estivill X, Gasparini P, Hutchin T, Nance WE, Sartorato EL, Smith RJ, Van Camp G, Avraham KB, Petit C, Moreno F 2003 Prevalence and evolutionary origins of the del(GJB6-D13S1830) mutation in the DFNB1 locus in hearing-impaired subjects: a multicenter study. *Am J Hum Genet* 73:1452-1458
16. Walsh T, Walsh V, Vreugde S, Hertzano R, Shahin H, Haika S, Lee MK, Kanaan M, King MC, Avraham KB 2002 From flies' eyes to our ears: mutations in a human class III myosin cause progressive nonsyndromic hearing loss *DFNB30*. *Proc Natl Acad Sci USA* 99:7518-7523
17. Brownstein ZN, Dror AA, Gilony D, Migirov L, Hirschberg K, Avraham KB 2008 A novel *SLC26A4* (PDS) deafness mutation retained in the endoplasmic reticulum. *Arch Otolaryngol Head Neck Surg* 134:403-407
18. Vahava O, Morell R, Lynch ED, Weiss S, Kagan ME, Ahituv N, Morrow JE, Lee MK, Skvorak AB, Morton CC, Blumenfeld A, Frydman M, Friedman TB, King MC, Avraham KB 1998 Mutation in transcription factor *POU4F3* associated with inherited progressive hearing loss in humans. *Science* 279:1950-1954
19. Brownstein Z, Goldfarb A, Levi H, Frydman M, Avraham KB 2006 Chromosomal mapping and phenotypic characterization of hereditary otosclerosis linked to the *OTSC4* locus. *Arch Otolaryngol Head Neck Surg* 132:416-424
20. Brownstein Z, Friedlander Y, Peritz E, Cohen T 1991 Estimated number of loci for autosomal recessive severe nerve deafness within the Israeli Jewish population, with implications for genetic counseling. *Am J Med Genet* 41:306-312
21. Kelley PM, Abe S, Askew JW, Smith SD, Usami S, Kimberling WJ 1999 Human connexin 30 (*GJB6*), a candidate gene for nonsyndromic hearing loss: molecular cloning, tissue-specific expression, and assignment to chromosome 13q12. *Genomics* 62:172-176
22. Forge A, Becker D, Casalotti S, Edwards J, Marziano N, Nevill G 2003 Gap junctions in the inner ear: comparison of distribution patterns in different vertebrates and assessment of connexin composition in mammals. *J Comp Neurol* 467:207-231
23. Forge A, Becker D, Casalotti S, Edwards J, Evans WH, Lench N, Souter M 1999 Gap junctions and connexin expression in the inner ear. *Novartis Found Symp* 219:134-150; discussion 151-136
24. Beltramello M, Piazza V, Bukauskas FF, Pozzan T, Mammano F 2005 Impaired permeability to Ins(1,4,5)P<sub>3</sub> in a mutant connexin underlies recessive hereditary deafness. *Nat Cell Biol* 7:63-69
25. Grifa A, Wagner CA, D'Ambrosio L, Melchionda S, Bernardi F, Lopez-Bigas N, Rabionet R, Arbones M, Monica MD, Estivill X, Zelante L, Lang F, Gasparini P 1999 Mutations in *GJB6* cause nonsyndromic autosomal dominant deafness at *DFNA3* locus. *Nat Genet* 23:16-18
26. Dahl E, Manthey D, Chen Y, Schwarz HJ, Chang YS, Lalley PA, Nicholson BJ, Willecke K 1996 Molecular cloning and functional expression of mouse connexin-30, a gap junction gene highly expressed in adult brain and skin. *J Biol Chem* 271:17903-17910
27. Denoyelle F, Marlin S, Weil D, Moatti L, Chauvin P, Garabedian EN, Petit C 1999 Clinical features of the prevalent form of childhood deafness, *DFNB1*, due to a connexin-26 gene defect: implications for genetic counselling. *Lancet* 353:1298-1303
28. Kelsell DP, Dunlop J, Stevens HP, Lench NJ, Liang JN, Parry G, Mueller RF, Leigh IM 1997 Connexin 26 mutations in hereditary non-syndromic sensorineural deafness. *Nature* 387:80-83
29. Lerer I, Sagi M, Malamud E, Levi H, Raas-Rothschild A, Abeliovich D 2000 Contribution of connexin 26 mutations to nonsyndromic deafness in Ashkenazi patients and the variable phenotypic effect of the mutation 167delT. *Am J Med Genet* 95:53-56
30. Lerer I, Sagi M, Ben-Neriah Z, Wang T, Levi H, Abeliovich D 2001 A deletion mutation in *GJB6* cooperating with a *GJB2* mutation in trans in non-syndromic deafness: a novel founder mutation in Ashkenazi Jews. *Hum Mutat* 18:460
31. Del Castillo I 2002 A deletion involving the connexin 30 gene in nonsyndromic hearing impairment. *N Engl J Med* 346:243-249
32. Pallares-Ruiz N, Blanchet P, Mondain M, Claustres M, Roux AF 2002 A large deletion including most of *GJB6* in recessive non syndromic deafness: a digenic effect? *Eur J Hum Genet* 10:72-76
33. Common JE, Bitner-Glindzicz M, O'Toole EA, Barnes MR, Jenkins L, Forge A, Kelsell DP 2005 Specific loss of connexin 26 expression in ductal sweat gland epithelium associated with the deletion mutation del(GJB6-D13S1830). *Clin Exp Dermatol* 30:688-693
34. Snoeckx RL, Huygen PL, Feldmann D, Marlin S, Denoyelle F, Waligora J, Mueller-Malesinska M, Pollak A, Ploski R, Murgia A, Orzan E, Castorina P, Ambrosetti U, Nowakowska-Szyrwinska E, Bal J, Wiszniewski W, Jancek AR, Nekahm-Heis D, Seeman P, Bendova O, Kenna MA, Frangulov A, Rehm HL, Tekin M, Incesulu A, Dahl HH, du Sart D, Jenkins L, Lucas D, Bitner-Glindzicz M, Avraham KB, Brownstein Z, del Castillo I, Moreno F, Blin N, Pfister M, Sziklai I, Toth T, Kelley PM, Cohn ES, Van Maldergem L, Hilbert P, Roux AF, Mondain M, Hoefsloot LH, Cremers CW, Lopponeu T, Lopponeu H, Parving A, Gronskov K, Schrijver I, Roberson J, Gualandri F, Martini A, Lina-Granade G, Pallares-Ruiz N, Correia C, Fialho G, Cryns K, Hilgert N, Van de Heyning P, Nishimura CJ, Smith RJ, Van Camp G 2005 *GJB2* mutations and degree of hearing loss: a multicenter study. *Am J Hum Genet* 77:945-957
35. Albert S, Blons H, Jonard L, Feldmann D, Chauvin P, Loundon N, Sergeant-Allaoui A, Houang M, Joannard A, Schmerber S, Delobel B, Leman J, Jourmel H, Catros H, Dollfus H, Eliot MM, David A, Calais A, Drouin-Garraud V, Obstoy MF, Tran Ba Huy P, Lacombe D, Duriez F, Francannet C, Bitoun P, Petit C, Garabedian EN, Couderc R, Marlin S, Denoyelle F 2006 *SLC26A4* gene is frequently involved in nonsyndromic hearing impairment with enlarged vestibular aqueduct in Caucasian populations. *Eur J Hum Genet* 14:773-779
36. Alagramam KN, Yuan H, Kuehn MH, Murcia CL, Wayne S, Srisailopathy CR, Lowry RB, Knaus R, Van Laer L, Bernier FP, Schwartz S, Lee C, Morton CC, Mullins RF, Ramesh A, Van Camp G, Hageman GS, Woychik RP, Smith RJ, Hageman GS 2001 Mutations in the novel protocadherin *PCDH15* cause Usher syndrome type 1F. *Hum Mol Genet* 10:1709-1718
37. Ahmed ZM, Riazuddin S, Bernstein SL, Ahmed Z, Khan S, Griffith AJ, Morell RJ, Friedman TB, Wilcox ER 2001 Mutations of the protocadherin gene *PCDH15* cause Usher syndrome type 1F. *Am J Hum Genet* 69:25-34
38. Alagramam KN, Murcia CL, Kwon HY, Pawlowski KS, Wright CG, Woychik RP 2001 The mouse Ames Waltzer hearing-loss mutant is caused by mutation of *Pcdh15*, a novel protocadherin gene. *Nat Genet* 27:99-102
39. Kazmierczak P, Sakaguchi H, Tokita J, Wilson-Kubalek EM, Milligan RA, Muller U, Kachar B 2007 Cadherin 23 and protocadherin 15 interact to form tip-link filaments in sensory hair cells. *Nature* 449:87-91
40. Ben-Yosef T, Ness SL, Madeo AC, Bar-Lev A, Wolfman JH, Ahmed ZM, Desnick RJ, Willner JP, Avraham KB, Ostrer H, Oddoux C, Griffith AJ, Friedman TB 2003 A mutation of *PCDH15* among Ashkenazi Jews with the type 1 Usher syndrome. *N Engl J Med* 348:1664-1670
41. Brownstein Z, Ben-Yosef T, Dagan O, Frydman M, Abeliovich D, Sagi M, Abraham FA, Taitelbaum-Swead R, Shohat M, Hildesheimer M, Friedman TB, Avraham KB 2004 The R245X mutation of *PCDH15* in Ashkenazi Jewish children diagnosed with nonsyndromic hearing loss foreshadows retinitis pigmentosa. *Pediatr Res* 55:995-1000
42. Petit C 2001 Usher syndrome: from genetics to pathogenesis. *Annu Rev Genomics Hum Genet* 2:271-297
43. Lefevre G, Michel V, Weil D, Lepelletier L, Bizard E, Wolfrum U, Hardelin JP, Petit C 2008 A core cochlear phenotype in *USH1* mouse mutants implicates fibrous links of the hair bundle in its cohesion, orientation and differential growth. *Development* 135:1427-1437
44. Verpy E, Leibovici M, Zwaenepoel I, Liu XZ, Gal A, Salem N, Mansour A, Blanchard S, Kobayashi I, Keats BJ, Slim R, Petit C 2000 A defect in harmonin, a PDZ domain-containing protein expressed in the inner ear sensory hair cells, underlies Usher syndrome type 1C. *Nat Genet* 26:51-55
45. Zwaenepoel I, Verpy E, Blanchard S, Meins M, Apfelstedt-Sylla E, Gal A, Petit C 2001 Identification of three novel mutations in the *USH1C* gene and detection of thirty-one polymorphisms used for haplotype analysis. *Hum Mutat* 17:34-41
46. Erkman L, McEvilly RJ, Luo L, Ryan AK, Hooshmand F, O'Connell SM, Keithley EM, Rapaport DH, Ryan AF, Rosenfeld MG 1996 Role of transcription factors Brn-3.1 and Brn-3.2 in auditory and visual system development. *Nature* 381:603-606
47. Xiang M, Gan L, Li D, Chen ZY, Zhou L, O'Malley BW Jr, Klein W, Nathans J 1997 Essential role of POU-domain factor Brn-3c in auditory and vestibular hair cell development. *Proc Natl Acad Sci USA* 94:9445-9450
48. Weiss S, Gottfried I, Mayrose I, Khare SL, Xiang M, Dawson SJ, Avraham KB 2003 The *DFNA15* deafness mutation affects *POU4F3* protein stability, localization, and transcriptional activity. *Mol Cell Biol* 23:7957-7964
49. Frydman M, Vreugde S, Nageris BI, Weiss S, Vahava O, Avraham KB 2000 Clinical characterization of genetic hearing loss caused by a mutation in the *POU4F3* transcription factor. *Eur Arch Otorhinolaryngol* 126:633-637
50. Collin RW, Chellappa R, Pauw RJ, Vriend G, Oostrik J, van Druenen W, Huygen PL, Admiraal R, Hoefsloot LH, Cremers FP, Xiang M, Cremers CW, Kremer H 2008 Missense mutations in *POU4F3* cause autosomal dominant hearing impairment *DFNA15* and affect subcellular localization and DNA binding. *Hum Mutat* 29:545-554
51. Goh JP, Chan LL, Tan TY 2002 MRI of cochlear otosclerosis. *Br J Radiol* 75:502-505
52. Lopez-Gonzalez MA, Delgado F 2000 Oral vaccine in otosclerosis. *Med Hypotheses* 54:216-220
53. Sakihara Y, Parving A 1999 Clinical otosclerosis, prevalence estimates and spontaneous progress. *Acta Otolaryngol* 119:468-472
54. Zhao F, Wada H, Koike T, Ohya K, Kawase T, Stephens D 2002 Middle ear dynamic characteristics in patients with otosclerosis. *Ear Hear* 23:150-158
55. Vartiainen E 1999 Sex differences in patients with hearing impairments caused by otosclerosis. *Eur Arch Otorhinolaryngol* 126:431-433
56. Young TL, Ives E, Lynch E, Person R, Snook S, MacLaren L, Cater T, Griffin A, Fernandez B, Lee MK, King MC 2001 Non-syndromic progressive hearing loss *DFNA38* is caused by heterozygous missense mutation in the Wolfram syndrome gene *WFS1*. *Hum Mol Genet* 10:2509-2514
57. Robertson NG, Lu L, Heller S, Merchant SN, Eavey RD, McKenna M, Nadol JB Jr, Miyamoto RT, Linthicum FH Jr, Lubianca Neto JF, Hudspeth AJ, Seidman CE, Morton CC, Seidman JG 1998 Mutations in a novel cochlear gene cause *DFNA9*, a human nonsyndromic deafness with vestibular dysfunction. *Nat Genet* 20:299-303
58. Prezant TR, Agopian JV, Bohlman MC, Bu X, Oztas S, Qiu WQ, Arnos KS, Cortopassi GA, Jaber L, Rotter JJ, Shohat M, Fischel-Ghodsian N 1993 Mitochondrial ribosomal RNA mutation associated with both antibiotic-induced and nonsyndromic deafness. *Nat Genet* 4:289-294
59. Verhoeven K, Van Laer L, Kirschhofer K, Legan PK, Hughes DC, Schatteman I, Verstreken M, Van Hauwe P, Coucke P, Chen A, Smith RJ, Somers T, Officiers FE, Van de Heyning P, Richardson GP, Wachtler F, Kimberling WJ, Willems PJ, Govaerts PJ, Van Camp G 1998 Mutations in the human alpha-tectorin gene cause autosomal dominant non-syndromic hearing impairment. *Nat Genet* 19:60-62
60. Varga R, Kelley PM, Keats BJ, Starr A, Leal SM, Cohn E, Kimberling WJ 2003 Non-syndromic recessive auditory neuropathy is the result of mutations in the otoferlin (*OTOF*) gene. *J Med Genet* 40:45-50
61. Rodriguez-Ballesteros M, del Castillo FJ, Martin Y, Moreno-Pelayo MA, Morera C, Prieto F, Marco J, Morant A, Gallo-Teran J, Morales-Angulo C, Navas C, Trinidad C, Tapia MC, Moreno F, del Castillo I 2003 Auditory neuropathy in patients carrying mutations in the otoferlin gene (*OTOF*). *Hum Mutat* 22:451-456
62. Martinez-Monedero R, Oshima K, Heller S, Edge AS 2007 The potential role of endogenous stem cells in regeneration of the inner ear. *Hear Res* 227:48-52
63. Raphael Y, Kim YH, Osuni Y, Izumikawa M 2007 Non-sensory cells in the deafened organ of Corti: approaches for repair. *Int J Dev Biol* 51:649-654