

ARTICLE

# Five novel loci for inherited hearing loss mapped by SNP-based homozygosity profiles in Palestinian families

Hashem Shahin<sup>1,5</sup>, Tom Walsh<sup>\*,2,5</sup>, Amal Abu Rayyan<sup>1</sup>, Ming K Lee<sup>2</sup>, Jake Higgins<sup>2,6</sup>, Diane Dickel<sup>3</sup>, Kristen Lewis<sup>3</sup>, James Thompson<sup>3</sup>, Carl Baker<sup>3</sup>, Alex S Nord<sup>3</sup>, Sunday Stray<sup>2</sup>, David Gurwitz<sup>4</sup>, Karen B Avraham<sup>4</sup>, Mary-Claire King<sup>2,3</sup> and Moien Kanaan<sup>1</sup>

In communities with high rates of consanguinity and consequently high prevalence of recessive phenotypes, homozygosity mapping with SNP arrays is an effective approach for gene discovery. In 20 Palestinian kindreds with prelingual nonsyndromic hearing loss, we generated homozygosity profiles reflecting linkage to the phenotype. Family sizes ranged from small nuclear families with two affected children, one unaffected sibling, and parents to multigenerational kindreds with 12 affected relatives. By including unaffected parents and siblings and screening 250 K SNP arrays, even small nuclear families yielded informative profiles. In 14 families, we identified the allele responsible for hearing loss by screening a single candidate gene in the longest homozygous region. Novel alleles included missense, nonsense, and splice site mutations of *CDH23*, *MYO7A*, *MYO15A*, *OTOF*, *PJVK*, *Pendrin/SLC26A4*, *TECTA*, *TMHS*, and *TMPRSS3*, and a large genomic deletion of *Otoancorin (OTOA)*. All point mutations were rare in the Palestinian population (zero carriers in 288 unrelated controls); the carrier frequency of the *OTOA* genomic deletion was 1%. In six families, we identified five genomic regions likely to harbor novel genes for human hearing loss on chromosomes 1p13.3 (DFNB82), 9p23–p21.2/p13.3–q21.13 (DFNB83), 12q14.3–q21.2 (DFNB84; two families), 14q23.1–q31.1, and 17p12–q11.2 (DFNB85).

*European Journal of Human Genetics* (2010) 18, 407–413; doi:10.1038/ejhg.2009.190; published online 4 November 2009

**Keywords:** hearing; deafness; DFNB; homozygosity mapping; mutation; protein modeling

## INTRODUCTION

Recessively inherited phenotypes are common in the Palestinian population as the result of a high frequency of consanguineous marriages.<sup>1,2</sup> Palestinian families expressing such traits have cooperated with research studies to identify the critical genes for many phenotypes. In particular, Palestinian families with nonsyndromic hearing loss (NSHL) have been a valuable genetic resource for the identification of genes essential for normal hearing.<sup>3–8</sup> Homozygosity mapping with SNPs has proven an effective approach to the identification of these genes, because persons with a recessive phenotype and born of a consanguineous union are likely to be homozygous both for the causative mutation and for polymorphisms in the genomic region surrounding the mutation. In this study, we used high-density SNP arrays to genotype the DNA of 155 relatives from 20 consanguineous families with autosomal recessive NSHL. Deafness-associated homozygosity profiles revealed loci likely to harbor novel genes for NSHL in 6 families and homozygous mutations in known genes for NSHL in 14 families. In all families, the deafness-causing mutation was present in the longest homozygous segment of the deaf individuals' genomes.

## MATERIALS AND METHODS

### Subjects

Subjects with prelingual bilateral hearing loss ( $n=238$ ) were ascertained through schools for deaf children in the West Bank: Eftah School for the Deaf in Bethlehem, Al Amad School for the Deaf in Hebron, and the Princess Basma Rehabilitation Center for Deaf Children in Jerusalem. In addition, families were referred to the Hereditary Research Laboratory at Bethlehem University from the clinic of Dr Wael Salhab and from the Dar Al-Kalima Audiology Clinic in Bethlehem by Mr Michael Rahil. Families were contacted and informed consent was obtained from parents with assent from older children. The project was approved by the Human Subjects Committee of Bethlehem University and by the Human Subjects Division of the University of Washington. Pedigrees were constructed with 80% of families reporting consanguineous marriages. Audiological exams were performed to exclude hearing loss attributed to infection or trauma and to evaluate the severity and laterality of the hearing loss. Additional ascertainment was carried out for the extended families of 20 probands reporting affected relatives (Figure 1). For these 20 kindreds, all family members willing to participate were enrolled (Supplementary Table 1). Complete physical, visual, and vestibular exams were carried out to exclude syndromic deafness. Population controls comprised 288 Palestinian adults with normal hearing and came from two sources: (i) persons attending West Bank clinics for preventive health services ( $n=201$ ) and (ii) Palestinian individuals


<sup>1</sup>Department of Life Sciences, Bethlehem University, Bethlehem, Palestine; <sup>2</sup>Division of Medical Genetics, Department of Medicine, University of Washington, Seattle, WA, USA; <sup>3</sup>Department of Genome Sciences, University of Washington, Seattle, WA, USA; <sup>4</sup>Department of Human Molecular Genetics and Biochemistry, Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel

\*Correspondence: Dr T Walsh, Division of Medical Genetics, Department of Medicine, University of Washington, 1959 NE Pacific St, Box 357720, Seattle, WA 98195-7720, USA. Tel: +1 206 616 4296; Fax: +1 206 616 4295; E-mail: twalsh@uw.edu

<sup>5</sup>These authors contributed equally to this work.

<sup>6</sup>Current address: Department of Human Genetics, University of Michigan, Ann Arbor, MI 48109, USA

Received 22 June 2009; revised 23 September 2009; accepted 1 October 2009; published online 4 November 2009



**Figure 1** Inherited prelingual hearing loss in Palestinian families. Hearing loss in affected individuals (filled symbols) is prelingual with thresholds >90 dB. Double bars between parents indicate marriages within the same *hamula*, or clan. Only enrolled individuals and their parents are shown. Based on complete pedigree information, segregation of hearing loss is consistent with recessive inheritance in each family.

living in Israel who contributed DNA to the National Laboratory for the Genetics of Israeli Populations at Tel Aviv University ( $n=87$ ).

### Excluding known Palestinian deafness alleles

Genomic DNA was prepared by salting out.<sup>9</sup> Genomic DNA from 238 probands and 58 additional deaf relatives from 20 extended families were fully sequenced for both exons of connexin 26, as described.<sup>10</sup> In addition, all individuals were tested for deafness-associated alleles previously identified in families of Palestinian origin: *Otoancorin.1067A>T*, *SLC26A4.716T>A*, *SLC26A4.1001G>T*, *TMPRSS3.988delA*, *TRIOBP.1039C>T*, *TRIOBP.1741C>T*,<sup>8,11</sup> *Otoferlin.2416T>A*,<sup>3</sup> *TECTA.IVS9(+1)G>A*,<sup>4</sup> *Otoancorin.IVS12(+2)G>T*,<sup>5</sup> the  $\beta$ -satellite insertion of *TMPRSS3*,<sup>6</sup> and *Whirlin.2332C>T*.<sup>7</sup>

### Homozygosity mapping

Genomic DNA samples (250 ng) were hybridized on Affymetrix GeneChip (Santa Clara, CA, USA) Human Mapping 250K NspI arrays according to the manufacturer's protocol (<http://www.affymetrix.com>). Arrays were scanned using the Affymetrix Gene Chip Scanner 3000 with the Gene Chip Operating System 1.4 (GCOS). Data were analyzed using the Gene Chip DNA Analysis Software 2.0 (GDAS) and classified by chromosome and sorted on the basis of the physical position of the SNPs. Homozygous regions were defined as segments where SNPs were homozygous for one allele among affected relatives but discordant (heterozygous or homozygous of the complement allele) among unaffected relatives in the pedigree. Enhanced stringency requirements for SNP quality were applied, with only SNPs with a quality score better than 0.06 from GDAS and GCOS being retained, thus using about 67% of the SNPs in homozygosity analyses. A segment was considered continuously homozygous if there was no more than one heterozygous SNP in any 10-SNP window. Homozygous segments of 2 Mb or longer were retained for further analysis. LOD scores<sup>12</sup> were calculated for these homozygous segments for each family, exploiting genotypes of all informative SNPs. The LOD score for a family depends on the family's structure and parameters of the recessive model and is the same for all informative homozygous regions in the genome.

### Sequencing known genes

Previously described deafness genes<sup>13</sup> mapping within homozygous regions greater than 2 Mb were evaluated by Sanger DNA sequencing of PCR amplicons. Primers were designed to encompass all annotated exons and 50–100 bp of flanking intronic sequence.<sup>8</sup> Primers for *Cadherin23*, *ESRRB*, *Otoferlin*, *Pejvakin*, *Myosin 7A*, *Myosin 15A*, *SLC26A4*, *TECTA*, *TMCI*, *TMHS*, and *TMPRSS3* are available on request. For families C, AN, and CB, both genomic DNA and RNA from white cells were available; for the other families, only genomic DNA was available for analysis. As each new mutation was identified, the additional 218 deaf probands and 288 hearing controls were screened by PCR-RFLP and direct sequencing.

### Analysis of copy number variation

DNA samples from affected individuals were PCR amplified with unique STSs from the nonduplicated portion of the *Otoancorin* gene, exons 1–19 (chr16: 21 597 336–21 647 927). Primers are available on request. Genomic boundaries of the deletion in family DO and in Palestinian controls were determined by high-resolution arrayCGH on NimbleGen 2.1 million probe arrays. Genomic samples were analyzed<sup>14</sup> using NA15724 as the reference genome for hybridizations. In controls, genomic deletion was evaluated by SYBR Green real-time PCR. Briefly, 10 ng of genomic DNA was amplified with primer sets located in exon 6 of *Otoancorin* and in intron 12 of *Albumin* (which served as an endogenous control) on an Applied Biosystems (Foster City, CA, USA) 7300 Real-Time PCR Instrument. Samples were run in quadruplicate alongside a known 2-copy control and a known 1-copy carrier from family DO. Raw  $C_t$  values were collected and the relative fold difference between test samples and controls was determined using the  $2^{-\Delta\Delta C_t}$  method.<sup>15</sup> Samples with a 0.5-fold decrease in signal at *Otoancorin* relative to the known two-copy sample were analyzed by arrayCGH to determine break points.

## RESULTS

In this cohort of 286 families, hearing loss is attributed to connexin 26 mutations in 8% of families (23/286) and to other previously

described alleles of Palestinian origin in 9% of families (25/286). From the families of the unresolved 238 probands, we selected 20 informative kindreds in which parents reported consanguineous marriages and patterns of hearing loss were consistent with recessive inheritance (Figure 1). From these 20 kindreds, 78 individuals with NSHL, 28 of their hearing siblings, and 49 parents enrolled in the project (Supplementary Table 1).


Each participant provided a sample of whole blood from which DNA was extracted and genotyped on Affymetrix 250K SNP arrays. We generated homozygosity profiles specific to the deaf individuals in each family, defining a deafness-associated homozygous peak as a contiguous genomic region >2 Mb across which SNPs were homozygous for one allele among affected relatives but discordant (heterozygous or homozygous for the alternate allele) among unaffected parents or sibs. For nine families, a single homozygous peak was observed, in five families two peaks were observed, and in six families multiple peaks were observed (Figure 2).

In 16 of the 20 families, the longest homozygous region included a gene known to be associated with hearing loss (Table 1). For each family, the relevant gene was fully sequenced from genomic DNA of the proband, revealing in 14 families alleles that were very likely to be deleterious. These included premature truncating alleles of *Pejvakin*, *TECTA*, *TMPRSS3*, and *Otoferlin*, a splice site mutation leading to a large in-frame deletion of the *Pendrin/SLC26A4* message, and non-synonymous mutations in *MYO15A*, *MYO7A*, *CDH23*, and *TMHS*, which were further evaluated by splicing analysis and molecular modeling. Detailed analyses of these point mutations are presented in Supplementary Information. One family (DO) harbored a large genomic deletion of *Otoancorin*. For three families (DE, BG, C), homozygous regions harbored known deafness genes, but full sequence analysis of these candidate genes did not reveal any rare alleles segregating with hearing loss. These linkage regions likely harbor as yet undiscovered genes for hearing loss. Finally, in three other families (CG, DP, CN), hearing loss mapped to homozygous regions that did not include known genes for human hearing loss, thereby also likely reflecting new recessive deafness loci in the Palestinian population. In summary, six of the 20 families are likely to represent five novel genes for NSHL.

### Homozygosity for a genomic deletion of *Otoancorin* at 16p12

The largest homozygous peak in family DO was a 5.4 Mb region of chromosome 16p12.1–p12.2, including the gene *Otoancorin*. The genomic sequence flanking *Otoancorin* has multiple clusters of high-identity segmental duplications, which likely gave rise to a deletion spanning the *Otoancorin* gene through the nonallelic homologous recombination (Figure 3).<sup>16</sup> The largest *Otoancorin* isoform comprises 28 exons. Exons 1–19 map to a single location in the human genome, but exons 20–28 are located in a 68 kb region that is segmentally duplicated, at greater than 99% identity, at chr16: 22 464 054–22 532 760, approximately 820 kb centromeric of the original locus.

From the genomic DNA of DO2, DO3, DO4, and DO5 (all of whom are deaf), PCR amplification of *Otoancorin* exons 1–19 failed to produce a product (Figures 3a and b). In contrast, amplification of these exons from genomic DNA of DO1 and DO6 yielded products of normal size. This pattern suggests that the deaf individuals are homozygous for a genomic deletion spanning these exons. DNA from the same individuals was further analyzed by high-density genome-wide arrayCGH, revealing a deletion of approximately 500 kb that completely deletes the *Otoancorin* gene (Figure 3c). Deaf individuals in family DO were homozygous for the deletion and their parents heterozygous for the deletion. The deletion is flanked



**Figure 2** Genomic regions of homozygosity associated with hearing loss. Each colored bar represents a genomic region homozygous for the same haplotype in all deaf individuals in a family, and heterozygous or homozygous for the alternate allele among unaffected parents or siblings in the same family. Chromosome arms are indicated on X-axes; lengths in Mb of the homozygous regions are indicated on Y-axes. Green bars represent homozygous regions with a deleterious mutation in a known deafness gene, as indicated in Table 1. Red bars represent homozygous regions likely to harbor novel deafness genes as indicated in Table 2. Blue bars indicate other regions of homozygosity.

**Table 1 Genomic regions homozygous in deaf individuals revealing mutations in known genes for hearing loss**

Kindred	Peaks > 2 Mb	Longest homozygous region		DFN gene in region			Homozygous mutation in family		Frequency of mutation in the Palestinian population	
		Location NCBI	build 36.1 (hg18)	DFN gene in region	Nucleotide change	Amino-acid change	Heterozygotes in 288 controls	Homozygotes in 218 unrelated deaf		
AM	5	19.4	chr2: 14,733,739–34,121,329	OTOF	c4157 C>T	Arg 577 STOP	0	0		
A	1	1.9	chr2: 177,555,791–179,471,954	PJKV	c762 C>T	Arg 136 STOP	0	1		
DD	2	14.6	chr6: 23,791,422–38,418,869	TMHS	c1 A>G	Met 1 Val	0	0		
CB	5	5.2	chr7: 105,307,599–110,520,953	SLC26A4	c1341 ΔG	Δ aa 422–447	0	0		
G	1	19.8	chr10: 53,479,033–73,231,404	CDH23	c1427 C>T	Pro 346 Ser	0	0		
DA	3	19.2	chr10: 71,683,886–90,894,091	CDH23	c1428 C>T	Pro 346 Leu	0	0		
AB	1	11.7	chr10: 72,774,642–84,459,990	CDH23	c2065 C>T	Pro 559 Ser	0	1		
J	3	26.4	chr11: 69,115,451–95,560,647	MYO7A	c6487 G>A	Gly 2163 Ser	0	2		
BQ	1	11.4	chr11: 110,396,930–121,744,055	TECTA	c4857 C>A	Cys 1619 STOP	0	See BI		
BI	4	5.9	chr11: 117,111,049–123,047,230	TECTA	c4857 C>A	Cys 1619 STOP	0	2		
DO	2	5.4	chr16: 21,035,675–26,477,060	OTOA	Complete Δ	Complete Δ	3	0		
P	1	10.1	chr17: 16,806,979–26,862,502	MYO15A	c7545 G>T	Asp 2403 fsX2414	0	See AN		
AN	1	13.6	chr17: 17,292,606–30,871,427	MYO15A	c7545 G>T	Asp 2403 fsX2414	0	5		
IB	2	5.4	chr21: 40,357,695–45,755,537	TPR53	c783T>A	Cys 194 STOP	0	0		

by segmental duplications located at chr16: 21 314 758–21 419 562 and chr16: 21 746 863–21 851 057. The identity between these two loci is approximately 99.7%, suggesting that the nonallelic homologous recombination is the likely mechanism of deletion, as has been described for other genomic regions.<sup>16</sup> The deletion was observed as a heterozygote in 1.0% of unrelated Palestinian controls (3/288). A comparison of the arrayCGH intensity values suggests that the mutations in family DO and in the controls have the same genomic breakpoints (Figure 3c).

### Novel deafness loci


In six families, homozygosity mapping suggests that hearing loss is due to mutations in as-yet-unidentified genes. The six families represent five new loci (ie, two families map to the same region). We identified the critical regions of linkage, calculated LOD scores for each of the six families, and reserved DFNB loci (HUGO Gene Nomenclature Committee) for families with LOD scores greater than 3.0 (Table 2).

**DFNB82.** Hearing loss in family CG maps to a 3.1 Mb region of chromosome 1p13.3 with an LOD score of 5.16 (Supplementary Figure 3a). The DFNB82 region overlaps the DFNB32 locus by 833 kb.<sup>17</sup> DFNB32 was described in a Tunisian family with profound congenital hearing loss; the critical gene is not yet known. In DNA from family CG, we sequenced the two genes in the region of overlap (VAV3 and SLC25A24), but found no novel variants segregating with NSHL. It is therefore likely that DFNB82 and DFNB32 reflect different genes for NSHL.

**DFNB83.** Hearing loss in family DE maps to two regions of chromosome 9: a 9p region of 16.5 Mb and a pericentromeric region of 28.5 Mb (excluding the unannotated 15 Mb gap at the centromere; supplementary Figure 3b). The LOD score is 3.07. The chromosome 9 pericentromeric region includes the NSHL gene *TMCI*.<sup>18</sup> The 9p region includes the *MTAP* gene, which is disrupted by a *de novo* balanced translocation in a child with syndromic deafness.<sup>19</sup> We sequenced all exons and conserved intronic regions of *TMCI* and *MTAP* in family DE, but did not identify novel variants segregating with deafness in either gene. The DFNA47 locus maps within the DFNB83 region at chr9:13 046 167–21 980 675.<sup>20</sup> The gene responsible for DFNA47 is not known. It is possible that mutations of the same gene will be responsible for recessive and dominant hearing loss.

**DFNB84.** Deaf individuals in families CN and DP have overlapping regions of homozygosity and identical haplotypes over a 12.5 Mb region of chromosome 12q (chr12: 74 498 486–86 977 739) with an LOD score of 3.45 (supplementary Figure 3c). No genes for inherited hearing loss have been described in this genomic interval and it therefore represents a novel locus for recessive NSHL.

**DFNB85.** Family C is a large kindred with 42 individuals enrolled. Hearing loss in family C maps to an 11.1 Mb region of chromosome 17p12–q11.2 with an LOD score of 7.25 (supplementary Figure 3d). *MYO15A* lies in this region and was therefore an excellent candidate for the NSHL gene in this family. However, full sequencing of *MYO15A* from genomic DNA and lymphoblast cDNA from homozygous affected persons and heterozygous obligate carrier relatives revealed no functional mutations. Furthermore, the two alleles of *MYO15A* were expressed at the same levels in lymphoblast cDNA from obligate carriers, as determined by the evaluation of cSNP rs854772 in exon 40. Although it is possible that *MYO15A* may harbor a cryptic



**Figure 3** Genomic deletion of *Otoancorin* (*OTOA*). **(a)** Family DO pedigree and PCR amplification of a unique STS in *OTOA* intron 17 at chr16: 21 645 309. The STS is present in the unaffected mother and sibling and absent from all affected siblings. **(b)** Presence (+) or absence (-) of multiple STSs within and flanking *OTOA* in genomic DNA from unaffected mother DO1 and deaf child DO5. **(c)** High-resolution arrayCGH of 16p12.2 (chr16: 21 200 000–22 000 000) in genomic DNA from Palestinian control individuals HGDP4921 and HGDP5043 and from family DO unaffected mother DO1 and deaf child DO5. Deviations from zero of  $\log_2$  ratios of probe intensities are depicted by vertical lines, with ratios > 1.5 SDs from the mean ratio colored red and green to represent relative losses and gains, respectively. Segmental duplications of increasing similarity (90–98, 98–99, and >99%) are represented by gray, yellow, and orange bars, respectively. The size of the *OTOA* locus (black bar) is 82 kb; the size of the deletion is 320–550 kb. In the Palestinian hearing population, the carrier frequency of the deletion is 1% (3/288 controls).

**Table 2** Chromosomal regions of new loci for nonsyndromic hearing loss

DFNB	Kindred		Chromosome		Size (Mb)	Flanking markers		LOD score
			Start (hg18)	End (hg 18)				
82	CG	1	108 108 849	111 251 931	3.1	rs17542571	rs1936942	5.16
83	DE	9	9 959 023	26 430 090	16.5	rs4742645	rs1571364	3.07
		9	34 401 026	77 904 900	28.5	rs10738927	rs12001902	
84	CN/DP	12	74 498 486	86 977 739	12.5	rs10879987	rs2471512	3.45
85	C	17	15 212 798	26 490 848	11.1	rs230884	rs12603885	7.25
-	BG	14	57 474 255	80 664 133	23.2	rs41514351	rs3873938	

Sizes of linked regions exclude gaps at centromeres.

mutation in a regulatory element driving cochlear-specific expression, clustering of functionally related genes suggests that it is likely that a second gene for NSHL is among the 165 other genes in this region.

**Chromosome 14q locus.** Homozygosity mapping of family BG revealed a single 23 Mb region of homozygosity on chromosome

14a (Supplementary Figure 3e). This region includes the gene *ESRRB* responsible for DFNB35.<sup>21</sup> Full sequencing of *ESRRB* in genomic DNA from affected relatives and carriers did not reveal any novel variants in the coding region or flanking regulatory regions. We hope to recruit additional informative relatives from family BG in an effort to reduce the size of the homozygous interval.

## DISCUSSION

For the analysis of recessively inherited hearing loss in these consanguineous families, homozygosity mapping with high-density SNP arrays yielded highly informative homozygosity profiles. In 14 families, the critical mutation was identified by sequencing a single gene mapping within the largest homozygous region. Even in small families with few deaf individuals (eg, families AM and CB), each with multiple homozygous peaks, the critical allele was still identifiable by searching exclusively in the largest region.

None of the ten newly described alleles explain a high proportion of hearing loss among Palestinian families. In the Palestinian hearing population, heterozygous carriers were observed for only one allele, the large genomic deletion of *OTOA*. This region of chromosome 16p12.2 represents a 'hot spot' for genomic deletion because of its location between two identical segmental duplications. This specific deletion and the reciprocal duplication have been observed in 0.5–1.0% of European controls.<sup>16</sup> It has recently been shown that *Stereocilin* (DFNB15) is also subject to genomic rearrangements mediated by segmental duplications at chromosome 15q15.3. A homozygous deletion of 90 kb, including *STRC* and four other genes, is responsible for a syndromic form of deafness with male infertility.<sup>22,23</sup> The carrier frequency of the *STRC* deletion has been estimated at 1.6%. It is possible that homozygous deletions of *OTOA* and *STRC* will be enriched in deaf individuals from other consanguineous communities.

The six families in which NSHL maps to five new loci represent an excellent opportunity to identify novel genes essential for normal hearing. The genomic regions harboring these genes are large (generally >10 Mb), but the critical gene can be identified readily with next generation sequencing approaches. Consanguineous kindreds, even small families, prove extremely valuable for the identification of novel alleles and novel loci. Of the families evaluated in this project, 30% are likely to reveal additional genes for hearing loss. This proportion suggests that a large number of genes critical to hearing remain to be discovered.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## ACKNOWLEDGEMENTS

We thank the families for their generous participation in this study. We thank Dr Wael Salhab for a clinical evaluation of the patients, and Michael Rahil of the Dar Al-Kalima Audiology Clinic for audiological analyses. We thank the technical staff of the Hereditary Research Laboratory (Bethlehem University), Rob Hall at the Center for Array Technology (University of Washington), Gregory Cooper, and Heather Mefford for helpful comments and suggestions. This study was supported by the NIH/NIDCD Grant R01DC005641.

- Jaber L, Halpern GJ, Shohat T: Trends in the frequencies of consanguineous marriages in the Israeli Arab community. *Clin Genet* 2000; **58**: 106–110.
- Zlotogora J, Barges S, Bisharat B, Shalev SA: Genetic disorders among Palestinian Arabs. 4: Genetic clinics in the community. *Am J Med Genet A* 2006; **140**: 1644–1646.
- Yasunaga S, Grati M, Cohen-Salmon M *et al*: A mutation in *OTOF*, encoding otoferlin, a FER-1-like protein, causes DFNB9, a nonsyndromic form of deafness. *Nat Genet* 1999; **21**: 363–369.
- Mustapha M, Weil D, Chardenoux S *et al*: An alpha-tectorin gene defect causes a newly identified autosomal recessive form of sensorineural pre-lingual non-syndromic deafness, DFNB21. *Hum Mol Genet* 1999; **8**: 409–412.
- Zwaenepoel I, Mustapha M, Leibovici M *et al*: Otoancorin, an inner ear protein restricted to the interface between the apical surface of sensory epithelia and their overlying acellular gels, is defective in autosomal recessive deafness DFNB22. *Proc Natl Acad Sci USA* 2002; **99**: 6240–6245.
- Scott HS, Kudoh J, Wattenhofer M *et al*: Insertion of beta-satellite repeats identifies a transmembrane protease causing both congenital and childhood onset autosomal recessive deafness. *Nat Genet* 2001; **27**: 59–63.
- Mburu P, Mustapha M, Varela A *et al*: Defects in whirlin, a PDZ domain molecule involved in stereocilia elongation, cause deafness in the whirler mouse and families with DFNB31. *Nat Genet* 2003; **34**: 421–428.
- Shahin H, Walsh T, Sobe T *et al*: Mutations in a novel isoform of TRIOBP that encodes a filamentous-actin binding protein are responsible for DFNB28 recessive non-syndromic hearing loss. *Am J Hum Genet* 2006; **78**: 144–152.
- Miller SA, Dykes DD, Polesky HF: A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 1988; **16**: 1215.
- Shahin H, Walsh T, Sobe T *et al*: Genetics of congenital deafness in the Palestinian population: multiple connexin 26 alleles with shared origins in the Middle East. *Hum Genet* 2002; **110**: 284–289.
- Walsh T, Abu Rayan A, Abu Sa'ed J *et al*: Genomic analysis of a heterogeneous Mendelian phenotype: multiple novel alleles for inherited hearing loss in the Palestinian population. *Hum Genomics* 2006; **2**: 203–211.
- Ott J: *Analysis of Human Genetic Linkage*, 3rd edn., Johns Hopkins Univ Press: Baltimore.
- Hereditary Hearing Loss Homepage. (<http://webh01.ua.ac.be/hhh/>).
- Selzer RR, Richmond TA, Pofahl NJ *et al*: Analysis of chromosome breakpoints in neuroblastoma at sub-kilobase resolution using fine-tiling oligonucleotide array CGH. *Genes Chromosomes Cancer* 2005; **44**: 305–319.
- Livak KJ, Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-ΔΔCT</sup> Methods. *Methods* 2001; **25**: 402–408.
- Itsara A, Cooper GM, Baker C *et al*: Population analysis of large copy number variants and hotspots of human genetic disease. *Am J Hum Genet* 2009; **84**: 148–161.
- Masmoudi S, Tlili A, Majava M *et al*: Mapping of a new autosomal recessive nonsyndromic hearing loss locus (DFNB32) to chromosome 1p13.3–22.1. *Eur J Hum Genet* 2003; **11**: 185–188.
- Kurima K, Peters LM, Yang Y *et al*: Dominant and recessive deafness caused by mutations of a novel gene, *TMC1*, required for cochlear hair-cell function. *Nat Genet* 2002; **30**: 277–284.
- Williamson RE, Darrow KN, Michaud S *et al*: Methylthioadenosine phosphorylase (MTAP) in hearing: Gene disruption by chromosomal rearrangement in a hearing impaired individual and model organism analysis. *Am J Med Genet A* 2007; **143A**: 1630–1639.
- D'Adamo P, Donaudy F, D'Eustacchio A, Di Iorio E, Melchionda S, Gasparini P: A new locus (DFNA47) for autosomal dominant non-syndromic inherited hearing loss maps to 9p21–22 in a large Italian family. *Eur J Hum Genet* 2003; **11**: 121–124.
- Collin RW, Kalay E, Tariq M *et al*: Mutations of *ESRRB* encoding estrogen-related receptor beta cause autosomal-recessive nonsyndromic hearing impairment DFNB35. *Am J Hum Genet* 2008; **82**: 125–138.
- Zhang Y, Malekpour M, Al-Madani N *et al*: Sensorineural deafness and male infertility: a contiguous gene deletion syndrome. *J Med Genet* 2007; **44**: 233–240.
- Knijnenburg J, Oberstein SA, Frei K *et al*: A homozygous deletion of a normal variation locus in a patient with hearing loss from non-consanguineous parents. *J Med Genet* 2009; **46**: 412–417.

Supplementary Information accompanies the paper on European Journal of Human Genetics website (<http://www.nature.com/ejhg>)