

Interlayer Registry Determines the Sliding Potential of Layered Metal Dichalcogenides: The case of 2H-MoS₂

Adi Blumberg, Uri Keshet, Inbal Zaltsman, and Oded Hod

Department of Chemical Physics, School of Chemistry, the Sackler Faculty of Exact Sciences,
Tel-Aviv University, Tel-Aviv 69978, Israel

Supporting Information: Coordinates of some representative structures

2H-MoS₂ is a layered crystal, where each layer consists of three sub-layers of S-Mo-S (Fig. S1). The crystal unit cell has the symmetry of the P6₃/mmc space group, with lattice parameters of $a = b = 3.1612 \text{ \AA}$ and $c = 12.2985 \text{ \AA}$.^{1-2,3} For the construction of the bilayer system studied in the present manuscript we used the following partial coordinates representation:⁴⁻⁵

$$\text{Mo: } (1/3)A + (2/3)B + (1/4)C$$

$$\text{S: } (2/3)A + (1/3)B - (1/2-u)C$$

$$\text{S: } (2/3)A + (1/3)B + (1-u)C$$

$$\text{Mo: } (2/3)A + (1/3)B + (3/4)C$$

$$\text{S: } (1/3)A + (2/3)B + uC$$

$$\text{S: } (1/3)A + (2/3)B + (1.5-u)C$$

$$\text{Where: } A = \left(-\sqrt{3} \frac{a}{2}, \frac{a}{2}, 0 \right), B = \left(\sqrt{3} \frac{b}{2}, \frac{b}{2}, 0 \right), C = (0, 0, c), u = 0.621$$




Fig. S1 Different views of MoS₂ crystal structure: (a) X-Y plain, (b) Y-Z plain and (c) X-Z plain.

The resulting coordinates of some high symmetry configurations (see Fig. 3 of the main text) are given below:

AA' configuration:

S	(-0.912560,	4.741800,	1.488119)
Mo	(0.912560,	4.741800,	3.074625)
S	(-0.912560,	4.741800,	4.661132)
S	(0.912560,	4.741800,	7.637369)
Mo	(-0.912560,	4.741800,	9.223875)
S	(0.912560,	4.741800,	10.810382)




Fig. S2 (a) AA'

AB₁ configuration:

S	(-0.912560,	4.741800,	1.488119)
Mo	(0.912560,	4.741800,	3.074625)
S	(-0.912560,	4.741800,	4.661132)
S	(2.737680,	4.741800,	7.637369)
Mo	(0.912560,	4.741800,	9.223875)
S	(2.737680,	4.741800,	10.810382)




Fig. S2 (b) AB₁

AB₂ configuration:

S	(-0.912560,	1.580600,	1.488119)
Mo	(0.912560,	1.580600,	3.074625)
S	(-0.912560,	1.580600,	4.661132)
S	(1.837043,	6.321606,	7.637369)
Mo	(0.011923,	6.321606,	9.223875)
S	(1.837043,	6.321606,	10.810382)




Fig. S2 (c) AB₂

- (1) Hu, J. J.; Zabinski, J. S.; Bultman, J. E.; Sanders, J. H.; Voevodin, A. A. *Tribol Lett* **2006**, *24*, 127.
- (2) Castro-Guerrero, C. F.; Deepak, F. L.; Ponce, A.; Cruz-Reyes, J.; Valle-Granados, M. D.; Fuentes-Moyado, S.; Galvan, D. H.; Jose-Yacaman, M. *Catalysis Science & Technology* **2011**, *1*, 1024.
- (3) Entry 37-1492, Joint Committee on Powder Diffraction Standards. International Centre for Diffraction Data, 1601 Park Lane, Swarthmore, PA 19081-2389, USA.
- (4) Bronsema, K. D.; De Boer, J. L.; Jellinek, F. *Zeitschrift für anorganische und allgemeine Chemie* **1986**, *540*, 15.
- (5) Dickinson, R. G.; Pauling, L. *Journal of the American Chemical Society* **1923**, *45*, 1466.