

Weiszfeld's Method: Old and New Results

Amir Beck · Shoham Sabach

Received: 25 February 2014 / Accepted: 9 May 2014 / Published online: 31 May 2014
© Springer Science+Business Media New York 2014

Abstract In 1937, the 16-years-old Hungarian mathematician Endre Weiszfeld, in a seminal paper, devised a method for solving the Fermat–Weber location problem—a problem whose origins can be traced back to the seventeenth century. Weiszfeld's method stirred up an enormous amount of research in the optimization and location communities, and is also being discussed and used till these days. In this paper, we review both the past and the ongoing research on Weiszfeld's method. The existing results are presented in a self-contained and concise manner—some are derived by new and simplified techniques. We also establish two new results using modern tools of optimization. First, we establish a non-asymptotic sublinear rate of convergence of Weiszfeld's method, and second, using an exact smoothing technique, we present a modification of the method with a proven better rate of convergence.

Keywords Complexity analysis · Fermat–Weber problem · Gradient method · Localization theory · Weiszfeld's method

Mathematics Subject Classification 49-02 · 90C25 · 90C30 · 90-02 · 65K05 · 97N60

A. Beck (✉)
Department of Industrial Engineering and Management,
Technion—Israel Institute of Technology, 32000 Haifa, Israel
e-mail: becka@ie.technion.ac.il

S. Sabach
School of Mathematical Sciences, Tel Aviv University,
69978 Tel Aviv, Israel
e-mail: ssabach@gmail.com

1 Introduction

One of the most fundamental location problems is the so-called Fermat–Weber problem, which consists of finding a point that minimizes the sum of its weighted distances to a given finite set of anchor points. The problem is credited to the well-known French mathematician Pierre de Fermat, who at the beginning of the seventeenth century posed the following question:


Given three points in a plane, find a fourth point such that the sum of its distances to the three given points is as small as possible.

The Italian physicist and mathematician Evangelista Torricelli (mostly known for inventing the barometer) found a construction method of this point by ruler and compass, and it is therefore also called “the Toricelli point”; see Fig. 1 for an illustration of Torricelli’s construction for the case where the triangle has all angles less than 120° .

At the beginning of the twentieth century, the German economist Alfred Weber incorporated weights, and was able to treat facility location problems with more than 3 facilities, and the problem was consequently called “the Fermat–Weber problem.” Other names for the problem are “the Fermat problem,” “the Weber problem,” “the Fermat–Torricelli problem,” “the Steiner problem,” and many more variants. More details on the history of the Fermat–Weber problem can be found, for example, in [1], as well as in the survey papers [2] and the second part of the book [3]. More information on the geometric aspects of the Fermat–Weber problem as well as variations and open problems can be found in [4].

The purpose of this paper is *not* to present another review on the Fermat–Weber problem but rather to focus on a very simple algorithm designed for solving it, suggested in 1937 by the 16-years- old Hungarian mathematician Endre Vaszonyi

Fig. 1 Given a (proper) triangle, formed by three points a_1 , a_2 , and a_3 , construct three equilateral triangles such that each contains one of the edges from the triangle $a_1 a_2 a_3$. Then, circumscribe each equilateral triangle. The unique point of intersection of these three circles is the point, that yields the minimum distance to the points a_1 , a_2 , and a_3 ; it is called “the Torricelli Point” and denoted by x^*



Weiszfeld [5]. Quite interestingly, as indicated by Weiszfeld himself in [6], the focus of Weiszfeld's paper was *not* to design an algorithm for solving the Fermat–Weber problem but rather to prove a mathematical theorem, and in fact the all notion of an “algorithm” was unfamiliar to him, as he himself indicated in [6]: “*the word algorithm was unknown to me and to most mathematicians.*” The theorem itself was not new and was already established by Sturm [7] in 1884. Weiszfeld's paper provided three different proofs for this theorem and, in the first proof, he defined a sequence that was supposed to converge to the optimal solution of the Fermat–Weber problem.

Weiszfeld's method stirred up an enormous amount of research and had an impact on researchers from the optimization, as well as the location fields. The contribution of this paper is threefold. First, we will review the intriguing story of the algorithm ever since its derivation in 1937 until today and present the current status of convergence analysis. The presentation is self-contained, so the main convergence results will be presented with proofs—some of them are new and simplified. Our second contribution is to provide—using modern tools of optimization—a nonasymptotic sublinear rate of convergence analysis of the method. Finally, noting that Weiszfeld's method is essentially a gradient method, our third contribution will be to present an accelerated version of the method, based on a combination of an exact smoothing technique and an optimal gradient method; the resulting method is shown to have an improved rate of convergence.

The paper is organized as follows. The next section is devoted to the description of the problem and of Weiszfeld's method. In Sect. 3, we present the original paper of Weiszfeld and describe what was actually proven in that seminal paper (monotonicity of the sequence of function values), as well as pinpoint the critical mistake in the analysis. The method remained mainly unknown until 1962; in Sect. 4, we describe several papers that reinvented the method as well as tell the story of how the original Weiszfeld's paper was discovered by Harold Kuhn. In Sect. 5 we discuss the paper of Kuhn from 1973; we provide a different and simplified proof of its convergence theorem without requiring the usual assumption of noncollinearity of the anchors. We also present Kuhn's example on why the method can potentially “get stuck” at nonoptimal anchor points and recall his incorrect statement: the number of “bad” starting points (those leading to anchor points) is denumerable. Beginning from Kuhn's statement in 1973 on the number of “bad” starting points, we present in Sect. 6 the attempts of resolving this issue until its final closure in 2002. Several modifications of Weiszfeld's method, in which the only difference is the way the method operates on anchor points, are presented in Sect. 7, along with a method to pick the starting point in a way that ensures avoiding anchor points; a review of more elaborate modifications of the method concludes the section. Section 8 uses the simplified analysis of the previous sections, as well as modern analysis of gradient-based methods, to establish a nonasymptotic sublinear rate of convergence of the sequence of function values generated by Weiszfeld's method. In Sect. 9, we develop an accelerated version of the method, which is based on a combination of an exact smoothing technique, and the employment of an optimal gradient method. The new method is an accelerated version with an improved rate of convergence. The paper ends in Sect. 10, where the impact of Weiszfeld's method on other problems,

different than the Fermat–Weber problem, is explored. For the convenience of the reader, Appendix A contains a list of the most frequent notations used throughout the paper.

2 Problem Formulation and Weiszfeld’s Method

The Fermat–Weber problem, described verbally at the beginning of the paper, can be formulated mathematically as the problem of seeking $\mathbf{x} \in \mathbb{R}^d$ that solves

$$\min_{\mathbf{x}} \left\{ f(\mathbf{x}) = \sum_{i=1}^m \omega_i \|\mathbf{x} - \mathbf{a}_i\| \right\}, \quad (\text{FW})$$

where $\omega_i > 0, i = 1, 2, \dots, m$, are given *weights* and the vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m \in \mathbb{R}^d$ are given *anchors*.

To understand the result that Weiszfeld aimed to prove, let us first write down the expression of the gradient of the objective function of problem (FW):

$$\nabla f(\mathbf{x}) = \sum_{i=1}^m \omega_i \frac{\mathbf{x} - \mathbf{a}_i}{\|\mathbf{x} - \mathbf{a}_i\|}, \quad \mathbf{x} \notin \mathcal{A},$$

where $\mathcal{A} = \{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m\}$ denotes the set of anchors. Note that the gradient is only defined on points different from the anchors. The theorem that Weiszfeld re-established was the following (see [5,7]).

Theorem 2.1 (Weiszfeld’s original result) *Suppose that the anchors are not collinear. Then,*

- (a) *Problem (FW) has a unique optimal solution.*
- (b) *Let \mathbf{x}^* be the optimal solution of problem (FW). If $\mathbf{x}^* \notin \mathcal{A}$, then*

$$\nabla f(\mathbf{x}^*) = \sum_{i=1}^m \omega_i \frac{\mathbf{x}^* - \mathbf{a}_i}{\|\mathbf{x}^* - \mathbf{a}_i\|} = \mathbf{0}. \quad (1)$$

If $\mathbf{x}^ = \mathbf{a}_i$, for some $i \in \{1, 2, \dots, m\}$, then the following inequality holds:*

$$\left\| \sum_{j=1, j \neq i}^m \omega_j \frac{\mathbf{x}^* - \mathbf{a}_j}{\|\mathbf{x}^* - \mathbf{a}_j\|} \right\| \leq \omega_i. \quad (2)$$

As was noted in the introduction, the theorem was not new and was already established by Sturm [7] in 1884. Some comments about this theorem are required. The anchors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$ are said to be *collinear* iff they reside on the same line, *i.e.*, there exist $\mathbf{y}, \mathbf{d} \in \mathbb{R}^d$ and $t_1, t_2, \dots, t_m \in \mathbb{R}$ such that $\mathbf{a}_i = \mathbf{y} + t_i \mathbf{d}, i = 1, 2, \dots, m$. In the collinear case, it can be shown that the optimal solution of problem (FW) is a median of the anchors, meaning that the optimal solution is attained at (at least) one of the

anchor points; see [8] for further details. It is important to note that Weiszfeld did not analyze the weighted problem, but rather assumed that all the weights are equal to 1. We present the method and results in the weighted case, whose analysis is almost identical. Nowadays, this theorem seems like an elementary result and is a direct consequence of basic convex analysis. Indeed, when the anchors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$ are not collinear, the objective function f is strictly convex, and thus the optimal solution is unique [9, Theorem 3.4.2]. The optimality condition (1) is just the necessary and sufficient optimality condition $\nabla f(\mathbf{x}^*) = \mathbf{0}$ for unconstrained convex minimization problems at points, for which the objective function is differentiable, and the condition (2) is the necessary and sufficient optimality condition $\mathbf{0} \in \partial f(\mathbf{a}_i)$ at points of non-differentiability. These optimality conditions are satisfied also in the collinear case. More on the proof of the theorem, as well as extensions to more general settings, can be found in the paper [10] and in Chapter II of [3].

To present the method, assume that the anchors are not collinear and that \mathbf{x}^* is the unique optimal solution. We begin by writing explicitly the optimality condition $\nabla f(\mathbf{x}^*) = \mathbf{0}$ under the assumption that $\mathbf{x}^* \notin \mathcal{A}$:

$$\nabla f(\mathbf{x}^*) = \sum_{i=1}^m \omega_i \frac{\mathbf{x}^* - \mathbf{a}_i}{\|\mathbf{x}^* - \mathbf{a}_i\|} = \mathbf{0}.$$

The next step is to “extract” \mathbf{x}^* (disregarding the dependency in $\|\mathbf{x}^* - \mathbf{a}_i\|$ from \mathbf{x}^* , $i = 1, 2, \dots, m$) and to obtain the relation

$$\mathbf{x}^* = \frac{1}{\sum_{i=1}^m \frac{\omega_i}{\|\mathbf{x}^* - \mathbf{a}_i\|}} \sum_{i=1}^m \frac{\omega_i \mathbf{a}_i}{\|\mathbf{x}^* - \mathbf{a}_i\|} \tag{3}$$

or

$$\mathbf{x}^* = T(\mathbf{x}^*),$$

where the operator $T : \mathbb{R}^d \setminus \mathcal{A} \rightarrow \mathbb{R}^d$ is defined by

$$T(\mathbf{x}) := \frac{1}{\sum_{i=1}^m \frac{\omega_i}{\|\mathbf{x} - \mathbf{a}_i\|}} \sum_{i=1}^m \frac{\omega_i \mathbf{a}_i}{\|\mathbf{x} - \mathbf{a}_i\|}. \tag{4}$$

We have thus shown that, for any $\mathbf{y} \in \mathbb{R}^d \setminus \mathcal{A}$,

$$\mathbf{y} = T(\mathbf{y}) \text{ if and only if } \nabla f(\mathbf{y}) = \mathbf{0}. \tag{5}$$

Weiszfeld’s method is just a fixed point method for solving the relation (3).

Weiszfeld’s Method

Initialization. $\mathbf{x}_0 \in \mathbb{R}^d \setminus \mathcal{A}$.

General Step ($k = 0, 1, \dots$)

$$\mathbf{x}_{k+1} = T(\mathbf{x}_k). \tag{6}$$

3 The Original Paper of Weiszfeld

Weiszfeld's paper [5] was originally written in French; an English translation of the paper can be found in the recent paper [11]. The translation made by Frank Plastria contains, in addition, many interesting comments and observations. Taking a close look at the algorithm, one apparent fault is the fact that it is actually not well defined. Weiszfeld assumed that the initial vector \mathbf{x}_0 is different from any anchor point, that is, $\mathbf{x}_0 \notin \mathcal{A}$. However, this is not enough to ensure that the sequence $\{\mathbf{x}_k\}_{k \geq 0}$ generated by the method is well defined, since it might happen that a certain iterate \mathbf{x}_k will belong to the anchor set \mathcal{A} , resulting with a division by zero in the computation of the next iterate \mathbf{x}_{k+1} . This situation can occur even if the optimal solution does not belong to \mathcal{A} . This error was recognized later by Kuhn and Kuenne [12] in 1962 (see also the discussion in Sect. 4).

Putting aside the issue of “getting stuck” at nonoptimal anchor points, Weiszfeld was able to prove the monotonicity of the sequence of function values. We will repeat his arguments but will use the notation used by Beck and Teboulle in [13] that will serve us later on in the new analysis of the method. We begin by providing a different presentation of Weiszfeld's method. Define the auxiliary function $h : \mathbb{R}^d \times \mathbb{R}^d \setminus \mathcal{A} \rightarrow \mathbb{R}$ by

$$h(\mathbf{x}, \mathbf{y}) := \sum_{i=1}^m \omega_i \frac{\|\mathbf{x} - \mathbf{a}_i\|^2}{\|\mathbf{y} - \mathbf{a}_i\|}, \quad \mathbf{x} \in \mathbb{R}^d, \mathbf{y} \in \mathbb{R}^d \setminus \mathcal{A}. \quad (7)$$

Given an iteration \mathbf{x}_k , it is not difficult to show that next iterate $\mathbf{x}_{k+1} = T(\mathbf{x}_k)$ is determined as the minimizer of the function

$$s_k(\mathbf{x}) = h(\mathbf{x}, \mathbf{x}_k) = \sum_{i=1}^m \omega_i \frac{\|\mathbf{x} - \mathbf{a}_i\|^2}{\|\mathbf{x}_k - \mathbf{a}_i\|}$$

over \mathbb{R}^d . Indeed, $s_k(\cdot)$ is a strongly convex function, and its unique minimizer is determined by the optimality condition

$$\nabla s_k(\mathbf{x}) = \mathbf{0}.$$

That is,

$$\sum_{i=1}^m \omega_i \frac{\mathbf{x} - \mathbf{a}_i}{\|\mathbf{x}_k - \mathbf{a}_i\|} = \mathbf{0},$$

which, after some simple algebraic manipulation, can be seen to be equivalent to the relation $\mathbf{x} = T(\mathbf{x}_k)$. In other words, what we have shown is that, for any $\mathbf{y} \in \mathbb{R}^d \setminus \mathcal{A}$,

$$T(\mathbf{y}) = \operatorname{argmin}_{\mathbf{x} \in \mathbb{R}^d} h(\mathbf{x}, \mathbf{y}).$$

We will now recall a technical lemma containing the latter property along with several other properties connecting the function f and the auxiliary function h , which will

be the key in proving the monotonicity of the sequence of function values (see [13, Lemma 1.1]).

Lemma 3.1 (Properties of the auxiliary function h) *The following properties hold.*

(i) For any $\mathbf{y} \in \mathbb{R}^d \setminus \mathcal{A}$,

$$h(\mathbf{y}, \mathbf{y}) = f(\mathbf{y}).$$

(ii) For any $\mathbf{x} \in \mathbb{R}^d$ and all $\mathbf{y} \in \mathbb{R}^d \setminus \mathcal{A}$,

$$h(\mathbf{x}, \mathbf{y}) \geq 2f(\mathbf{x}) - f(\mathbf{y}).$$

(iii) For any $\mathbf{y} \in \mathbb{R}^d \setminus \mathcal{A}$,

$$T(\mathbf{y}) = \operatorname{argmin}_{\mathbf{x} \in \mathbb{R}^d} h(\mathbf{x}, \mathbf{y}).$$

Proof (i) Follows by simple substitution.

(ii) First, note that, for every two real numbers $a \in \mathbb{R}$ and $b > 0$, the inequality

$$\frac{a^2}{b} \geq 2a - b$$

holds true. Therefore, for every $i = 1, 2, \dots, m$, we have

$$\frac{\|\mathbf{x} - \mathbf{a}_i\|^2}{\|\mathbf{y} - \mathbf{a}_i\|} \geq 2\|\mathbf{x} - \mathbf{a}_i\| - \|\mathbf{y} - \mathbf{a}_i\|.$$

Multiplying the latter inequality by ω_i and summing over $i = 1, 2, \dots, m$, the result follows.

(iii) Follows by the discussion prior to the lemma. □

Using Lemma 3.1, we are now able to prove the monotonicity property of the operator T with respect to f . The proof here relies on the same arguments of Weiszfeld [5] and of Kuhn [14].

Lemma 3.2 (Monotonicity property of T) *For every $\mathbf{y} \in \mathbb{R}^d \setminus \mathcal{A}$, we have*

$$f(T(\mathbf{y})) \leq f(\mathbf{y}), \tag{8}$$

and equality holds if and only if $T(\mathbf{y}) = \mathbf{y}$.

Proof From Lemma 3.1(iii), we have that $T(\mathbf{y}) = \operatorname{argmin}_{\mathbf{x} \in \mathbb{R}^d} h(\mathbf{x}, \mathbf{y})$, and by the strict convexity of the function $\mathbf{x} \rightarrow h(\mathbf{x}, \mathbf{y})$, one has

$$h(T(\mathbf{y}), \mathbf{y}) < h(\mathbf{x}, \mathbf{y}) \tag{9}$$

for every $\mathbf{x} \neq T(\mathbf{y})$. In particular, if $T(\mathbf{y}) \neq \mathbf{y}$, then

$$h(T(\mathbf{y}), \mathbf{y}) < h(\mathbf{y}, \mathbf{y}) = f(\mathbf{y}), \quad (10)$$

where the last equality follows from Lemma 3.1(i). Now, from Lemma 3.1(ii), we have

$$h(T(\mathbf{y}), \mathbf{y}) \geq 2f(T(\mathbf{y})) - f(\mathbf{y}),$$

which, combined with (10), establishes the desired strict monotonicity. \square

Since the general step of Weiszfeld's method is defined by $\mathbf{x}_{k+1} = T(\mathbf{x}_k)$, and since $\nabla f(\mathbf{x}^*) = \mathbf{0}$ if and only if $\mathbf{x}^* = T(\mathbf{x}^*)$ (see (5)), we can immediately conclude that, under the condition that the iterates of Weiszfeld's method do not belong to the anchor set \mathcal{A} , the method is nonincreasing and "gets stuck" only at optimal points.

Corollary 3.1 (Monotonicity of the sequence of function values) *Let $\{\mathbf{x}_k\}_{k \geq 0}$ be the sequence generated by Weiszfeld's method and assume that $\mathbf{x}_k \notin \mathcal{A}$ for any $k \geq 0$. Then, $f(\mathbf{x}_{k+1}) \leq f(\mathbf{x}_k)$ for any $k \geq 0$, and equality holds if and only if \mathbf{x}_k is an optimal solution of problem (FW).*

4 Reinventing the Wheel

Weiszfeld's method remained unknown until 1962. One evidence for this is the fact that the algorithm was rediscovered several times without any knowledge of Weiszfeld's earlier work. As far as we know, the first to rediscover the method was Miehle in 1958 [15], who studied an even more general problem, where the location of several points is to be determined. The paper is formulated in the two-dimensional plane, but the extension to \mathbb{R}^d is obvious. The derivation of the method in [15] is identical to the isolation process described in Sect. 1. Miehle, like Weiszfeld, did not treat the situation in which the method reaches an anchor point.

Four years later, in 1962, Kuhn and Kuenne rediscovered Weiszfeld's method for solving problem (FW) in the plane [12]. At the time of the writing of the paper, Kuhn and Kuenne were not aware of Weiszfeld's work, but an appendix called "added in proof" was added reading as follows.

[12, p. 33]: **Added in proof:** *In the period between the submission of this paper and its publication by the journal, the results have been extended and further literature bearing on the problem has been discovered. We shall sketch the nature of the extensions and list only those references that bear directly on the material developed in the body of the paper. First of all, the algorithm of the paper has been considered, independently of the present account, at least three times. The first published version seems to be that of E. Weiszfeld in the Tohoku Mathematical Journal...*

The story of how the authors were made aware of Weiszfeld's method is known to us since in 2002, Weiszfeld (who changed his name to Andrew Vazsonyi) recalled the following story from the 1960s.

[6, p. 12]: *After reading more, I discovered that a well-known mathematician, Harold W. Kuhn of Princeton University, had given a talk in Budapest on his discovery of an algorithm to solve the location problem. After the talk, a former colleague of mine walked to the blackboard and wrote in big letters: “VAZ-SONYI.”*

“Who is that?” Kuhn asked.

“The name of the Hungarian mathematician who discovered your algorithm thirty years ago”, my old friend said. “He lives in the United States but published his revolutionary approach under the name Weiszfeld.”

In the “Added in proof” section of [12], Kuhn and Kuenne mention that the convergence proof of Weiszfeld contains an error, since the iterates may belong to the anchor set \mathcal{A} . They continue to claim that it can be proven that either $\mathbf{x}_k \notin \mathcal{A}$ for all $k \geq 0$, and convergence to the optimal solution can be guaranteed, or that the method gets stuck at an anchor point ($\mathbf{x}_k \in \mathcal{A}$ for some $k \geq 0$). In addition, they hypothesize that the latter case may only occur in “at most a denumerable number of (starting) points in the convex hull of \mathcal{A} ”. Later on, we will return to these claims and check their validity (see Sect. 6).

Another observation, which appears in Kuhn and Kuenne [12], is that Weiszfeld’s method is, in fact, a gradient method. Indeed, a simple computation shows that an alternative representation of the operator T is given by

$$T(\mathbf{x}) = \mathbf{x} - \frac{1}{L(\mathbf{x})} \nabla f(\mathbf{x}) \quad (\mathbf{x} \notin \mathcal{A}), \tag{11}$$

where the operator $L : \mathbb{R}^d \setminus \mathcal{A} \rightarrow \mathbb{R}_{++}$ is defined by

$$L(\mathbf{x}) := \sum_{i=1}^m \frac{\omega_i}{\|\mathbf{x} - \mathbf{a}_i\|}. \tag{12}$$

Therefore, Weiszfeld’s method can be written as

$$\mathbf{x}_{k+1} = \mathbf{x}_k - \frac{1}{L(\mathbf{x}_k)} \nabla f(\mathbf{x}_k). \tag{13}$$

One year later, in 1963, unaware of Weiszfeld’s contribution, Cooper [16] also reinvented the method, but again for the more general problem of multiple locations in \mathbb{R}^2 . Cooper did not provide a convergence analysis, but mentioned that in his numerical tests, the method works very well in comparison to other methods. It seems that after 1963, researchers from the optimization, as well as the location communities, were very well aware of the method, and Weiszfeld’s original paper [5] got its rightful credit.

5 The Paper of Kuhn from 1973: The Beginning of a (Correct) Convergence Analysis

The 1973 paper of Kuhn [14] is a continuation of his joint paper with Kuenne from 1962 [12]. Besides re-establishing the monotonicity property of the sequence of objective function values (see Corollary 3.1), he was concerned with two theoretical questions—both mentioned in his “added in proof” section of the previous paper [12].

- A. Assuming that all the iterates do not belong to the anchor set \mathcal{A} , *i.e.*, $\mathbf{x}_k \notin \mathcal{A}$ for all $k \geq 0$, can the convergence to the optimal solution be proven?
- B. Is the number of starting points of the method, for which the method “gets stuck” at nonoptimal anchor points, denumerable?

Kuhn’s answer to both questions was yes. Unfortunately, as we will see later on, the answer to the second question was wrong. We will now explore in details each of the two theoretical questions A and B.

5.1 The Convergence of the Sequence

The theorem that Kuhn proved in [14] is now recalled.

Theorem 5.1 (Convergence of Weiszfeld’s method) *Let $\{\mathbf{x}_k\}_{k \geq 0}$ be a sequence generated by the Weiszfeld’s method. If $\mathbf{x}_k \notin \mathcal{A}$ for all $k \geq 0$, then the sequence $\{\mathbf{x}_k\}_{k \geq 0}$ converges to an optimal solution of problem (FW).*

Kuhn proved this theorem under the assumption that the anchors are not collinear. The assumption in the statement of Theorem 5.1, namely that the iterates do not belong to the anchor set \mathcal{A} , is a bit problematic since it is not clear how to guarantee that such a condition will hold. Probably the reason for such an assumption is the empirical observation that the method practically does not “get stuck” at nonoptimal anchor points. The arguments used in [14] for proving Theorem 5.1 are quite lengthy and technical. We will provide here a different proof that utilizes the relations between the objective function f and the auxiliary function h . This approach will be also rather beneficial since it will be the basis for the rate of convergence analysis, which will be discussed in Sect. 8. In addition, the proof does not require the assumption of collinearity of the anchors. Before proving the theorem, we will establish another property of the objective function f . This result is very similar to the so-called “descent lemma” for continuously differentiable functions (see, *e.g.*, [17]). However, its validity for the nonsmooth function f is far from being obvious.

Lemma 5.1 (Descent lemma for the Fermat–Weber objective function) *Suppose that $\mathbf{y} \notin \mathcal{A}$. Then,*

$$f(T(\mathbf{y})) \leq f(\mathbf{y}) + \langle \nabla f(\mathbf{y}), T(\mathbf{y}) - \mathbf{y} \rangle + \frac{L(\mathbf{y})}{2} \|T(\mathbf{y}) - \mathbf{y}\|^2. \quad (14)$$

Proof Note that the function $\mathbf{x} \mapsto h(\mathbf{x}, \mathbf{y})$ is quadratic with associated matrix $L(\mathbf{y})\mathbf{I}$. Therefore, its second-order Taylor expansion around \mathbf{y} is exact and can be written as

$$h(\mathbf{x}, \mathbf{y}) = h(\mathbf{y}, \mathbf{y}) + \langle \nabla_{\mathbf{x}} h(\mathbf{y}, \mathbf{y}), \mathbf{x} - \mathbf{y} \rangle + L(\mathbf{y}) \|\mathbf{x} - \mathbf{y}\|^2.$$

Since $h(\mathbf{y}, \mathbf{y}) = f(\mathbf{y})$ (see Lemma 3.1(i)) and $\nabla_{\mathbf{x}} h(\mathbf{y}, \mathbf{y}) = 2\nabla f(\mathbf{y})$ (simple computation), we have that

$$h(\mathbf{x}, \mathbf{y}) = f(\mathbf{y}) + 2 \langle \nabla f(\mathbf{y}), \mathbf{x} - \mathbf{y} \rangle + L(\mathbf{y}) \|\mathbf{x} - \mathbf{y}\|^2.$$

Substituting $\mathbf{x} = T(\mathbf{y})$ in the latter identity yields

$$h(T(\mathbf{y}), \mathbf{y}) = f(\mathbf{y}) + 2 \langle \nabla f(\mathbf{y}), T(\mathbf{y}) - \mathbf{y} \rangle + L(\mathbf{y}) \|T(\mathbf{y}) - \mathbf{y}\|^2.$$

Hence, from Lemma 3.1(ii), we obtain

$$2f(T(\mathbf{y})) - f(\mathbf{y}) \leq f(\mathbf{y}) + 2 \langle \nabla f(\mathbf{y}), T(\mathbf{y}) - \mathbf{y} \rangle + L(\mathbf{y}) \|T(\mathbf{y}) - \mathbf{y}\|^2.$$

Therefore

$$2f(T(\mathbf{y})) \leq 2f(\mathbf{y}) + 2 \langle \nabla f(\mathbf{y}), T(\mathbf{y}) - \mathbf{y} \rangle + L(\mathbf{y}) \|T(\mathbf{y}) - \mathbf{y}\|^2,$$

which readily implies (14). □

Remark 5.1 It is interesting to note that, in a way, the latter result mimics known results on continuously differentiable functions. Suppose that g is a continuously differentiable function over \mathbb{R}^d , and assume that its gradient ∇g is Lipschitz continuous with parameter L_g , meaning that

$$\|\nabla g(\mathbf{x}) - \nabla g(\mathbf{y})\| \leq L_g \|\mathbf{x} - \mathbf{y}\|, \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^d.$$

Then, the “descent lemma” for such function states that, for any $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$:

$$g(\mathbf{y}) \leq g(\mathbf{x}) + \langle \nabla g(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{L_g}{2} \|\mathbf{x} - \mathbf{y}\|^2.$$

This is one of the key inequalities used to analyze the convergence properties of gradient-based methods, see, for example, [13, 18]. Of course, in our case, f is not differentiable, but nonetheless a version of the descent lemma still holds when \mathbf{x} is specifically chosen as $T(\mathbf{y})$, and when $L(\mathbf{y})$ takes the role of the Lipschitz constant. Note also that as was already mentioned, Weiszfeld’s method is a gradient method with stepsize $1/L(\mathbf{x}_k)$, which is also an indication that $L(\mathbf{x}_k)$ serves as a substitute for the Lipschitz constant since the gradient method for finding the minimizer of g is known to converge when the stepsize is chosen as $1/L_g$.

Using the descent lemma for the function f , we can now prove the following lemma, stating an inequality that will be the basis for the convergence of the sequence, as well as for the rate of convergence analysis that will be derived in Sect. 8. Note that we do not require the assumption on the collinearity of the anchor points, and hence the optimal set is not necessarily a singleton, and it will be denoted from now on by X^* . The optimal value will be denoted by f^* .

Lemma 5.2 *Let $\{\mathbf{x}_k\}_{k \geq 0}$ be a sequence generated by Weiszfeld's method and assume that $\mathbf{x}_k \notin \mathcal{A}$ for all $k \geq 0$. Then, for any $\mathbf{x} \in \mathbb{R}^d$, the following inequality holds:*

$$f(\mathbf{x}_{k+1}) - f(\mathbf{x}) \leq \frac{L(\mathbf{x}_k)}{2} \left(\|\mathbf{x}_k - \mathbf{x}\|^2 - \|\mathbf{x}_{k+1} - \mathbf{x}\|^2 \right). \quad (15)$$

Proof Substituting $\mathbf{y} = \mathbf{x}_k$ in (14) and using the fact that $\mathbf{x}_{k+1} = T(\mathbf{x}_k)$, we obtain

$$f(\mathbf{x}_{k+1}) \leq f(\mathbf{x}_k) + \langle \nabla f(\mathbf{x}_k), \mathbf{x}_{k+1} - \mathbf{x}_k \rangle + \frac{L(\mathbf{x}_k)}{2} \|\mathbf{x}_{k+1} - \mathbf{x}_k\|^2. \quad (16)$$

By the gradient inequality we have that $f(\mathbf{x}_k) \leq f(\mathbf{x}) + \langle \nabla f(\mathbf{x}_k), \mathbf{x}_k - \mathbf{x} \rangle$ for any $\mathbf{x} \in \mathbb{R}^d$, which combined with (16) yields

$$\begin{aligned} f(\mathbf{x}_{k+1}) &\leq f(\mathbf{x}) + \langle \nabla f(\mathbf{x}_k), \mathbf{x}_k - \mathbf{x} \rangle + \langle \nabla f(\mathbf{x}_k), \mathbf{x}_{k+1} - \mathbf{x}_k \rangle + \frac{L(\mathbf{x}_k)}{2} \|\mathbf{x}_{k+1} - \mathbf{x}_k\|^2 \\ &= f(\mathbf{x}) + \langle \nabla f(\mathbf{x}_k), \mathbf{x}_{k+1} - \mathbf{x} \rangle + \frac{L(\mathbf{x}_k)}{2} \|\mathbf{x}_{k+1} - \mathbf{x}_k\|^2 \\ &= f(\mathbf{x}) + L(\mathbf{x}_k) \langle \mathbf{x}_k - \mathbf{x}_{k+1}, \mathbf{x}_{k+1} - \mathbf{x} \rangle + \frac{L(\mathbf{x}_k)}{2} \|\mathbf{x}_{k+1} - \mathbf{x}_k\|^2 \\ &= f(\mathbf{x}) + \frac{L(\mathbf{x}_k)}{2} \left(\|\mathbf{x}_k - \mathbf{x}\|^2 - \|\mathbf{x}_{k+1} - \mathbf{x}\|^2 \right), \end{aligned}$$

where the second equality follows from (13), and the last equality follows from the identity that

$$2 \langle \mathbf{w} - \mathbf{v}, \mathbf{u} - \mathbf{v} \rangle = \|\mathbf{w} - \mathbf{v}\|^2 - \|\mathbf{w} - \mathbf{u}\|^2 + \|\mathbf{u} - \mathbf{v}\|^2,$$

for any $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^d$. This completes the proof. \square

Let $\{\mathbf{x}_k\}_{k \geq 0}$ be the sequence generated by Weiszfeld's method with initial point \mathbf{x}_0 . In those cases that the left-hand side of (15) is nonnegative, that is, when $f(\mathbf{x}) \leq f(\mathbf{x}_k)$ for all $k \geq 0$, we get as a direct result of the latter lemma the so-called Fejér monotonicity of $\{\mathbf{x}_k\}_{k \geq 0}$ —a result which seems to be unknown in the literature.

Corollary 5.1 (Fejér monotonicity) *Let $\{\mathbf{x}_k\}_{k \geq 0}$ be a sequence generated by Weiszfeld's method and assume that $\mathbf{x}_k \notin \mathcal{A}$ for all $k \geq 0$. Then, for any $\mathbf{x} \in \mathbb{R}^d$ which satisfies $f(\mathbf{x}) \leq f(\mathbf{x}_k)$ for all $k \geq 0$, the following inequality holds:*

$$\|\mathbf{x}_{k+1} - \mathbf{x}\| \leq \|\mathbf{x}_k - \mathbf{x}\|.$$

Hence the sequence $\{\mathbf{x}_k\}_{k \geq 0}$ is bounded.

Proof Follows directly from (15) along with the fact that $f(\mathbf{x}_{k+1}) \geq f(\mathbf{x})$. The boundedness of the sequence then readily follows by the fact that $\|\mathbf{x}_k - \mathbf{x}\| \leq \|\mathbf{x}_0 - \mathbf{x}\|$ for any $k \geq 0$. \square

The convergence result of Kuhn, namely Theorem 5.1, can now be easily deduced from the Fejér monotonicity property of the sequence $\{\mathbf{x}_k\}_{k \geq 0}$.

Proof of Theorem 5.1 We will prove this result in two steps. First, we will prove that $\{\mathbf{x}_k\}_{k \geq 0}$ converges, and then we will show that its limit point is an optimal solution of problem (FW).

The sequence $\{\mathbf{x}_k\}_{k \geq 0}$ is bounded by Corollary 5.1. To prove the convergence of $\{\mathbf{x}_k\}_{k \geq 0}$, it only remains to show that all converging subsequences have the same limit. Suppose in contradiction that there exist two subsequences $\{\mathbf{x}_{k_j}\}_{j \geq 0}$ and $\{\mathbf{x}_{n_j}\}_{j \geq 0}$ converging to different limits $\tilde{\mathbf{x}}$ and $\bar{\mathbf{x}}$, respectively.

From Corollary 3.1, it follows that $f(\tilde{\mathbf{x}}) \leq f(\mathbf{x}_k)$ for all $k \geq 0$, and thus from Corollary 5.1, we get that the sequence $\{\|\mathbf{x}_k - \tilde{\mathbf{x}}\|\}_{k \geq 0}$ is nonincreasing. Since this sequence is also bounded from below, it converges to some $l_1 \in \mathbb{R}$. Clearly

$$\lim_{k \rightarrow \infty} \|\mathbf{x}_k - \tilde{\mathbf{x}}\| = \lim_{j \rightarrow \infty} \|\mathbf{x}_{k_j} - \tilde{\mathbf{x}}\| = 0.$$

However, on the other hand,

$$\lim_{k \rightarrow \infty} \|\mathbf{x}_k - \tilde{\mathbf{x}}\| = \lim_{j \rightarrow \infty} \|\mathbf{x}_{n_j} - \tilde{\mathbf{x}}\| = \|\bar{\mathbf{x}} - \tilde{\mathbf{x}}\|.$$

Hence $l_1 = 0 = \|\bar{\mathbf{x}} - \tilde{\mathbf{x}}\|$, which is obviously a contradiction. This proves that $\{\mathbf{x}_k\}_{k \geq 0}$ converges.

We denote by $\tilde{\mathbf{x}}$ the limit of $\{\mathbf{x}_k\}_{k \geq 0}$. Now we will prove that $\tilde{\mathbf{x}}$ is an optimal solution of problem (FW). It is clear that, if $\tilde{\mathbf{x}} \notin \mathcal{A}$, then, taking the limit as $k \rightarrow \infty$ in the equation $\mathbf{x}_{k+1} = T(\mathbf{x}_k)$, and using the continuity of the operator T at nonanchor points, we obtain that $\tilde{\mathbf{x}} = T(\tilde{\mathbf{x}})$. The optimality of $\tilde{\mathbf{x}}$ now follows immediately from (5). On the other hand, if $\tilde{\mathbf{x}} \in \mathcal{A}$, then there exists $j \in \{1, 2, \dots, m\}$ such that $\tilde{\mathbf{x}} = \mathbf{a}_j$. From Lemma 3.1(iii) we have

$$\nabla_{\mathbf{x}} h(\mathbf{x}_{k+1}, \mathbf{x}_k) = \mathbf{0},$$

which can be written explicitly as follows:

$$\sum_{i=1}^m \omega_i \frac{\mathbf{x}_{k+1} - \mathbf{a}_i}{\|\mathbf{x}_k - \mathbf{a}_i\|} = \mathbf{0}.$$

Thus,

$$\sum_{i=1, i \neq j}^m \omega_i \frac{\mathbf{x}_{k+1} - \mathbf{a}_i}{\|\mathbf{x}_k - \mathbf{a}_i\|} = -\omega_j \frac{\mathbf{x}_{k+1} - \mathbf{a}_j}{\|\mathbf{x}_k - \mathbf{a}_j\|},$$

and, after taking the norm on both sides, we get

$$\left\| \sum_{i=1, i \neq j}^m \omega_i \frac{\mathbf{x}_{k+1} - \mathbf{a}_i}{\|\mathbf{x}_k - \mathbf{a}_i\|} \right\| = \omega_j \frac{\|\mathbf{x}_{k+1} - \mathbf{a}_j\|}{\|\mathbf{x}_k - \mathbf{a}_j\|} \leq \omega_j, \tag{17}$$

where the inequality follows from the fact that the sequence $\{\|\mathbf{x}_k - \mathbf{a}_j\|\}_{k \geq 0}$ is non-increasing (see Corollary 5.1 using the fact that $f(\mathbf{a}_j) \leq f(\mathbf{x}_k)$ for all $k \geq 0$ from Corollary 3.1). Taking the limit in (17) as $k \rightarrow \infty$ yields the inequality

$$\left\| \sum_{i=1, i \neq j}^m \omega_i \frac{\tilde{\mathbf{x}} - \mathbf{a}_i}{\|\tilde{\mathbf{x}} - \mathbf{a}_i\|} \right\| \leq \omega_j,$$

which, by Theorem 2.1, shows that $\tilde{\mathbf{x}} = \mathbf{a}_j$ is an optimal solution of (FW). □

5.2 “Bad” Starting Points

As was already mentioned, even in the earlier paper of Kuhn and Kuenne [12] from 1962, it was obvious that Weiszfeld’s method can reach a nonoptimal anchor point. This situation is described in the literature as “getting stuck”, and starting points of the method leading to this situation are called “bad” starting points. In his 1973 paper, Kuhn gave an example of such a case, as we recall now.

Example 5.1 (Kuhn’s counterexample) The example is in the 2-dimensional space, and the anchors are $\mathbf{a}_1 = (-2, 0)^T$, $\mathbf{a}_2 = (-1, 0)^T$, $\mathbf{a}_3 = (1, 0)^T$, $\mathbf{a}_4 = (2, 0)^T$, $\mathbf{a}_5 = (0, 1)^T$, and $\mathbf{a}_6 = (0, -1)^T$. All the weights are one. It is easy to see that $\nabla f(0, 0) = \mathbf{0}$, so that the optimal solution of the problem is $\mathbf{x}^* = \mathbf{0}$. Kuhn then studied the behavior of the operator T on points on the x -axis given by $(x, 0)^T$:

$$\begin{aligned} T\left((x, 0)^T\right) &= \frac{1}{\sum_{i=1}^6 \frac{1}{\|(x, 0)^T - \mathbf{a}_i\|}} \sum_{i=1}^6 \frac{\mathbf{a}_i}{\|(x, 0)^T - \mathbf{a}_i\|} \\ &= \frac{\begin{pmatrix} \frac{-2}{|x+2|} + \frac{-1}{|x+1|} + \frac{1}{|x-1|} + \frac{2}{|x-2|} \\ 0 \end{pmatrix}}{\frac{1}{|x+2|} + \frac{1}{|x+1|} + \frac{1}{|x-1|} + \frac{1}{|x-2|} + \frac{2}{\sqrt{x^2+1}}} := \begin{pmatrix} g(x) \\ 0 \end{pmatrix}, \end{aligned}$$

and then Kuhn found a value $\alpha \in [0, 2]$ such that $g(\alpha) = 1$. This shows that $T((\alpha, 0)^T) = \mathbf{a}_3$. The plot of g over the interval $[0, 2]$ can be found in Fig. 2, where the solution, which is $\alpha = 1.6213$ (up to three digits of accuracy), is described.

Kuhn claimed that the number of “bad” starting points is always denumerable. The key argument used to show this result is quoted here.

[14, p. 107]: *If we insert T from a vertex \mathbf{a}_i , we must solve algebraic equations. Thus we obtain a finite number of \mathbf{x}_0 such that $T(\mathbf{x}_0) = \mathbf{a}_i$.*




Fig. 2 Kuhn’s construction of a “bad” starting point. $T((\alpha, 0)^T) = \mathbf{a}_3$, where α is the solution of the equation $g(\alpha) = 1$

However, this argument is incorrect, as will be explained in the following section.

6 Counting the “Bad” Starting Points

The flaw in Kuhn’s argument is that, actually, algebraic system of equations can have a continuum number of solutions. This was shown by Chandrasekaran and Tamir in their 1989 paper [19], where the following counterexample to Kuhn’s claim in \mathbb{R}^3 was given.

Example 6.1 (Chandrasekaran and Tamir’s counterexample to Kuhn) Consider the problem in \mathbb{R}^3 with anchors $\mathbf{a}_1 = (1, 0, 0)^T$, $\mathbf{a}_2 = (-1, 0, 0)^T$, $\mathbf{a}_3 = (0, 0, 0)^T$, $\mathbf{a}_4 = (0, 2, 0)^T$, and $\mathbf{a}_5 = (0, -2, 0)^T$. Let $\omega_1 = \omega_2 = \omega_3 = \omega_5 = 1$, and $\omega_4 = 3$. Consider the point \mathbf{a}_3 , which is not optimal since

$$f(\mathbf{a}_3) = 10 > f(0, 1, 0) = 7 + 2\sqrt{2}.$$

To show that the algebraic system $T(\mathbf{x}) = \mathbf{a}_3$ has an infinite number of solutions, consider the points of the form $\mathbf{x} = (0, y, z)^T$. Then, $T(\mathbf{x}) = \mathbf{a}_3$ is equivalent to the system

$$\frac{3\mathbf{a}_4}{\|\mathbf{x} - \mathbf{a}_4\|} + \frac{\mathbf{a}_5}{\|\mathbf{x} - \mathbf{a}_5\|} = 0.$$

That is,

$$\frac{1}{\sqrt{(y-2)^2+z^2}}(0,6,0)^T + \frac{1}{\sqrt{(y+2)^2+z^2}}(0,-2,0)^T = (0,0,0)^T,$$

which is the same as

$$36(y+2)^2 + 36z^2 - 4((y-2)^2 + z^2) = 0.$$

After some simple algebraic manipulation, we conclude that all the points on the circle $(y+2.5)^2 + z^2 = 2.25$ solve the system $T(\mathbf{x}) = \mathbf{a}_3$.

The latter example is special in the sense that the anchors, although not collinear, reside in a lower dimensional affine subspace. Chandrasekaran and Tamir conjectured that such a situation is the only one in which a continuum of “bad” starting points can occur.

[19, p. 295]: “In view of the above examples we conjecture that, if the non-collinearity is replaced by the stronger assumption that the convex hull of the points $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$ is of full dimension, then the algebraic system $T(\mathbf{x}) = \mathbf{a}_i$ has a finite number of solutions for $i = 1, 2, \dots, m$. Phrased differently, the conjecture is that, for each $i = 1, 2, \dots, m$, there is a finite number of solutions to $T(\mathbf{x}) = \mathbf{a}_i$ in the minimal affine set containing the points $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$.”

To write it explicitly, the first conjecture of Chandrasekaran and Tamir is formulated as follows.

Conjecture 6.1 (Chandrasekaran and Tamir conjecture) When the affine hull of the anchor set \mathcal{A} is the entire space \mathbb{R}^d , the number of solutions of the system $T(\mathbf{x}) = \mathbf{a}_i$, for any $i = 1, 2, \dots, m$, is finite, and hence there is a denumerable number of “bad” starting points.

As quoted above, Chandrasekaran and Tamir also had a more general conjecture that the number of “bad” starting points is denumerable when the starting point is restricted a priori to be in the affine hull of the anchor set \mathcal{A} . Conjecture 6.1 was resolved in 1995 by Brimberg [20]. More precisely, Brimberg proved the following result (written in this paper’s terminology).

Theorem 6.1 (cf. [20, Theorem 1, p. 75]) *The set of starting points \mathbf{x}_0 , which will terminate the sequence generated by Weiszfeld’s method at some anchor point \mathbf{a}_i , $i = 1, 2, \dots, m$, after a finite number of iterations is denumerable if the affine hull of \mathcal{A} is \mathbb{R}^d .*

At this point, one would think that the issue of “counting” the number of bad starting points was resolved. However, Brimberg also claimed that, in fact, the number of “bad” starting points is denumerable *if and only if* the affine hull of \mathcal{A} is \mathbb{R}^d . The “only if” claim is not part of Chandrasekaran and Tamir’s conjecture. Later on, in 2002, Cánovas, Cañavate, and Marín showed in [21] that the “only if” claim is not

correct by providing two counterexamples. We will present the first counterexample which shows a setting in which the number of “bad” starting points is denumerable (in fact zero), even though the affine hull of \mathcal{A} is not the entire space \mathbb{R}^d .

Example 6.2 (Counterexample to Brimberg’s “only if” part) Consider the unweighted Fermat–Weber problem with $m = n = 3$ given by the anchors $\mathbf{a}_1 = (1, 0, 0)^T$, $\mathbf{a}_2 = (0, 1, 0)^T$ and $\mathbf{a}_3 = (0, 0, 0)$. Note that \mathbf{a}_1 , \mathbf{a}_2 , and \mathbf{a}_3 are the extreme points of $\text{conv}(\mathcal{A})$, and that, for any $\mathbf{x} \notin \mathcal{A}$,

$$T(\mathbf{x}) = \frac{1}{\sum_{i=1}^3 \frac{1}{\|\mathbf{x} - \mathbf{a}_i\|}} \sum_{i=1}^3 \frac{1}{\|\mathbf{x} - \mathbf{a}_i\|} \mathbf{a}_i.$$

Hence, $T(\mathbf{x})$ is a convex combination of \mathbf{a}_1 , \mathbf{a}_2 and \mathbf{a}_3 with positive coefficients. However, this means that $T(\mathbf{x})$ cannot be equal to any of the extreme points, that is, to an anchor point. Therefore, the system $T(\mathbf{x}) = \mathbf{a}_i$ has no solutions for any $i = 1, 2, 3$, and in this case, Weiszfeld’s method is well defined for any starting point which is not in \mathcal{A} .

7 Bypassing the Anchor Points: Modifying the Method

7.1 The Modified Weiszfeld’s Method

Since the issue of bumping into anchor points is an important issue in the convergence analysis of the method, these instances should be treated. In this section, we will begin by reviewing a modification of the method in which the problem of reaching an anchor point is treated in a “surgical” manner. Specifically, the class of methods that we consider coincides with Weiszfeld’s method when the current iterate \mathbf{x}_k , for some $k \geq 0$, is not an anchor point; and when the iterate is equal to some \mathbf{a}_i , then the next iterate will be equal to \mathbf{a}_i if \mathbf{a}_i is optimal, and equal to another point, with a smaller function value when \mathbf{a}_i is not optimal. To test the optimality of the anchor points, we are required to define the following quantities (see the optimality conditions in Theorem 2.1):

$$\mathbf{R}_j := \sum_{i=1, i \neq j}^m \omega_i \frac{\mathbf{a}_j - \mathbf{a}_i}{\|\mathbf{a}_i - \mathbf{a}_j\|}, \quad j = 1, 2, \dots, m.$$

The anchor point \mathbf{a}_j , $j = 1, 2, \dots, m$, is optimal if and only if

$$\|\mathbf{R}_j\| \leq w_j.$$

The general scheme for the modified approach can be written as follows.

Modified Weiszfeld's Method**Initialization.** $\mathbf{x}_0 \in \mathbb{R}^d$.**General Step** ($k = 0, 1, \dots$)

$$\mathbf{x}_{k+1} = \tilde{T}(\mathbf{x}_k) = \begin{cases} T(\mathbf{x}_k), & \mathbf{x}_k \notin \mathcal{A}, \\ \mathbf{a}_j, & \mathbf{x}_k = \mathbf{a}_j \ (1 \leq j \leq m) \text{ and } \|\mathbf{R}_j\| \leq \omega_j \\ S(\mathbf{a}_j), & \mathbf{x}_k = \mathbf{a}_j \ (1 \leq j \leq m) \text{ and } \|\mathbf{R}_j\| > \omega_j. \end{cases} \quad (18)$$

The new operator \tilde{T} coincides with the usual Weiszfeld's operator T at nonanchor points. At nonoptimal anchor points, another operator, which is denoted by S , is invoked. The question that arises is of course how to define the operator S on a specific nonoptimal anchor point \mathbf{a}_j , $j = 1, 2, \dots, m$. However, at this point, we will just require that, for a nonoptimal anchor point \mathbf{a}_j , the point $S(\mathbf{a}_j)$ will have a smaller function value than \mathbf{a}_j . Under this condition, based on Kuhn's convergence result (see Theorem 5.1), we can prove convergence of the sequence generated by the modified Weiszfeld's method. Note that we do not require the assumption of noncollinearity that is always assumed in the literature.

Theorem 7.1 (Convergence of the modified Weiszfeld's method) *Let $\{\mathbf{x}_k\}_{k \geq 0}$ be a sequence generated by the modified Weiszfeld's method. Assume that $f(S(\mathbf{a}_j)) < f(\mathbf{a}_j)$ for any $\mathbf{a}_j \in \mathcal{A}$ for which $\|\mathbf{R}_j\| > \omega_j$. Then, the sequence $\{\mathbf{x}_k\}_{k \geq 0}$ converges to an optimal solution of problem (FW).*

Proof There are two options. Either the sequence “gets stuck” at a fixed point of the method, i.e., at a point $\tilde{\mathbf{x}}$ for which $\tilde{T}(\tilde{\mathbf{x}}) = \tilde{\mathbf{x}}$. If $\tilde{\mathbf{x}} \notin \mathcal{A}$, then by (5), it follows that $\nabla f(\tilde{\mathbf{x}}) = 0$, and hence the method converges to an optimal solution $\tilde{\mathbf{x}}$. If $\tilde{\mathbf{x}} = \mathbf{a}_j$ for some $j \in \{1, 2, \dots, m\}$, then, since $S(\mathbf{a}_j) \neq \mathbf{a}_j$ if \mathbf{a}_j is not optimal, it follows that \mathbf{a}_j must be optimal, and we conclude that once again the sequence generated by the method converges to an optimal solution $\tilde{\mathbf{x}}$. The second option is when the sequence does not get stuck at a fixed point. In this case, the sequence generated by the method is strictly monotone: $f(\mathbf{x}_{k+1}) < f(\mathbf{x}_k)$ for all $k \geq 0$ (this follows from Corollary 3.1 and the assumption that $f(S(\mathbf{a}_j)) < f(\mathbf{a}_j)$ for any $\mathbf{a}_j \in \mathcal{A}$ for which $\|\mathbf{R}_j\| > \omega_j$). Therefore, the iterates are different from each other and thus, since there are only a finite number of anchor points, it follows that there exists a positive integer K such that $\mathbf{x}_k \notin \mathcal{A}$ for all $k \geq K$. The sequence $\{\mathbf{x}_k\}_{k \geq K}$ is the sequence generated by Weiszfeld's method with initial point \mathbf{x}_K , and since none of its elements is in \mathcal{A} , it follows from Theorem 5.1 that it converges to an optimal solution of problem (FW). \square

Note that since, in practice, Weiszfeld's method does not actually reach anchor points, the modified method is actually the same as Weiszfeld's method for all practical purposes.

7.2 Choosing the Operator S

The most natural way to define S on an anchor point $\mathbf{a}_j, j = 1, 2, \dots, m$, is to find a descent direction of f at \mathbf{a}_j , and take a step along this direction. To find such a descent direction, note that the objective function f can be written as

$$f(\mathbf{x}) = \omega_j \|\mathbf{x} - \mathbf{a}_j\| + f_j(\mathbf{x}),$$

where

$$f_j(\mathbf{x}) := \sum_{i=1, i \neq j}^m \omega_i \|\mathbf{x} - \mathbf{a}_i\|.$$

Therefore, taking a direction $\mathbf{d} \in \mathbb{R}^d$ satisfying $\|\mathbf{d}\| = 1$, we can define the function

$$\alpha(t) = f(\mathbf{a}_j + t\mathbf{d}) = \omega_j t \|\mathbf{d}\| + f_j(\mathbf{a}_j + t\mathbf{d}) = \omega_j t + f_j(\mathbf{a}_j + t\mathbf{d}).$$

The directional derivative of f at \mathbf{a}_j in the direction of \mathbf{d} is given by (note that f_j is differentiable at \mathbf{a}_j):

$$f'(\mathbf{a}_j; \mathbf{d}) = \alpha'(0) = \omega_j + f'_j(\mathbf{a}_j; \mathbf{d}) = \omega_j + \langle \nabla f_j(\mathbf{a}_j), \mathbf{d} \rangle.$$

The smallest directional derivative is attained at $\mathbf{d} = \mathbf{d}_j$, where $\mathbf{d}_j = -\nabla f_j(\mathbf{a}_j) / \|\nabla f_j(\mathbf{a}_j)\|$, which is the steepest descent direction. Since $\nabla f_j(\mathbf{a}_j) = \mathbf{R}_j$, we can summarize by writing that the steepest descent direction of f at \mathbf{a}_j is

$$\mathbf{d}_j = -\frac{\mathbf{R}_j}{\|\mathbf{R}_j\|}.$$

In all of the papers that deal with this slightly modified Weiszfeld’s method, the operator S is chosen as

$$S(\mathbf{a}_j) = \mathbf{a}_j + t_j \mathbf{d}_j,$$

where t_j is some well chosen stepsize. Several choices of the stepsize were discussed in the literature. Ostresh [8] (1978) considered the following stepsize:

$$t_j = t_O := c \frac{\|\mathbf{R}_j\| - \omega_j}{L(\mathbf{a}_j)}, \quad [\text{Ostresh [8]}]$$

where $c \in [1, 2]$, and the definition of the operator L (originally given in (12)) is extended to include also anchor points:

$$L(\mathbf{x}) = \begin{cases} \sum_{i=1}^m \frac{\omega_i}{\|\mathbf{x} - \mathbf{a}_i\|}, & \mathbf{x} \notin \mathcal{A}, \\ \sum_{i=1, i \neq j}^m \frac{\omega_i}{\|\mathbf{a}_j - \mathbf{a}_i\|}, & \mathbf{x} = \mathbf{a}_j \ (1 \leq j \leq m). \end{cases} \quad (19)$$

A totally different analysis of Vardi and Zhang [22] (2001) results with the stepsize

$$t_j = t_{VZ} := \frac{\|\mathbf{R}_j\| - \omega_j}{L(\mathbf{a}_j)}, \quad [\text{Vardi and Zhang [22]}]$$

meaning that this is the stepsize of Ostresh with $c = 1$. The following different stepsize was considered by Rautenbach et. al. [23] (2004),

$$t_j = t_R := \min \left\{ \frac{s_j}{2}, \frac{\|\mathbf{R}_j\| - \omega_j}{4L(\mathbf{a}_j)} \right\}, \quad [\text{Rautenbach et. al. [23]}]$$

where $s_j := \min \{\|\mathbf{a}_j - \mathbf{a}_i\| : i \neq j, 1 \leq i \leq m\}$. Obviously, the largest step is the one given by Ostresh when c is taken to be 2, and the smallest step is the one given by Rautenbach. We will now give a proof that, indeed, the stepsize given by Vardi and Zhang results with a decrease in the function value. We also give an explicit expression for a lower bound on the amount of decrease resulting from taking the Vardi-Zhang stepsize. This new result will be important later on in establishing the complexity results of the method (see Sect. 8). The technical and lengthy proof of the lemma can be found in Appendix B.

Lemma 7.1 *Suppose that \mathbf{a}_j , for some $j \in \{1, 2, \dots, m\}$, is not an optimal solution of problem (FW), meaning that $\|\mathbf{R}_j\| > \omega_j$. Then*

$$f(\mathbf{a}_j) - f(\mathbf{a}_j + t_j \mathbf{d}_j) \geq \frac{(\|\mathbf{R}_j\| - \omega_j)^2}{2L(\mathbf{a}_j)},$$

where $t_j = t_{VZ} = (\|\mathbf{R}_j\| - \omega_j) / L(\mathbf{a}_j)$.

7.3 Choosing the Starting Point

The modified method was devised in order to relax the assumption that $\mathbf{x}_k \notin \mathcal{A}$ for all $k \geq 0$ in Kuhn’s convergence result. However, we can use the operator S to carefully pick a starting point that will guarantee that the iterates will not coincide with anchor points. The simple procedure for choosing the starting point, which we call “the SP method”, is now described.

The SP method

(a) Let

$$p \in \operatorname{argmin} \{f(\mathbf{a}_i) : 1 \leq i \leq m\}.$$

(b) If \mathbf{a}_p is optimal (easily checked by verifying that $\|\mathbf{R}_p\| \leq \omega_p$), then the output is \mathbf{a}_p with an indication that it is the optimal solution. Otherwise, if \mathbf{a}_p is not optimal, the output is the point $S(\mathbf{a}_p)$.

The important property of the SP method is that, if there is no optimal anchor point, then the output of the method is a starting point \mathbf{x}_0 which satisfies

$$f(\mathbf{x}_0) < \min \{f(\mathbf{a}_1), f(\mathbf{a}_2), \dots, f(\mathbf{a}_m)\}.$$

The latter inequality, along with the monotonicity property of the sequence of function values (see Corollary 3.1) implies that $\mathbf{x}_k \notin \mathcal{A}$ for all $k \geq 0$, and hence the convergence

of the sequence to an optimal solution is assured by Theorem 5.1. We summarize this discussion in the following corollary. The underlying assumption is that there are no optimal anchor points (otherwise, the SP method will produce the optimal solution). In addition, under this assumption the anchors are necessarily not collinear, implying that the optimal solution is unique.

Corollary 7.1 *Suppose that there is no optimal anchor point for problem (FW). Let $\{\mathbf{x}_k\}_{k \geq 0}$ be the sequence generated by Weiszfeld’s method with starting point \mathbf{x}_0 produced by the SP method. Then, $\mathbf{x}_k \notin \mathcal{A}$ for all $k \geq 0$ and $\mathbf{x}_k \rightarrow \mathbf{x}^*$ as $k \rightarrow \infty$.*

7.4 Further Modifications of Weiszfeld’s Method

Aside of the rather local modifications of Weiszfeld’s method mentioned in the previous subsections, many more modifications were suggested in the literature. In the 1978 paper [8], Ostresh suggested to accelerate the Weiszfeld’s method by using the following idea. Let $\lambda \in [1, 2]$ and consider the operator

$$T_\lambda(\mathbf{y}) = \mathbf{y} + \lambda(T(\mathbf{y}) - \mathbf{y}).$$

Ostresh proved the convergence of the modified method defined by $\mathbf{x}_{k+1} = T_\lambda(\mathbf{x}_k)$ for any choice of $\lambda \in [1, 2]$, and under a suitable treatment of the anchor points. The use of a stepsize can also be found in [24] (1984). In [25] (1992), Drezner uses an adaptive method in order to choose a different λ at each iteration, so that the method reads as $\mathbf{x}_{k+1} = T_{\lambda_k}(\mathbf{x}_k)$. Another acceleration technique was proposed by Drezner in [26] (1995), where Steffensen’s method, which is a general acceleration scheme for fixed point method, was tested numerically against Weiszfeld’s method with several stepsize strategies for choosing λ .

Deviating from the main focus of this paper, which is Weiszfeld’s method, we note that the Fermat–Weber problem can also be solved by more sophisticated methods than Weiszfeld’s method. For example, Calamai and Conn [27] and Overton [28] solved the more general problem of minimizing a sum of norms of affine functions by Newton methods combined with an active set approach. In addition, the problem can be recast as a second-order cone programming and solved via interior point methods [29,30]. A specially devised primal-dual interior point method for the minimization of the sum of Euclidean norms was analyzed by Andersen et. al. in [31].

There are, of course, many variations of Weiszfeld’s method when the problem to be solved is not the Fermat–Weber problem, but this will be the subject of Sect. 10.

8 Rate of Convergence of Weiszfeld’s Method

Local and asymptotic rate of convergence of Weiszfeld’s method was discussed in the 1974 paper of Katz [32]. However, there does not seem to be a global nonasymptotic rate of convergence analysis of Weiszfeld’s method in the literature. The main objective of this section is to derive such rate of convergence, but before, we would like to recall

several related results on global nonasymptotic rate of convergence for gradient-based methods. This type of results is usually of the form

$$f(\mathbf{x}_k) - f^* \leq \frac{C}{k^\theta},$$

where $C, \theta > 0$ are constants. For example, for nonsmooth convex problems we can employ, under some unrestrictive conditions, the subgradient method to solve problem (FW). This will result with a rate of convergence of $O(1/\sqrt{k})$ (that is, $\theta = 1/2$), see e.g., [18,33]. In this case, since the method is not monotone, the bound is on $\min_{n=1,2,\dots,k} f(\mathbf{x}_n) - f^*$, rather than on $f(\mathbf{x}_k) - f^*$. If the problem is smooth, then the gradient method can be employed, and the convergence rate will accelerate to $O(1/k)$ (corresponding to $\theta = 1$). Another option is to use an “optimal” gradient method, which uses the memory of the last two iterations. These methods have a rate of convergence of $O(1/k^2)$ (corresponding to $\theta = 2$). For a wealth of fundamental results on these issues, see for instance, [13,18,34,35].

In principal, since problem (FW) is nonsmooth, and since Weiszfeld’s method is a type of gradient method, it seems logical to assume that only the inferior $O(1/\sqrt{k})$ can be established. However, as we shall see, when choosing the starting point carefully, we can actually prove a rate of convergence of $O(1/k)$ even though the problem is nonsmooth. Later on, we will even show that, by modifying the method, we can establish the fast $O(1/k^2)$ rate of convergence.

Throughout this section, we assume that the starting point is chosen according to the SP method with the Vardi-Zhang stepsize strategy. That is,

$$\mathbf{x}_0 = \mathbf{a}_p + t_p \mathbf{d}_p, \tag{20}$$

where
$$p \in \operatorname{argmin} \{f(\mathbf{a}_i) : 1 \leq i \leq m\}, \tag{21}$$

and
$$\mathbf{d}_p = -\frac{\mathbf{R}_p}{\|\mathbf{R}_p\|}, \quad t_p = \frac{\|\mathbf{R}_p\| - \omega_p}{L(\mathbf{a}_p)}. \tag{22}$$

The only assumption on the data is that there is no optimal anchor point. Otherwise, the SP method will produce an optimal solution. Note also that, in this setting, the anchors are necessarily not collinear, and hence there exists a unique optimal solution \mathbf{x}^* . Note that in the premise of the following theorem, which will be the basis for the main convergence analysis, we assume that the sequence $\{L(\mathbf{x}_k)\}_{k \geq 0}$ is upper bounded. Later on, in Lemma 8.2, we will show the validity of this assumption, as well as find an explicit expression for the upper bound.

Theorem 8.1 (Sublinear rate of convergence of Weiszfeld’s method) *Let $\{\mathbf{x}_k\}_{k \geq 0}$ be the sequence generated by Weiszfeld’s method with \mathbf{x}_0 chosen by (20)–(22). Then, for any $k \geq 0$, we have*

$$f(\mathbf{x}_k) - f^* \leq \frac{\bar{L}}{2k} \|\mathbf{x}_0 - \mathbf{x}^*\|^2, \tag{23}$$

where \bar{L} is an upper bound of the sequence $\{L(\mathbf{x}_k)\}_{k \geq 0}$.

Proof From (15) with $k = n$, we have

$$f(\mathbf{x}_{n+1}) - f^* \leq \frac{L(\mathbf{x}_n)}{2} \left(\|\mathbf{x}_n - \mathbf{x}^*\|^2 - \|\mathbf{x}_{n+1} - \mathbf{x}^*\|^2 \right).$$

Therefore, by the Fejér monotonicity of the sequence $\{\mathbf{x}_k\}_{k \geq 0}$ (see Corollary 5.1) and the boundedness of the sequence $\{L(\mathbf{x}_k)\}_{k \geq 0}$, we get

$$f(\mathbf{x}_{n+1}) - f^* \leq \frac{\bar{L}}{2} \left(\|\mathbf{x}_n - \mathbf{x}^*\|^2 - \|\mathbf{x}_{n+1} - \mathbf{x}^*\|^2 \right).$$

Summing this inequality over $n = 0, \dots, k - 1$ gives

$$\begin{aligned} \sum_{n=0}^{k-1} (f(\mathbf{x}_{n+1}) - f^*) &\leq \frac{\bar{L}}{2} \sum_{n=0}^{k-1} \left(\|\mathbf{x}_n - \mathbf{x}^*\|^2 - \|\mathbf{x}_{n+1} - \mathbf{x}^*\|^2 \right) \\ &= \frac{\bar{L}}{2} \left(\|\mathbf{x}_0 - \mathbf{x}^*\|^2 - \|\mathbf{x}_k - \mathbf{x}^*\|^2 \right) \\ &\leq \frac{\bar{L}}{2} \|\mathbf{x}_0 - \mathbf{x}^*\|^2. \end{aligned}$$

Since the sequence $\{f(\mathbf{x}_k)\}_{k \geq 0}$ is nonincreasing (see Corollary 3.1) we have

$$k(f(\mathbf{x}_k) - f^*) \leq \sum_{n=0}^{k-1} (f(\mathbf{x}_{n+1}) - f^*) \leq \frac{\bar{L}}{2} \|\mathbf{x}_0 - \mathbf{x}^*\|^2,$$

that is,

$$f(\mathbf{x}_k) - f^* \leq \frac{\bar{L}}{2k} \|\mathbf{x}_0 - \mathbf{x}^*\|^2.$$

This proves the desired result. □

All that is left is to find an explicit upper bound of the sequence $\{L(\mathbf{x}_k)\}_{k \geq 0}$. For that, we will use the following result, which establishes a lower bound on the distance between the anchor points to those points with smaller function values than $f(\mathbf{x}_0)$, where \mathbf{x}_0 is chosen by (20)–(22).

Lemma 8.1 (Lower bound on $\|\mathbf{a}_i - \mathbf{x}\|$) *Let \mathbf{x}_0 be chosen by (20)–(22). Then, for any $i = 1, 2, \dots, m$ and any \mathbf{x} satisfying $f(\mathbf{x}) \leq f(\mathbf{x}_0)$, the inequality*

$$\|\mathbf{x} - \mathbf{a}_i\| \geq \frac{f(\mathbf{a}_i) - f(\mathbf{x}_0)}{\omega}$$

holds with $\omega = \sum_{i=1}^m \omega_i$.

Proof From the fact that $f(\mathbf{x}) \leq f(\mathbf{x}_0)$, the convexity of f and the Cauchy-Schwarz inequality it follows, for any $i = 1, 2, \dots, m$ and any \mathbf{x} , that

$$f(\mathbf{a}_i) - f(\mathbf{x}_0) \leq f(\mathbf{a}_i) - f(\mathbf{x}) \leq \langle \mathbf{v}_i, \mathbf{a}_i - \mathbf{x} \rangle \leq \|\mathbf{v}_i\| \|\mathbf{a}_i - \mathbf{x}\|, \tag{24}$$

where $\mathbf{v}_i \in \partial f(\mathbf{a}_i)$. Note that the subdifferential set $\partial f(\mathbf{a}_i)$ can be written as

$$\partial f(\mathbf{a}_i) = \left\{ \sum_{j=1, j \neq i}^m \omega_j \frac{\mathbf{a}_i - \mathbf{a}_j}{\|\mathbf{a}_i - \mathbf{a}_j\|} + \omega_i \mathbf{z}_i : \|\mathbf{z}_i\| \leq 1 \right\}.$$

Therefore, there exists $\tilde{\mathbf{z}}_i$ such that $\|\tilde{\mathbf{z}}_i\| \leq 1$ and

$$\mathbf{v}_i = \sum_{j=1, j \neq i}^m \omega_j \frac{\mathbf{a}_i - \mathbf{a}_j}{\|\mathbf{a}_i - \mathbf{a}_j\|} + \omega_i \tilde{\mathbf{z}}_i.$$

Hence, from the triangle inequality we get

$$\begin{aligned} \|\mathbf{v}_i\| &= \left\| \sum_{j=1, j \neq i}^m \omega_j \frac{\mathbf{a}_i - \mathbf{a}_j}{\|\mathbf{a}_i - \mathbf{a}_j\|} + \omega_i \tilde{\mathbf{z}}_i \right\| \leq \sum_{j=1, j \neq i}^m \omega_j \left\| \frac{\mathbf{a}_i - \mathbf{a}_j}{\|\mathbf{a}_i - \mathbf{a}_j\|} \right\| \\ &\quad + \omega_i \|\tilde{\mathbf{z}}_i\| \leq \sum_{i=1}^m \omega_i = \omega, \end{aligned}$$

which combined with (24), yields the inequality

$$f(\mathbf{a}_i) - f(\mathbf{x}_0) \leq \omega \|\mathbf{a}_i - \mathbf{x}\|.$$

The desired result now follows by dividing the last inequality by ω . □

We can now find an upper bound of the sequence $\{L(\mathbf{x}_k)\}_{k \geq 0}$. This is done in next lemma.

Lemma 8.2 (Upper bound of the sequence $\{L(\mathbf{x}_k)\}_{k \geq 0}$) *Let $\{\mathbf{x}_k\}_{k \geq 0}$ be the sequence generated by Weiszfeld’s method with \mathbf{x}_0 chosen by (20)–(22). Then, for any $k \geq 0$, we have*

$$L(\mathbf{x}_k) \leq \frac{2L(\mathbf{a}_p) \omega^2}{(\|\mathbf{R}_p\| - \omega_p)^2}, \tag{25}$$

where $\omega = \sum_{i=1}^m \omega_i$.

Proof By the monotonicity of the sequence $\{f(\mathbf{x}_k)\}_{k \geq 0}$ (see Corollary 3.1) it follows that $f(\mathbf{x}_k) \leq f(\mathbf{x}_0)$ for all $k \geq 0$. Therefore, from Lemma 8.1 it follows that for any $i \in \{1, 2, \dots, m\}$, we have

$$\frac{1}{\|\mathbf{x}_k - \mathbf{a}_i\|} \leq \frac{\omega}{f(\mathbf{a}_i) - f(\mathbf{x}_0)}.$$

Since $f(\mathbf{x}_0) < f(\mathbf{a}_p) \leq f(\mathbf{a}_i)$ for all $1 \leq i \leq m$, we deduce that

$$\frac{1}{\|\mathbf{x}_k - \mathbf{a}_i\|} \leq \frac{\omega}{f(\mathbf{a}_p) - f(\mathbf{x}_0)}.$$

Thus,

$$L(\mathbf{x}_k) = \sum_{i=1}^m \frac{\omega_i}{\|\mathbf{x}_k - \mathbf{a}_i\|} \leq \sum_{i=1}^m \frac{\omega\omega_i}{f(\mathbf{a}_p) - f(\mathbf{x}_0)} = \frac{\omega^2}{f(\mathbf{a}_p) - f(\mathbf{x}_0)},$$

which, along with Lemma 7.1, implies the desired result. □

Combining Theorem 8.1 and Lemma 8.2, we finally obtain the following rate of convergence result of Weiszfeld’s method.

Theorem 8.2 (Sublinear rate of convergence of the Weiszfeld’s method - Explicit version) *Let $\{\mathbf{x}_k\}_{k \geq 0}$ be the sequence generated by Weiszfeld’s method with \mathbf{x}_0 chosen by (20)–(22). Then, for any $k \geq 0$, we have*

$$f(\mathbf{x}_k) - f^* \leq \frac{M}{2k} \|\mathbf{x}_0 - \mathbf{x}^*\|^2, \tag{26}$$

where $M = 2L(\mathbf{a}_p)\omega^2/(\|\mathbf{R}_p\| - \omega_p)^2$.

9 Acceleration via Optimal Schemes

The question that now arises is whether we can find an acceleration of Weiszfeld’s method with a better theoretical rate of convergence than that of the original Weiszfeld’s method. Recall that, in Sect. 8, we showed that Weiszfeld’s method with a specific choice of a starting point converges in terms of function values to the optimal value in a rate of $O(1/k)$. The natural idea is now to use an accelerated gradient-based scheme in order to assure convergence with the faster rate of $O(1/k^2)$. Unfortunately, accelerated schemes such as Nesterov’s optimal method [35] and FISTA [34] are not monotone, meaning that the sequence of function values is not necessarily nonincreasing. This causes a genuine theoretical difficulty to employ the accelerated schemes on the Fermat–Weber problem, since the monotonicity was a crucial argument in showing that Weiszfeld’s method does not get stuck in anchor points.

We are therefore led to consider an additional and different idea. Instead of bypassing the anchor points, which are the points of nondifferentiability, we will simply eliminate them, by using the idea of *smoothing*.

Given a minimization problem

$$\min \left\{ q(\mathbf{x}) : \mathbf{x} \in \mathbb{R}^d \right\}, \tag{M}$$

where the objective function q is nonsmooth, the idea of smoothing is to replace the objective with a smooth approximation q_μ and solve the problem

$$\min \left\{ q_\mu(\mathbf{x}) : \mathbf{x} \in \mathbb{R}^d \right\}, \tag{M_\mu}$$

where μ is the so-called “smoothing parameter” that controls the level of smoothness and the proximity of the approximation q_μ to q . A more precise definition can be found, for example, in [36]. Some earlier works on smoothing techniques are [37–40]. The more recent works [36,41] show that complexity results can be obtained by employing a fast gradient-based method with a rate of $O(1/k^2)$ on the smooth counterpart (M_μ) . In these works, the smooth problem (M_μ) is not equivalent to the original problem (M) but serves only as an approximation. The complexity result states that, by employing a fast gradient-based method on the smooth problem with a special choice of the smoothing parameter, an ε -optimal solution can be obtained after $O(1/\varepsilon)$ iterations, which corresponds to a method with an $O(1/k)$ rate of convergence. This result is not impressive in the context of the Fermat–Weber problem, for which we have already shown in Sect. 8 that Weiszfeld’s method, with a specially chosen starting point, shares this rate of convergence. The challenge is therefore to present a method with an $O(1/k^2)$ rate of convergence. For that, we present here an *exact smoothing* scheme, in which the original problem (FW) is replaced by a different problem that is equivalent to problem (FW) in the sense that its minimizer is exactly the same as the minimizer of the original problem.

We assume that there is no optimal anchor point (otherwise, as usual, it is trivial to find it). In this case, anchors are necessarily not collinear, implying that the objective function is strictly convex, and there exists a unique optimizer \mathbf{x}^* .

We begin by combining two simple but essential results. First, from Lemma 8.1 we can find a lower bound on the distance of \mathbf{x}^* from each of the anchor points:

$$\|\mathbf{x}^* - \mathbf{a}_i\| \geq \frac{f(\mathbf{a}_i) - f(\mathbf{w})}{\omega}, \quad \forall i \in \{1, 2, \dots, m\}, \tag{27}$$

where \mathbf{w} chosen in the same way as the starting point (20)–(22) is picked. More precisely, we take

$$\mathbf{w} = \mathbf{a}_p + t_p \mathbf{d}_p, \tag{28}$$

where

$$p \in \operatorname{argmin} \{ f(\mathbf{a}_i) : 1 \leq i \leq m \}, \tag{29}$$

and

$$\mathbf{d}_p = -\frac{\mathbf{R}_p}{\|\mathbf{R}_p\|}, \quad t_p = \frac{\|\mathbf{R}_p\| - \omega_p}{L(\mathbf{a}_p)}. \tag{30}$$

Second, we can find a smooth function upper bounding the norm function that coincides with the norm function outside a specified ball. Indeed, let us denote the norm function by $g(\mathbf{x}) := \|\mathbf{x}\|$ and let $r > 0$. Then, we define the following function, which we refer to as a *smooth approximation*

$$g_r(\mathbf{x}) = \begin{cases} \|\mathbf{x}\|, & \|\mathbf{x}\| \geq r, \\ \frac{\|\mathbf{x}\|^2}{2r} + \frac{r}{2}, & \|\mathbf{x}\| < r. \end{cases}$$

The smooth approximation function g_r enjoys two essential properties: (i) it is continuously differentiable with Lipschitz gradient with constant $1/r$, and (ii) it is an upper bound on g .

Lemma 9.1 (Properties of g_r) *Let $r > 0$. Then*

- (i) $g_r(\mathbf{x}) \geq g(\mathbf{x})$ for any $\mathbf{x} \in \mathbb{R}^d$.
- (ii) g_r is continuously differentiable, and its gradient is Lipschitz continuous with constant $1/r$.

Proof (i) Clearly from the definition of g_r that we have to show the result only when $\|\mathbf{x}\| < r$. Indeed, in this case we have the following identity:

$$g_r(\mathbf{x}) - g(\mathbf{x}) = \frac{\|\mathbf{x}\|^2}{2r} + \frac{r}{2} - \|\mathbf{x}\| = \left(\frac{\|\mathbf{x}\|}{\sqrt{2r}} - \sqrt{\frac{r}{2}} \right)^2.$$

Hence $g_r(\mathbf{x}) \geq g(\mathbf{x})$ for any $\mathbf{x} \in \mathbb{R}^d$.

- (ii) First, we will write the gradient of g_r :

$$\nabla g_r(\mathbf{x}) = \begin{cases} \frac{\mathbf{x}}{\|\mathbf{x}\|}, & \|\mathbf{x}\| \geq r \\ \frac{\mathbf{x}}{r}, & \|\mathbf{x}\| < r. \end{cases}$$

Thus, clearly that g_r is continuously differentiable. In order to prove the Lipschitz continuity of ∇g_r , we will use the fact that ∇g_r can be written as

$$\nabla g_r = \frac{1}{r} P_B,$$

where P_B is the orthogonal projection operator onto the closed and convex ball $B = \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\| \leq r\}$. We can now use the fact the orthogonal projection operator P_B is nonexpansive, meaning that it is Lipschitz continuous with constant 1 (see [17, Proposition 2.1.3(c), p. 201]). This proves the desired result. □

Remark 9.1 The function g_r is closely related to the so-called Huber function [42], given by

$$H_r(\mathbf{x}) = \begin{cases} \|\mathbf{x}\| - \frac{r}{2}, & \|\mathbf{x}\| \geq r, \\ \frac{\|\mathbf{x}\|^2}{2r}, & \|\mathbf{x}\| < r. \end{cases}$$

In fact, $g_r(\mathbf{x}) + \frac{r}{2} = H_r(\mathbf{x})$. We can deduce part (ii) of Lemma 9.1 from the known properties of the Huber function (see, e.g., [36,41]), but we have chosen to provide a self-contained proof. In addition, since the Huber function is convex, the convexity of g_r follows.

Now, motivated by the inequalities (27), we will define the following smooth approximation of the Fermat–Weber objective function:

$$f_s(\mathbf{x}) := \sum_{i=1}^m \omega_i g_{b_i}(\mathbf{x} - \mathbf{a}_i),$$

where

$$b_i = \frac{f(\mathbf{a}_i) - f(\mathbf{w})}{\omega}, \quad i = 1, 2, \dots, m.$$

Now we have the following lemma stating that f_s is an exact convex smoothing counterpart of f .

Lemma 9.2 (Properties of the exact smoothing function f_s) *Let \mathbf{w} be defined by (28)–(30). The following properties hold:*

- (i) f_s is convex over \mathbb{R}^d .
- (ii) $f_s(\mathbf{x}) \geq f(\mathbf{x})$ for any $\mathbf{x} \in \mathbb{R}^d$.
- (iii) The optimal solution of the problem

$$\min_{\mathbf{x} \in \mathbb{R}^d} f_s(\mathbf{x})$$

is \mathbf{x}^* - the optimal solution of problem (FW).

- (iv) f_s is continuously differentiable, and its gradient is Lipschitz continuous with constant

$$L_s = \omega \sum_{i=1}^m \frac{\omega_i}{f(\mathbf{a}_i) - f(\mathbf{w})}.$$

Moreover, the Lipschitz constant L_s can be bounded from above by

$$L_s \leq \frac{2L(\mathbf{a}_p)\omega^2}{(\|\mathbf{R}_p\| - \omega_p)^2}.$$

Proof (i) Follows by the fact that f_s is the weighted sum of the convex functions g_{b_i} , $i = 1, 2, \dots, m$.

- (ii) First, from Lemma 9.1(i), it follows that $g_{b_i}(\mathbf{x}) \geq g(\mathbf{x})(= \|\mathbf{x}\|)$ for all $\mathbf{x} \in \mathbb{R}^d$. Hence, for any $\mathbf{x} \in \mathbb{R}^d$:

$$f_s(\mathbf{x}) = \sum_{i=1}^m \omega_i g_{b_i}(\mathbf{x} - \mathbf{a}_i) \geq \sum_{i=1}^m \omega_i g(\mathbf{x} - \mathbf{a}_i) = \sum_{i=1}^m \omega_i \|\mathbf{x} - \mathbf{a}_i\| = f(\mathbf{x}).$$

- (iii) From (27), the inequality $\|\mathbf{x}^* - \mathbf{a}_i\| \geq b_i$ holds for any $1 \leq i \leq m$ and hence $g_{b_i}(\mathbf{x}^* - \mathbf{a}_i) = \|\mathbf{x}^* - \mathbf{a}_i\|$ for any $1 \leq i \leq m$. Consequently,

$$f_s(\mathbf{x}^*) = \sum_{i=1}^m \omega_i g_{b_i}(\mathbf{x}^* - \mathbf{a}_i) = \sum_{i=1}^m \omega_i \|\mathbf{x}^* - \mathbf{a}_i\| = f(\mathbf{x}^*).$$

Let $\mathbf{x} \in \mathbb{R}^d$ be different from than \mathbf{x}^* . Since \mathbf{x}^* is the strict global minimum of f over \mathbb{R}^d , it follows that $f(\mathbf{x}^*) < f(\mathbf{x})$. Hence,

$$f_s(\mathbf{x}^*) = f(\mathbf{x}^*) < f(\mathbf{x}) \leq f_s(\mathbf{x}),$$

where the last inequality follows from part (ii). We therefore conclude that \mathbf{x}^* is also the unique minimizer of f_s over \mathbb{R}^d .

- (iv) For each $1 \leq i \leq m$, by Lemma 9.1(ii) it follows that g_{b_i} is continuously differentiable with gradient which is Lipschitz with constant $1/b_i$. Therefore, by its definition, the function f_s is continuously differentiable with gradient which is Lipschitz continuous with constant

$$\sum_{i=1}^m \frac{\omega_i}{b_i} = \omega \sum_{i=1}^m \frac{\omega_i}{f(\mathbf{a}_i) - f(\mathbf{w})}.$$

Since $f(\mathbf{a}_i) \geq f(\mathbf{a}_p)$ for all $1 \leq i \leq m$ (follows from (29)), we can estimate the Lipschitz constant by

$$\omega \sum_{i=1}^m \frac{\omega_i}{f(\mathbf{a}_i) - f(\mathbf{w})} \leq \omega \sum_{i=1}^m \frac{\omega_i}{f(\mathbf{a}_p) - f(\mathbf{w})} = \frac{\omega^2}{f(\mathbf{a}_p) - f(\mathbf{w})} \leq \frac{2L(\mathbf{a}_p)\omega^2}{(\|\mathbf{R}_p\| - \omega_p)^2},$$

where the last inequality follows from Lemma 7.1.

□

The effect of the smoothing operation is illustrated in Fig. 3.

Now we know that the nonsmooth Fermat–Weber problem can be replaced by the smooth counterpart

$$\min_{\mathbf{x} \in \mathbb{R}^d} f_s(\mathbf{x}), \tag{FW_s}$$

which has the same optimal solution. We can invoke one of the optimal gradient-based methods for solving smooth convex optimization problems. One of the simplest options is Nesterov’s method from 1983 [35]. We begin by explicitly writing the accelerated method with a constant stepsize version.

Fast Weiszfeld Method (constant stepsize)

Initialization. $\mathbf{y}_1 = \mathbf{x}_0 \in \mathbb{R}^d$ and $t_1 = 1$.

General Step ($k = 1, 2, \dots$) Compute

$$\begin{aligned} \mathbf{x}_k &= \mathbf{y}_k - \frac{1}{L_s} \nabla f_s(\mathbf{y}_k), \\ t_{k+1} &= \frac{1 + \sqrt{1 + 4t_k^2}}{2}, \\ \mathbf{y}_{k+1} &= \mathbf{x}_k + \left(\frac{t_k - 1}{t_{k+1}} \right) (\mathbf{x}_k - \mathbf{x}_{k-1}). \end{aligned}$$




Fig. 3 The left image describes the contour lines of the unweighted Fermat–Weber function with anchors $\mathbf{a}_1 = (1, 2)^T$, $\mathbf{a}_2 = (2, 1)^T$, and $\mathbf{a}_3 = (5, 5)^T$. The anchors are denoted by the filled squares, while the empty square stands for the optimal solution. In the right image, the contour lines of a smoothed function are given. Here we have chosen to replace each of the norm functions $g(\mathbf{x}) = \|\mathbf{x}\|$ by the smooth counterpart $g_{1/2}(\mathbf{x})$

A possible drawback of this basic fast scheme is that the Lipschitz constant L_s can be too conservative (*i.e.*, large). We therefore also consider a version with a backtracking stepsize rule. This version was considered in the context of the more general composite model [34].

Fast Weiszfeld Method (backtracking stepsize)

Initialization. $L_0 > 0$, some $\eta > 1$, and $\mathbf{x}_0 \in \mathbb{R}^d$. Set $\mathbf{y}_1 = \mathbf{x}_0$ and $t_1 = 1$.

General Step ($k = 1, 2, \dots$) Find the smallest non-negative integer i_k such that, with $\bar{L} = \eta^{i_k} L_{k-1}$,

$$f_s \left(\mathbf{y}_k - \frac{1}{\bar{L}} \nabla f_s(\mathbf{y}_k) \right) \leq f_s(\mathbf{y}_k) - \frac{1}{2\bar{L}} \|\nabla f_s(\mathbf{y}_k)\|^2.$$

Set $L_k = \eta^{i_k} L_{k-1}$ and compute

$$\begin{aligned} \mathbf{x}_k &= \mathbf{y}_k - \frac{1}{L_k} \nabla f_s(\mathbf{y}_k), \\ t_{k+1} &= \frac{1 + \sqrt{1 + 4t_k^2}}{2}, \\ \mathbf{y}_{k+1} &= \mathbf{x}_k + \left(\frac{t_k - 1}{t_{k+1}} \right) (\mathbf{x}_k - \mathbf{x}_{k-1}). \end{aligned}$$

The following convergence result for both the constant stepsize scheme as well as the backtracking version was proved in [34].

Theorem 9.1 (Convergence of the fast Weiszfeld method) *Let $\{\mathbf{x}_k\}_{k \geq 0}$ and $\{\mathbf{y}_k\}_{k \geq 0}$ be two sequences which are generated by one of the fast Weiszfeld method. Then, for any $k \geq 1$, we have*

$$f_s(\mathbf{x}_k) - f^* \leq \frac{2\alpha L_s \|\mathbf{x}_0 - \mathbf{x}^*\|^2}{(k + 1)^2},$$

where $\alpha = 1$ for the constant stepsize setting and $\alpha = \eta$ for the backtracking stepsize setting.

Since $f_s(\mathbf{x}) \geq f(\mathbf{x})$ for any $\mathbf{x} \in \mathbb{R}^d$, we can immediately conclude an $O(1/k^2)$ rate of convergence result of the original objective function to the optimal value.

Corollary 9.1 (Convergence of the fast Weiszfeld’s method - original function values) *Let $\{\mathbf{x}_k\}_{k \geq 0}$ and $\{\mathbf{y}_k\}_{k \geq 0}$ be two sequences which are generated by one of the fast Weiszfeld methods. Then, for any $k \geq 1$, we have*

$$f(\mathbf{x}_k) - f^* \leq \frac{2\alpha L_s \|\mathbf{x}_0 - \mathbf{x}^*\|^2}{(k + 1)^2},$$

where $\alpha = 1$ for the constant stepsize setting and $\alpha = \eta$ for the backtracking stepsize setting.

10 Extensions, Open Questions and Perspectives

Weiszfeld’s method had an impact on the development of many numerical methods for solving various problems, and its influence was not restricted to the Fermat–Weber problem. Perhaps the most natural generalization of the Fermat–Weber problem (and Weiszfeld’s method as well) is to the problem of multi-facility location. In fact, one of the first papers that dealt with Weiszfeld’s method (without knowing it...), was Miehle’s paper from 1958 [15], where he considered a problem of finding the locations of two points in \mathbb{R}^2 . The general multi-facility location problem was considered by Radó in [43] (1988), where the problem was formulated as

$$\min \sum_{j=1}^n \sum_{i=1}^m w_{ji} \|\mathbf{x}_j - \mathbf{a}_i\| + \sum_{j=1}^n \sum_{\ell=1}^n v_{j\ell} \|\mathbf{x}_j - \mathbf{x}_\ell\|.$$

Here, $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m \in \mathbb{R}^d$ are the fixed anchors, and $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n \in \mathbb{R}^d$ are the n locations that we seek to find. The nonnegative numbers w_{ji} and v_{jk} are given weights. Radó considered the following generalization of Weiszfeld’s method. To construct the $(k + 1)$ -th iterate from the k -th iterate, like in Weiszfeld’s method, each of the norm expressions $\|\mathbf{x}\|$ is replaced by the term $\|\mathbf{x}\|^2 / \|\mathbf{x}^k\|$. That is, at the k -th iteration the following minimization problem is solved:

$$\min_{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n} \sum_{j=1}^n \sum_{i=1}^m w_{ji} \frac{\|\mathbf{x}_j - \mathbf{a}_i\|^2}{\|\mathbf{x}_i^k - \mathbf{a}_j\|} + \sum_{j=1}^n \sum_{\ell=1}^n v_{j\ell} \frac{\|\mathbf{x}_j - \mathbf{x}_\ell\|^2}{\|\mathbf{x}_j^k - \mathbf{x}_\ell^k\|}.$$

The solution to the above convex problem, which is next iterate $\mathbf{x}_1^{k+1}, \mathbf{x}_2^{k+1}, \dots, \mathbf{x}_n^{k+1}$, is attained at its unique stationary point, which is the solution of following linear system of equations:

$$\sum_{i=1}^m w_{ji} \frac{\mathbf{x}_j - \mathbf{a}_i}{\|\mathbf{x}_i^k - \mathbf{a}_j\|} + \sum_{j=1}^n \sum_{\ell=1}^n v_{j\ell} \frac{\mathbf{x}_j - \mathbf{x}_\ell}{\|\mathbf{x}_j^k - \mathbf{x}_\ell^k\|} = 0, \quad j = 1, 2, \dots, n.$$

Some convergence properties of this generalized Weiszfeld's method were studied in [43]. Another type of a multi-facility location problem that involves also clustering, was considered by Iyigun and Ben-Israel in [44] (2010), and also by Teboulle in [45] (2007). The authors constructed a generalization of Weiszfeld's method and found several properties such as monotonicity; of course, since the problem is nonconvex, it does not seem possible to construct a method that is guaranteed to converge to the global optimal solution.

Another type of generalizations of the Fermat–Weber problem leading to corresponding generalizations of Weiszfeld's method is concerned with replacing the Euclidean norm in the objective function by another norm. For example, in [46] (2010) Katz and Vogl considered a Weiszfeld method which solves a version of the Fermat–Weber problem in which the Euclidean norms are replaced with weighted Euclidean norms. Cooper extended Weiszfeld method in [47] (1968) to consider a Fermat–Weber type problem in which the objective is to minimize a weighted sum of *powers* of the Euclidean norms:

$$\min_{\mathbf{x} \in \mathbb{R}^d} \sum_{i=1}^n \omega_i \|\mathbf{x} - \mathbf{a}_i\|^K,$$

where $K > 0$. Of course, the problem is convex only when $K \geq 1$. This extension of the problem was also studied by Chen in [24] (1984). An extension of both the Fermat–Weber problem and the Weiszfeld's method for p -norms was considered by Morris in [48] (1981).

We would also like to point out a known generalization of the Fermat–Weber problem which was suggested by Erdős in [49] for the case when $m = d + 1$ and when $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_{d+1}$ are affinely independent. The problem Erdős considered takes the form

$$\min \frac{\sum_{i=1}^{d+1} \|\mathbf{x} - \mathbf{a}_i\|}{\sum_{i=1}^{d+1} \text{dist}(\mathbf{x}, F_i)},$$

where F_1, F_2, \dots, F_{d+1} are the $(d - 1)$ -dimensional facets of the simplex $\text{conv}\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_{d+1}\}$. As opposed to the Fermat–Weber problem, this generalization is a non-convex problem and is therefore in principal difficult to solve.

A totally different location problem, in which Weiszfeld-type ideas were used, is the *source localization problem*. In this problem we are given measurements of the distances of a source in an unknown location \mathbf{x} from the m anchors:

$$\|\mathbf{x} - \mathbf{a}_i\| \approx d_i, \quad i = 1, 2, \dots, m.$$

One formulation of the problem consists in finding the \mathbf{x} resulting with the minimum sum of squared errors:

$$\min_{\mathbf{x} \in \mathbb{R}^d} \sum_{i=1}^m (\|\mathbf{x} - \mathbf{a}_i\| - d_i)^2. \tag{SL}$$

Despite its apparent resemblance to Fermat–Weber, problem (SL) is quite different in the sense that it is a difficult nonconvex problem. One of the methods studied by Beck et al in [50] (2008) uses the idea of Weiszfeld to replace the norm terms of the form $\|\mathbf{x} - \mathbf{a}_i\|$ with the expressions $\|\mathbf{x} - \mathbf{a}_i\|^2 / \|\mathbf{x}_k - \mathbf{a}_i\|$, thus resulting with the following iterative scheme:

$$\mathbf{x}_{k+1} \in \operatorname{argmin}_{\mathbf{x} \in \mathbb{R}^d} \sum_{i=1}^m \left(\frac{\|\mathbf{x} - \mathbf{a}_i\|^2}{\|\mathbf{x}_k - \mathbf{a}_i\|} - d_i \right)^2.$$

The convergence of this scheme (to stationary points) was studied in [50]. An open question in this context is whether this approach can be extended to the more general sensor network localization problem where we are given a sensor network with m anchors and n sensors. The locations of the anchors are given by the *known* vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m \in \mathbb{R}^d$; the locations of the n sensors are decision variables vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n \in \mathbb{R}^d$. We assume that we are given noisy measurements of some of the distances between pairs of sensors and between pairs of sensors and anchors:

$$\|\mathbf{x}_j - \mathbf{x}_t\| \approx d_{jt}, \quad (j, t) \in \mathcal{N}, \tag{31}$$

$$\|\mathbf{x}_j - \mathbf{a}_i\| \approx w_{ji}, \quad (j, i) \in \mathcal{M}, \tag{32}$$

where

$$\begin{aligned} \mathcal{N} &\subseteq \{(j, t) : j \neq t, j, t = 1, 2, \dots, n\}, \\ \mathcal{M} &\subseteq \{(j, i) : i = 1, 2, \dots, m, j = 1, 2, \dots, n\}, \end{aligned}$$

are the subsets of pairs of indices corresponding to the sensor/sensor and anchor/sensor distance measurements. We assume that, if $(j, t) \in \mathcal{N}$ for some $j \neq t$, then $(t, j) \notin \mathcal{N}$. A possible modeling of the problem (see *e.g.*, Biswas et al [51]) is via the minimization problem:

$$\min_{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n} \left\{ f(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) \equiv \sum_{(j,t) \in \mathcal{N}} (\|\mathbf{x}_j - \mathbf{x}_t\| - d_{jt})^2 + \sum_{(j,i) \in \mathcal{M}} (\|\mathbf{x}_j - \mathbf{a}_i\| - w_{ji})^2 \right\}. \tag{SNL}$$

Problem (SNL) is a nonconvex problem, and hence finding its global optimal solution is generally speaking a difficult task. The question that arises is whether the Weiszfeld-type techniques used in [50] can be used to construct an efficient solution method. Obviously, since the problem is nonconvex, the main objective from a theoretical point of view is to prove convergence or rate of convergence to a stationary point.

We also note that *the iteratively reweighted least squares method*, which is one of the most popular optimization algorithms for solving a wide variety of problems involving norms, is essentially based on ideas from Weiszfeld’s method. As an illustration of the method, consider the problem of robust regression which consists of solving the l_1 -norm problem

$$\min_{\mathbf{x} \in \mathbb{R}^d} \left\{ \|\mathbf{Ax} - \mathbf{b}\|_1 = \sum_{i=1}^m |\mathbf{a}_i^T \mathbf{x} - b_i| \right\},$$

where $\mathbf{A} \in \mathbb{R}^{m \times n}$, $\mathbf{b} \in \mathbb{R}^m$, and $\mathbf{a}_1^T, \mathbf{a}_2^T, \dots, \mathbf{a}_m^T$ are the rows of the matrix \mathbf{A} . At each iteration, the absolute values $|\mathbf{a}_i^T \mathbf{x} - b_i|$ are replaced by $(\mathbf{a}_i^T \mathbf{x} - b_i)^2 / |\mathbf{a}_i^T \mathbf{x}_k - b_i|$, and the $(k + 1)$ -th is determined from the k -th iteration by the update formulas:

$$\mathbf{x}_{k+1} \in \operatorname{argmin} \left\{ \sum_{i=1}^m \frac{(\mathbf{a}_i^T \mathbf{x} - b_i)^2}{|\mathbf{a}_i^T \mathbf{x}_k - b_i|} \right\}.$$

This scheme was studied in the context of various types of models, and the literature of this method covers hundreds of paper that were written in the past 60 years; the list of references [52–57] is just a very small representative sample of works dealing with various applications and theoretical properties.

We end this section by recalling that the Fermat–Weber problem is associated with what is considered to be the oldest example of constrained extremum problems duality in the literature (see the paper of Kuhn [58] for a historical account). Here we consider the original problem of Fermat with three points in the plane, and without weights. The primal and dual problems are

- **Primal.** Fermat–Weber problem: given three points in the plane, $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$, find a point $\mathbf{x} \in \mathbb{R}^2$ minimizing the sum of distances to $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$.
- **Dual.** Find the equilateral triangle with maximal altitude circumscribing the triangle with vertices $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$.

The optimal values of the two problems are actually the same, that is, the altitude of the largest equilateral triangle circumscribing the given triangle is equal to the sum of the distances of the Fermat–Torricelli point from the three vertices. This duality is also called *Fasbender’s duality* since it was discovered by Fasbender [59] in 1846. As was pointed out in [58], this duality is essentially equivalent to Lagrangian duality. For the general Fermat–Weber problem (FW), the Lagrangian dual problem is given by

$$\max_{\mathbf{u}_1, \dots, \mathbf{u}_m \in \mathbb{R}^d} \left\{ \sum_{i=1}^m \mathbf{a}_i^T \mathbf{u}_i : \sum_{i=1}^m \mathbf{u}_i = \mathbf{0}, \|\mathbf{u}_j\| \leq \omega_j, j = 1, 2, \dots, m \right\}. \quad (\text{D})$$

For the original Fermat–Torricelli problem ($m = 3, \omega_i = 1, n = 2$), when all the angles are smaller than 120° , the relation between the optimal dual variables $(\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3)$ and the primal variables vector (\mathbf{x}) is given by $\mathbf{u}_i = \frac{\mathbf{a}_i - \mathbf{x}}{\|\mathbf{a}_i - \mathbf{x}\|}$. In addition, the i -th side of the largest circumscribing equilateral triangle passes through \mathbf{a}_i and is perpendicular to \mathbf{u}_i . An interesting line of research will be to understand the Weiszfeld’s method from the point of view duality. Thus, it will be interesting to derive a dual form of Weiszfeld’s method, and to find generalizations of this dual method that are able to cope with models different than (D).

11 Conclusions

In this paper we reviewed the intriguing story of Weiszfeld’s method beginning from its development in 1937. All the convergence results were presented in a self-contained manner, and some of the proofs are new and simplified. Two new results were derived: the first is a nonasymptotic rate of convergence of the sequence of function values generated by Weiszfeld’s method, and the second is an acceleration of the method based on an exact smoothed formulation and an optimal gradient-based method.

Acknowledgments We would like to thank an anonymous reviewer and the editor in chief for their useful comments which helped to improve the presentation of the paper. This work was partially supported by ISF grant #25312.

12 Appendix A: Notations

Following is a list of notations that are used throughout the paper.

- $\mathcal{A} = \{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m\}$ —the set of anchors.
- $\omega_1, \omega_2, \dots, \omega_m$ —given positive weights.
- $\omega = \sum_{i=1}^m \omega_i$ —sum of weights.
- $f(\mathbf{x}) = \sum_{i=1}^m \omega_i \|\mathbf{x} - \mathbf{a}_i\|$ —the Fermat–Weber objective function.
- \mathbf{x}^* —an optimal solution of the Fermat–Weber problem. If the anchors are not collinear, then \mathbf{x}^* is the *unique* optimal solution.
- X^* —the optimal solution set of the Fermat–Weber problem. When the anchors are not collinear, X^* is the singleton $\{\mathbf{x}^*\}$.
- f^* —the optimal value of the Fermat–Weber problem.
- $T(\mathbf{x}) = \frac{1}{\sum_{i=1}^m \frac{\omega_i}{\|\mathbf{x} - \mathbf{a}_i\|}} \sum_{i=1}^m \frac{\omega_i \mathbf{a}_i}{\|\mathbf{x} - \mathbf{a}_i\|}$ —the operator defining Weiszfeld’s method.
- $h(\mathbf{x}, \mathbf{y}) := \sum_{i=1}^m \omega_i \frac{\|\mathbf{x} - \mathbf{a}_i\|^2}{\|\mathbf{y} - \mathbf{a}_i\|}$ —an auxiliary function used to analyze Weiszfeld’s method.
- $L(\mathbf{x}) = \sum_{i=1}^m \frac{\omega_i}{\|\mathbf{x} - \mathbf{a}_i\|}$ —serves as a kind of “Lipschitz” constant, and the operator T can be written as taking a gradient step with stepsize $1/L(\mathbf{x})$: $T(\mathbf{x}) = \mathbf{x} - 1/L(\mathbf{x}) \nabla f(\mathbf{x})$.
- $\mathbf{R}_j = \sum_{i=1, i \neq j}^m \omega_i (\mathbf{a}_j - \mathbf{a}_i) / \|\mathbf{a}_i - \mathbf{a}_j\|, j = 1, 2, \dots, m$. An important property related to \mathbf{R}_j is that \mathbf{a}_j is optimal if and only if $\|\mathbf{R}_j\| \leq \omega_j$.
- $\mathbf{d}_j = -\mathbf{R}_j / \|\mathbf{R}_j\|$ —the steepest descent direction of f at \mathbf{a}_j .

13 Appendix B: Proof of Lemma 7.1

Let $\mathbf{x} := \mathbf{a}_j + t_j \mathbf{d}_j$. Then, from the definition of \mathbf{x} , we have that $\mathbf{d}_j = (1/t_j) (\mathbf{x} - \mathbf{a}_j)$ and hence

$$\begin{aligned} -\frac{1}{t_j} \|\mathbf{x} - \mathbf{a}_j\|^2 &= \frac{1}{t_j} \|\mathbf{x} - \mathbf{a}_j\|^2 - 2 \left\langle \mathbf{x} - \mathbf{a}_j, \frac{\mathbf{x} - \mathbf{a}_j}{t_j} \right\rangle \\ &= \frac{1}{t_j} \|\mathbf{x} - \mathbf{a}_j\|^2 - 2 \langle \mathbf{x} - \mathbf{a}_j, \mathbf{d}_j \rangle \\ &= \frac{1}{t_j} \|\mathbf{x} - \mathbf{a}_j\|^2 - 2 \left\langle \mathbf{x} - \mathbf{a}_j, -\frac{\mathbf{R}_j}{\|\mathbf{R}_j\|} \right\rangle \\ &= \frac{1}{t_j} \|\mathbf{x} - \mathbf{a}_j\|^2 + \frac{2}{\|\mathbf{R}_j\|} \langle \mathbf{x} - \mathbf{a}_j, \mathbf{R}_j \rangle. \end{aligned} \tag{33}$$

Now, we will expand the first term of the right-hand side of (33)

$$\begin{aligned} \frac{1}{t_j} \|\mathbf{x} - \mathbf{a}_j\|^2 &= \frac{L(\mathbf{a}_j)}{\|\mathbf{R}_j\| - \omega_j} \|\mathbf{x} - \mathbf{a}_j\|^2 \\ &= \frac{1}{\|\mathbf{R}_j\| - \omega_j} \left(\sum_{i=1, i \neq j}^m \frac{\omega_i}{\|\mathbf{a}_j - \mathbf{a}_i\|} \right) \|\mathbf{x} - \mathbf{a}_j\|^2 \\ &= \frac{1}{\|\mathbf{R}_j\| - \omega_j} \sum_{i=1, i \neq j}^m \omega_i \frac{\|\mathbf{x} - \mathbf{a}_j\|^2}{\|\mathbf{a}_j - \mathbf{a}_i\|} \\ &= \frac{1}{\|\mathbf{R}_j\| - \omega_j} \sum_{i=1, i \neq j}^m \omega_i \left(\frac{\|\mathbf{x} - \mathbf{a}_i\|^2}{\|\mathbf{a}_j - \mathbf{a}_i\|} + 2 \frac{\langle \mathbf{x} - \mathbf{a}_i, \mathbf{a}_i - \mathbf{a}_j \rangle}{\|\mathbf{a}_j - \mathbf{a}_i\|} + \frac{\|\mathbf{a}_i - \mathbf{a}_j\|^2}{\|\mathbf{a}_j - \mathbf{a}_i\|} \right) \\ &= \frac{1}{\|\mathbf{R}_j\| - \omega_j} \sum_{i=1, i \neq j}^m \omega_i \left(\frac{\|\mathbf{x} - \mathbf{a}_i\|^2}{\|\mathbf{a}_j - \mathbf{a}_i\|} + 2 \frac{\langle \mathbf{x} - \mathbf{a}_i, \mathbf{a}_i - \mathbf{a}_j \rangle}{\|\mathbf{a}_j - \mathbf{a}_i\|} + \|\mathbf{a}_j - \mathbf{a}_i\| \right). \end{aligned}$$

The middle term can be also written as follows

$$\begin{aligned} 2 \frac{\langle \mathbf{x} - \mathbf{a}_i, \mathbf{a}_i - \mathbf{a}_j \rangle}{\|\mathbf{a}_j - \mathbf{a}_i\|} &= 2 \frac{\langle \mathbf{x} - \mathbf{a}_j, \mathbf{a}_i - \mathbf{a}_j \rangle}{\|\mathbf{a}_j - \mathbf{a}_i\|} + 2 \frac{\langle \mathbf{a}_j - \mathbf{a}_i, \mathbf{a}_i - \mathbf{a}_j \rangle}{\|\mathbf{a}_j - \mathbf{a}_i\|} \\ &= 2 \frac{\langle \mathbf{x} - \mathbf{a}_j, \mathbf{a}_i - \mathbf{a}_j \rangle}{\|\mathbf{a}_j - \mathbf{a}_i\|} - 2 \|\mathbf{a}_j - \mathbf{a}_i\|. \end{aligned}$$

Thus, from the definition of \mathbf{R}_j , we have

$$\frac{1}{t_j} \|\mathbf{x} - \mathbf{a}_j\|^2 = \frac{1}{\|\mathbf{R}_j\| - \omega_j} \sum_{i=1, i \neq j}^m \omega_i \left(\frac{\|\mathbf{x} - \mathbf{a}_i\|^2}{\|\mathbf{a}_j - \mathbf{a}_i\|} + 2 \frac{\langle \mathbf{x} - \mathbf{a}_j, \mathbf{a}_i - \mathbf{a}_j \rangle}{\|\mathbf{a}_j - \mathbf{a}_i\|} - \|\mathbf{a}_j - \mathbf{a}_i\| \right)$$

$$\begin{aligned}
 &= \frac{1}{\|\mathbf{R}_j\| - \omega_j} \left(\sum_{i=1, i \neq j}^m \omega_i \frac{\|\mathbf{x} - \mathbf{a}_i\|^2}{\|\mathbf{a}_j - \mathbf{a}_i\|} \right. \\
 &\quad \left. + 2 \left\langle \mathbf{x} - \mathbf{a}_j, \sum_{i=1, i \neq j}^m \omega_i \frac{\mathbf{a}_i - \mathbf{a}_j}{\|\mathbf{a}_j - \mathbf{a}_i\|} \right\rangle - f(\mathbf{a}_j) \right) \\
 &= \frac{1}{\|\mathbf{R}_j\| - \omega_j} \left(\sum_{i=1, i \neq j}^m \omega_i \frac{\|\mathbf{x} - \mathbf{a}_i\|^2}{\|\mathbf{a}_j - \mathbf{a}_i\|} - 2 \langle \mathbf{x} - \mathbf{a}_j, \mathbf{R}_j \rangle - f(\mathbf{a}_j) \right).
 \end{aligned}$$

Note that, by the fact that $\frac{a^2}{b} \geq 2a - b$ for any $a \in \mathbb{R}$ and $b \in \mathbb{R}_{++}$ we have

$$\frac{\|\mathbf{x} - \mathbf{a}_i\|^2}{\|\mathbf{a}_j - \mathbf{a}_i\|} \geq 2\|\mathbf{x} - \mathbf{a}_i\| - \|\mathbf{a}_j - \mathbf{a}_i\|.$$

Hence

$$\begin{aligned}
 \frac{1}{t_j} \|\mathbf{x} - \mathbf{a}_j\|^2 &\geq \frac{1}{\|\mathbf{R}_j\| - \omega_j} \left(\sum_{i=1, i \neq j}^m \omega_i (2\|\mathbf{x} - \mathbf{a}_i\| - \|\mathbf{a}_j - \mathbf{a}_i\|) \right. \\
 &\quad \left. - 2 \langle \mathbf{x} - \mathbf{a}_j, \mathbf{R}_j \rangle - f(\mathbf{a}_j) \right) \\
 &= \frac{1}{\|\mathbf{R}_j\| - \omega_j} \left(2 \sum_{i=1, i \neq j}^m \omega_i \|\mathbf{x} - \mathbf{a}_i\| - 2 \langle \mathbf{x} - \mathbf{a}_j, \mathbf{R}_j \rangle - 2f(\mathbf{a}_j) \right) \\
 &= \frac{2}{\|\mathbf{R}_j\| - \omega_j} \left(\sum_{i=1, i \neq j}^m \omega_i \|\mathbf{x} - \mathbf{a}_i\| - \langle \mathbf{x} - \mathbf{a}_j, \mathbf{R}_j \rangle - f(\mathbf{a}_j) \right).
 \end{aligned}$$

Plugging the last inequality in (33) yields

$$\begin{aligned}
 -\frac{1}{t_j} \|\mathbf{x} - \mathbf{a}_j\|^2 &= \frac{1}{t_j} \|\mathbf{x} - \mathbf{a}_j\|^2 + \frac{2}{\|\mathbf{R}_j\|} \langle \mathbf{x} - \mathbf{a}_j, \mathbf{R}_j \rangle \\
 &\geq \frac{2}{\|\mathbf{R}_j\| - \omega_j} \left(\sum_{i=1, i \neq j}^m \omega_i \|\mathbf{x} - \mathbf{a}_i\| - \langle \mathbf{x} - \mathbf{a}_j, \mathbf{R}_j \rangle - f(\mathbf{a}_j) \right) \\
 &\quad + \frac{2}{\|\mathbf{R}_j\|} \langle \mathbf{x} - \mathbf{a}_j, \mathbf{R}_j \rangle \\
 &= \frac{2}{\|\mathbf{R}_j\| - \omega_j} \left(\sum_{i=1, i \neq j}^m \omega_i \|\mathbf{x} - \mathbf{a}_i\| - f(\mathbf{a}_j) \right) \\
 &\quad - \left(\frac{2}{\|\mathbf{R}_j\| - \omega_j} - \frac{2}{\|\mathbf{R}_j\|} \right) \langle \mathbf{x} - \mathbf{a}_j, \mathbf{R}_j \rangle
 \end{aligned}$$

$$\begin{aligned}
&= \frac{2}{\|\mathbf{R}_j\| - \omega_j} \left(\sum_{i=1, i \neq j}^m \omega_i \|\mathbf{x} - \mathbf{a}_i\| - f(\mathbf{a}_j) \right) \\
&\quad - \frac{2\omega_j}{\|\mathbf{R}_j\| (\|\mathbf{R}_j\| - \omega_j)} \langle \mathbf{x} - \mathbf{a}_j, \mathbf{R}_j \rangle \\
&= \frac{2}{\|\mathbf{R}_j\| - \omega_j} \left(\sum_{i=1, i \neq j}^m \omega_i \|\mathbf{x} - \mathbf{a}_i\| - f(\mathbf{a}_j) \right) \\
&\quad + \frac{2\omega_j}{\|\mathbf{R}_j\| - \omega_j} \langle \mathbf{x} - \mathbf{a}_j, \mathbf{d}_j \rangle \\
&= \frac{2}{\|\mathbf{R}_j\| - \omega_j} \left(\sum_{i=1, i \neq j}^m \omega_i \|\mathbf{x} - \mathbf{a}_i\| - f(\mathbf{a}_j) + \omega_j \langle \mathbf{x} - \mathbf{a}_j, \mathbf{d}_j \rangle \right) \\
&= \frac{2}{\|\mathbf{R}_j\| - \omega_j} \left(\sum_{i=1, i \neq j}^m \omega_i \|\mathbf{x} - \mathbf{a}_i\| - f(\mathbf{a}_j) + \omega_j \|\mathbf{x} - \mathbf{a}_j\| \right) \\
&= \frac{2}{\|\mathbf{R}_j\| - \omega_j} (f(\mathbf{x}) - f(\mathbf{a}_j)),
\end{aligned}$$

where the second equality from below follows from the fact that $1 = \|\mathbf{d}_j\| = \|\mathbf{x} - \mathbf{a}_j\| / t_j$. Hence,

$$f(\mathbf{a}_j) - f(\mathbf{x}) \geq \frac{\|\mathbf{R}_j\| - \omega_j}{2t_j} \|\mathbf{x} - \mathbf{a}_j\|^2 = t_j \frac{\|\mathbf{R}_j\| - \omega_j}{2} = \frac{(\|\mathbf{R}_j\| - \omega_j)^2}{2L(\mathbf{a}_j)},$$

where the first equality follows from the fact that $\|\mathbf{x} - \mathbf{a}_j\| = t_j$, and the last equality follows from the definition of t_j . \square

References

1. Eiselt, H.A., Marianov, V.: Pioneering developments in location analysis. In: Eiselt, H.A., Marianov, V. (eds.) *Foundations of Location Analysis*, International Series in Operations Research & Management Science, vol. 155, pp. 3–22. Springer, Berlin (2011)
2. Drezner, Z., Klamroth, K., Schöbel, A., Wesolowsky, G.O.: The Weber problem. In: Drezner, Z., Hamacher, H.W. (eds.) *Facility Location: Applications and Theory*, International Series in Operations Research & Management Science, pp. 1–36. Springer, Berlin (2002)
3. Boltyanski, Y., Martini, H., Soltan, V.: *Geometric Methods and Optimization Problems*, Combinatorial Optimization, vol. 4. Kluwer Academic Publishers, Dordrecht (1999)
4. Jahn, T., Kupitz, Y.S., Martini, H., Richter, C.: The Fermat-Torricelli problem, Part II: Geometric aspects, open and related problems, and new generalizations. *J. Optim. Theory Appl.* (to appear)
5. Weiszfeld, E.V.: Sur le point pour lequel la somme des distances de n points données est minimum. *Tohoku Math. J.* **43**, 355–386 (1937)
6. Vazsonyi, A.: Pure mathematics and the Weiszfeld algorithm. *Decis. Line* **33**, 12–13 (2002)
7. Sturm, R.: Ueber den Punkt kleinster Entfernungssumme von gegebenen Punkten. *J. Rein. Angew. Math.* **97**, 49–61 (1884)

8. Ostresh, L.M.: On the convergence of a class of iterative methods for solving the Weber location problem. *Oper. Res.* **26**, 597–609 (1978)
9. Bazaraa, M.S., Sherali, H.D., Shetty, C.M.: *Nonlinear Programming: Theory and Algorithms*, 3rd edn. Wiley, London (2006)
10. Kupitz, Y.S., Martini, H., Spirova, M.: The Fermat-Torricelli problem, part I: a discrete gradient-method approach. *J. Optim. Theory Appl.* **158**(2), 305–327 (2013)
11. Weiszfeld, E.: On the point for which the sum of the distances to n given points is minimum. *Ann. Oper. Res.* **167**, 7–41 (2009). Translated from the French original [Tohoku Math. J. 43 (1937), 355–386] and annotated by Frank Plastria
12. Kuhn, H.W., Kuenne, R.E.: An efficient algorithm for the numerical solution of the generalized Weber problem in spatial economics. *J. Reg. Sci.* **4**, 21–34 (1962)
13. Beck, A., Teboulle, M.: Gradient-based algorithms with applications to signal recovery problems. In: Palomar, D., Eldar, Y. (eds.) *Convex Optimization in Signal Processing and Communications*, pp. 139–162. Cambridge University Press, New York (2009)
14. Kuhn, H.W.: A note on Fermat’s problem. *Math. Program.* **4**, 98–107 (1973)
15. Miehle, W.: Link-length minimization in networks. *Oper. Res.* **6**, 232–243 (1958)
16. Cooper, L.: Location-allocation problems. *Oper. Res.* **11**, 331–343 (1963)
17. Bertsekas, D.P.: *Nonlinear Programming*, 2nd edn. Athena Scientific, Belmont, MA (1999)
18. Nesterov, Y.: *Introductory Lectures on Convex Optimization*. Kluwer Academic Publishers, Boston, MA (2004)
19. Chandrasekaran, R., Tamir, A.: Open questions concerning Weiszfeld’s algorithm for the Fermat-Weber location problem. *Math. Program.* **44**, 293–295 (1989)
20. Brimberg, J.: The Fermat-Weber location problem revisited. *Math. Program.* **71**, 71–76 (1995)
21. Cánovas, L., Cañavate, R., Marín, A.: On the convergence of the Weiszfeld algorithm. *Math. Program.* **93**, 327–330 (2002)
22. Vardi, Y., Zhang, C.H.: A modified Weiszfeld algorithm for the Fermat-Weber location problem. *Math. Program.* **90**, 559–566 (2001)
23. Rautenbach, D., Struzyna, M., Szegedy, C., Vygen, J.: *Weiszfeld’s Algorithm Revisited Once Again*. Forschungsinstitut für Diskrete Mathematik. Rheinische Friedrich-Wilhelms-Universität, Bonn, Report (2004)
24. Chen, R.: Solution of location problems with radial cost functions. *Comput. Math. Appl.* **10**, 87–94 (1984)
25. Drezner, Z.: A note on the Weber location problem. *Ann. Oper. Res.* **40**, 153–161 (1992)
26. Drezner, Z.: A note on accelerating the Weiszfeld procedure. *Locat. Sci.* **3**, 275–279 (1995)
27. Calamai, P.H., Conn, A.R.: A projected newton method for l_p norm location problems. *Math. Program.* **38**, 75–109 (1987)
28. Overton, M.L.: A quadratically convergent method for minimizing a sum of euclidean norms. *Math. Program.* **27**, 34–63 (1983)
29. Ben-Tal, A., Nemirovski, A.: *Lectures on Modern Convex Optimization*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA (2001)
30. Nesterov, Y., Nemirovski, A.: *Interior-point Polynomial Algorithms in Convex Programming*. SIAM Studies in Applied Mathematics. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA (1994)
31. Andersen, K.D., Christiansen, E., Conn, A.R., Overton, M.L.: An efficient primal-dual interior-point method for minimizing a sum of Euclidean norms. *SIAM J. Sci. Comp.* **22**, 243–262 (2000)
32. Katz, I.N.: Local convergence in Fermat’s problem. *Math. Program.* **6**, 89–104 (1974)
33. Polyak, B.T.: *Introduction to Optimization*. Translations Series in Mathematics and Engineering. Optimization Software Inc., Publications Division, New York (1987). Translated from the Russian. With a foreword by Dimitri P. Bertsekas
34. Beck, A., Teboulle, M.: A fast iterative shrinkage-thresholding algorithm for linear inverse problems. *SIAM J. Imaging Sci.* **2**, 183–202 (2009)
35. Nesterov, Y.: A method for solving the convex programming problem with convergence rate $O(1/k^2)$. *Dokl. Akad. Nauk SSSR* **269**, 543–547 (1983)
36. Beck, A., Teboulle, M.: Smoothing and first order methods: a unified framework. *SIAM J. Optim.* **22**, 557–580 (2012)
37. Ben-Tal, A., Teboulle, M.: A smoothing technique for nondifferentiable optimization problems. In: *Optimization* (Varetz, 1988), Lecture Notes in Math., vol. 1405, pp. 1–11. Springer, Berlin (1989)

38. Bertsekas, D.P.: Nondifferentiable optimization via approximation. *Math. Program. Stud.* pp. 1–25 (1975)
39. Bertsekas, D.P.: *Constrained Optimization and Lagrangian Multipliers*. Academic Press, New York (1982)
40. Moreau, J.J.: Proximité et dualité dans un espace Hilbertien. *Bull. Soc. Math. Fr.* **93**, 273–299 (1965)
41. Nesterov, Y.: Smooth minimization of non-smooth functions. *Math. Program.* **103**, 127–152 (2005)
42. Huber, P.J.: *Robust Statistics*. Wiley, New York (1981)
43. Radó, F.: The Euclidean multifacility location problem. *Oper. Res.* **36**, 485–492 (1988)
44. Ben-Israel, A., Iyigun, C.: A generalized Weiszfeld method for the multi-facility location problem. *Oper. Res. Lett.* **38**, 207–214 (2010)
45. Teboulle, M.: A unified continuous optimization framework for center-based clustering methods. *J. Mach. Learn. Res.* **8**, 65–102 (2007)
46. Katz, I.N., Vogl, S.R.: A Weiszfeld algorithm for the solution of an asymmetric extension of the generalized Fermat location problem. *Comput. Math. Appl.* **59**, 399–410 (2010)
47. Cooper, L.: An extension to the generalized Weber problem. *J. Reg. Sci.* **8**, 181–197 (1968)
48. Morris, J.G.: Convergence of the Weiszfeld algorithm for Weber problems using a generalized “distance” function. *Oper. Res.* **29**, 37–48 (1981)
49. Erdős, P.: Open problems. *J. Math. Phys. High Sch. Stud.* **43**(10), 444 (1993)
50. Beck, A., Teboulle, M., Chikishev, Z.: Iterative minimization schemes for solving the single source localization problem. *SIAM J. Optim.* **19**, 1397–1416 (2008)
51. Biswas, P., Lian, T.C., Wang, T.C., Ye, Y.: Semidefinite programming based algorithms for sensor network localization. *ACM Trans. Sen. Netw.* **2**(2), 188–220 (2006)
52. Beaton, A.E., Tukey, J.W.: The fitting of power series, meaning polynomials, illustrated on bandspectroscopic data. *Technometrics* **16**, 147–185 (1974)
53. Ben-Tal, A., Teboulle, M., Yang, W.H.: A least-squares-based method for a class of nonsmooth minimization problems with applications in plasticity. *Appl. Math. Optim.* **24**, 273–288 (1991)
54. Daubechies, I., DeVore, R., Fornasier, M., Güntürk, C.S.: Iteratively reweighted least squares minimization for sparse recovery. *Commun. Pure Appl. Math.* **63**, 1–38 (2010)
55. Lawson, C.L.: *Contributions to the Theory of Linear Least Maximum Approximation*. ProQuest LLC, Ann Arbor, MI (1961). Thesis (Ph.D.)-University of California, Los Angeles
56. O’Leary, D.P.: Robust regression computation using iteratively reweighted least squares. *SIAM J. Matrix Anal. Appl.* **11**, 466–480 (1990)
57. Rice, J.R., Usow, K.H.: The Lawson algorithm and extensions. *Math. Comp.* **22**, 118–127 (1968)
58. Kuhn, H.W.: On a pair of dual nonlinear programs. *Nonlinear Programming (NATO Summer School, Menton, 1964)*, pp. 37–54. North-Holland, Amsterdam (1967)
59. Fasbender, E.: Über die gleichseitigen Dreiecke welche um ein gegebenes Dreieck gelegt werden können. *J. F. Math* **30**, 230–231 (1846)