

# Pitching Pennies into a Magnet

Kirk T. McDonald

*Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544*

(Jan. 15, 1997)

## 1 Problem

If one pitches a penny into a large magnet, eddy currents are induced in the penny, and their interaction with the magnetic field results in a repulsive force, according to Lenz' law. Estimate the minimum velocity needed for a penny to enter a long, 1-T solenoid magnet whose diameter is 0.1 m.

You may suppose that the penny moves so that its axis always coincides with that of the magnet, and that gravity may be ignored. The speed of the penny is low enough that the magnetic field caused by the eddy currents may be neglected compared to that of the solenoid. Equivalently, you may assume that the magnetic diffusion time is small.

## 2 Solution

The penny has radius  $a$  and thickness  $\Delta z$ . For the motion as stated in the problem, the eddy current will flow in concentric rings about the center of the disk. Therefore, we first examine a ring of radius  $r$  and radial extent  $\Delta r$ .

The magnetic flux through the ring at position  $z$  is

$$\Phi \approx \pi r^2 B_z(0, z), \quad (1)$$

whose time rate of change is

$$\dot{\Phi} = \pi r^2 \dot{B}_z = \pi r^2 B'_z v, \quad (2)$$

where  $\dot{\phantom{x}}$  indicates differentiation with respect to time,  $'$  is differentiation with respect to  $z$ ,  $B_z$  stands for  $B_z(0, z)$ , and  $v$  is the velocity of the center of mass of the ring.

The penny has electrical conductivity  $\sigma$ . Its resistance to currents around the ring is

$$R = \frac{2\pi r}{\sigma \Delta r \Delta z}, \quad (3)$$

so the (absolute value of the) induced current is

$$I = \frac{\mathcal{E}}{R} = \frac{\dot{\Phi}}{R} = \frac{\sigma r B'_z v \Delta r \Delta z}{2}, \quad (4)$$

(in MKSA units).

The azimuthal eddy current interacts with the radial component of the magnetic field to produce the axial retarding force. Close to the magnetic axis, we estimate the radial field in term of the axial field according to

$$B_r(r, z) \approx r \frac{\partial B_r(0, z)}{\partial r} = -\frac{r}{2} \frac{\partial B_z(0, z)}{\partial z} \equiv -\frac{r B'_z}{2}, \quad (5)$$

as can be deduced from the Maxwell equation  $\nabla \cdot \mathbf{B} = 0$ , noting that on the magnetic axis  $\partial B_r / \partial r = \partial B_x / \partial x = \partial B_y / \partial y$ . Then, the retarding force on the ring is

$$\Delta F_z = 2\pi r B_r I = -\pi \sigma r^2 B_r B'_z v \Delta r \Delta z \approx -\frac{\pi \sigma r^3 (B'_z)^2 v \Delta r \Delta z}{2}. \quad (6)$$

Alternatively, we note that the kinetic energy lost by the penny appears as Joule heating. Hence, for the ring analyzed above,

$$v \Delta F_z = \frac{dU}{dt} = -I^2 R = -\frac{\pi \sigma r^3 (B'_z)^2 v^2 \Delta r \Delta z}{2}, \quad (7)$$

using eqs. (3) and (4), which agains leads to eq. (6).

The equation of motion of the ring is

$$dF_z = -\frac{\pi \sigma r^3 (B'_z)^2 v \Delta r \Delta z}{2} = m \dot{v} = 2\pi \rho r \Delta r \Delta z v' v, \quad (8)$$

where  $\rho$  is the mass density of the metal. We integrate this equation with respect to radius to find

$$-\frac{\pi \sigma a^4 (B'_z)^2 v \Delta z}{8} = \pi \rho a^2 \Delta z v' v, \quad (9)$$

After dividing out the common factor  $\pi a^2 \Delta z v$ , we find

$$v' = -\frac{\sigma a^2 (B'_z)^2}{8\rho}. \quad (10)$$

For an estimate, we note that the peak gradient of the axial field of a solenoid of diameter  $D$  is about  $B_0/D$ , and the gradient is significant over a region  $\Delta z \approx D$ . Hence, on entering a solenoid the jet velocity is reduced by

$$\Delta v \approx \frac{\sigma a^2 B_0^2}{8\rho D}. \quad (11)$$

The penny must have initial velocity  $v_0 > \Delta v$  to enter the magnet.

A copper penny has  $a \approx 1 \text{ cm} = 10^{-2} \text{ m}$ , density  $\approx 10 \text{ g/cm}^3 = 10^4 \text{ kg/m}^3$ , electrical resistivity  $\approx 10^{-6} \text{ } \Omega\text{-cm}$ , and therefore conductivity  $\approx 10^8 \text{ MKSA units}$ . The minimum velocity to enter a 1-T magnet with diameter  $D = 0.1 \text{ m}$  is then,

$$v_{\min} \approx \frac{10^8 \cdot (10^{-2})^2 \cdot 1^2}{8 \cdot 10^4 \cdot 0.1} \approx 1 \text{ m/s}. \quad (12)$$

### 3 Acknowledgment

The author thanks Ahmed Hassanein, Bob Palmer and Bob Weggel for discussions of this problem. The case of a sphere rather than a disk has been presented in [1].

### 4 References

- [1] J. Walker and W.H. Wells, *Drag Force on a Conducting Spherical Drop in a Nonuniform Magnetic Field*, ORNL/TN-6976 (Sept. 1979).