

A Combinatorial Characterization of the Testable Graph Properties: It's All About Regularity

Extended Abstract + Appendix

Noga Alon* Eldar Fischer† Ilan Newman‡ Asaf Shapira§

Abstract

A common thread in all the recent results concerning testing dense graphs is the use of Szemerédi's regularity lemma. In this paper we show that in some sense this is not a coincidence. Our first result is that the property defined by having any given Szemerédi-partition is testable with a constant number of queries. Our second and main result is a purely combinatorial characterization of the graph properties that are testable with a constant number of queries. This characterization (roughly) says that a graph property \mathcal{P} can be tested with a constant number of queries **if and only if** testing \mathcal{P} can be reduced to testing the property of satisfying one of finitely many Szemerédi-partitions. This means that in some sense, testing for Szemerédi-partitions is as hard as testing any testable graph property. We thus resolve one of the main open problems in the area of property-testing, which was first raised in the 1996 paper of Goldreich, Goldwasser and Ron [24] that initiated the study of graph property-testing. This characterization also gives an intuitive explanation as to what makes a graph property testable.

*Schools of Mathematics and Computer Science, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv, Israel, and IAS, Princeton, NJ. Email: nogaa@tau.ac.il. Research supported in part by a USA-Israeli BSF grant, by the Israel Science Foundation and by the Hermann Minkowski Minerva Center for Geometry at Tel Aviv University.

†Faculty of Computer Science, Technion – Israel institute of technology, Technion City, Haifa 32000, Israel. Research supported in part by an Israel Science Foundation grant number 55/03.

‡Department of Computer Science, Haifa University, Haifa 31905, Israel. Research supported in part by an Israel Science Foundation grant number 55/03.

§School of Computer Science, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv, Israel. Email: asafico@tau.ac.il. This work forms part of the author's Ph.D. thesis. Research supported in part by a Charles Clore Foundation Fellowship and by an IBM Ph.D. Fellowship.

1 Background

1.1 Basic definitions

The meta problem in the area of property testing is the following: Given a combinatorial structure S , distinguish between the case that S satisfies some property \mathcal{P} and the case that S is ϵ -far from satisfying \mathcal{P} . Roughly speaking, a combinatorial structure is said to be ϵ -far from satisfying some property \mathcal{P} if an ϵ -fraction of its representation should be modified in order to make S satisfy \mathcal{P} . The main goal is to design randomized algorithms, which look at a very small portion of the input, and using this information distinguish with high probability between the above two cases. Such algorithms are called *property testers* or simply *testers* for the property \mathcal{P} . Preferably, a tester should look at a portion of the input whose size is a function of ϵ only. Blum, Luby and Rubinfeld [10] were the first to formulate a question of this type, and the general notion of property testing was first formulated by Rubinfeld and Sudan [34], who were interested in studying various algebraic properties such as linearity of functions.

The main focus of this paper is the testing of properties of graphs. More specifically, we focus on property testing in the dense graph model as defined in [24]. In this case a graph G is said to be ϵ -far from satisfying a property \mathcal{P} , if one needs to add/delete at least ϵn^2 edges to G in order to turn it into a graph satisfying \mathcal{P} . A tester for \mathcal{P} should distinguish with high probability, say $2/3$, between the case that G satisfies \mathcal{P} and the case that G is ϵ -far from satisfying \mathcal{P} . Here we assume that the tester can query some oracle whether a pair of vertices, i and j , are adjacent in the input graph G . In what follows we will say that a tester for a graph property \mathcal{P} has *one-sided error* if it accepts any graph satisfying \mathcal{P} with probability 1 (and still rejects those that are ϵ -far with probability at least $2/3$). If the tester may reject graphs satisfying \mathcal{P} with non-zero probability then it is said to have *two-sided error*. The following notion of efficient testing will be the main focus of this paper:

Definition 1.1 (Testable) *A graph property \mathcal{P} is testable if there is a randomized algorithm T , that can distinguish with probability $2/3$ between graphs satisfying \mathcal{P} and graphs that are ϵ -far from satisfying \mathcal{P} , while making a number of edge queries which is bounded by some function $q(\epsilon)$ that is independent of the size of the input.*

The study of the notion of testability for combinatorial structures, and mainly the dense graph model, was introduced in the seminal paper of Goldreich, Goldwasser and Ron [24]. Graph property testing has also been studied in the *bounded-degree* model [26], and the newer *general density* model [30]. We note that in these models a property is usually said to be testable if the number of queries is $o(n)$. Following [24, 10, 34] property testing was studied in various other contexts such as boolean functions [4, 16, 19, 31], geometric objects [2, 11] and algebraic structures [10, 20, 8]. See the surveys [14, 33] for additional results and references.

1.2 Background on the characterization project

With this abundance of results on property testing, a natural question is what makes a combinatorial property testable. In particular, characterizing the testable graph properties was considered one of the main open problems in the area of property testing, and was raised already in the 1996 paper of Goldreich, Goldwasser and Ron [24], see also [22], [9] and [25]. In this paper we obtain for the first

time a characterization of the testable graph properties. We next discuss some results related to this problem.

A natural strategy toward obtaining a characterization of the testable graphs was to either prove the testability/non-testability of general families of graph properties or to obtain characterizations for special cases of testers. The main results of [24] was that a general family of so called “partition-problems” are all testable. These include the properties of being k -colorable, having a large cut and having a large clique. [25] gave a characterization of the partition-problems that can be tested with 1-sided error. They also proved that not all graph properties that are closed under edge-removal are testable. [12] studied property testing via the framework of *abstract combinatorial programs* and gave certain characterizations within this framework. [3] tried to obtain a *logical* characterization of the testable graph properties. More specifically, it was shown that every first order graph-property of type $\exists\forall$ (see [3]) is testable, while there are first-order graph properties of type $\forall\exists$ that are not testable. The main technical result of [3] was that certain abstract colorability properties are all testable. These results were generalized in [13]. In [6] it was shown that every graph property that is closed under removal of edges and vertices is testable. This result was extended in [7], where it was shown that in fact, being closed under vertex removal is already sufficient for being testable (see also [28]). [7] also contains a characterization of the graph properties that can be tested with *one-sided* error by certain restricted testers. Finally, [25] following [3], proved that a tester may be assumed to be non-adaptive (see Theorem 4.2), and [17] proved that if a graph property is testable then it is also possible to estimate how far is a given graph from satisfying the property (see Theorem 4). These last two results are key ingredients in the present paper.

2 The Main Result

2.1 Background on Szemerédi’s regularity lemma

Our main result in this paper gives a purely combinatorial characterization of the testable graph properties. As we have previously mentioned, the first properties that were shown to be testable in [24] were certain graph partition properties. As it turns out, our characterization relies on certain “enhanced” partition properties, whose existence is guaranteed by the celebrated regularity lemma of Szemerédi [35]. We start by introducing some standard definitions related to the regularity lemma. For a comprehensive survey about the regularity lemma the reader is referred to [27].

For every two nonempty disjoint vertex sets A and B of a graph G , we define $e(A, B)$ to be the number of edges of G between A and B . The *edge density* of the pair is defined by $d(A, B) = e(A, B)/|A||B|$.

Definition 2.1 (γ -regular pair) *A pair (A, B) is γ -regular, if for any two subsets $A' \subseteq A$ and $B' \subseteq B$, satisfying $|A'| \geq \gamma|A|$ and $|B'| \geq \gamma|B|$, the inequality $|d(A', B') - d(A, B)| \leq \gamma$ holds.*

Throughout the paper it will be useful to observe that in the above definition it is enough to require that $|d(A', B') - d(A, B)| \leq \gamma$ for sets $A' \subseteq A$ and $B' \subseteq B$, of sizes $|A'| = \gamma|A|$ and $|B'| = \gamma|B|$. A partition $\mathcal{A} = \{V_i \mid 1 \leq i \leq k\}$ of the vertex set of a graph is called an *equipartition* if $|V_i|$ and $|V_j|$ differ by no more than 1 for all $1 \leq i < j \leq k$ (so in particular every V_i has one of two possible sizes). The *order* of an equipartition denotes the number of partition classes (k above).

Definition 2.2 (γ -regular equipartition) An equipartition $\mathcal{B} = \{V_i \mid 1 \leq i \leq k\}$ of the vertex set of a graph is called γ -regular if all but at most $\gamma \binom{k}{2}$ of the pairs (V_i, V_j) are γ -regular.

In what follows an equipartition is said to refine another if every set of the former is contained in one of the sets of the latter. Szemerédi's regularity lemma can be formulated as follows.

Lemma 2.3 ([35]) For every m and $\gamma > 0$ there exists $T = T_{2,3}(m, \gamma)$ with the following property: If G is a graph with $n \geq T$ vertices, and \mathcal{A} is any equipartition of the vertex set of G of order at most m , then there exists a refinement \mathcal{B} of \mathcal{A} of order k , where $m \leq k \leq T$ and \mathcal{B} is γ -regular. In particular, for every m and $\gamma > 0$ there exists $T = T_{2,3}(m, \gamma)$ such that any graph with $n \geq T$ vertices, has a γ -regular equipartition of order k , where $m \leq k \leq T$.

The regularity lemma guarantees that every graph has a γ -regular equipartition of (relatively) small order. As it turns out in many applications of the regularity lemma, one is usually interested in the densities of the bipartite graphs connecting the sets V_i of the regular partitions. In fact, one important consequence of the regularity lemma is that in many cases knowing the densities connecting the sets V_i (approximately) tells us all we need to know about a graph. Roughly speaking, if a graph G has a regular partition of order k and we define a weighted graph $R(G)$, of size k , where the weight of edge (i, j) is $d(V_i, V_j)$, then by considering an appropriate property of $R(G)$ one can infer many properties of G . As the order of the equipartition is guaranteed to be bounded by a function of γ , this means that for many applications, any graph has an approximate description of *constant-complexity* (we will return to this aspect in a moment). As it turns out, this interpretation of the regularity lemma is the key to our characterization. We believe that our characterization of the testable graph properties is an interesting application of this aspect of the regularity lemma.

Given the above discussion it seems natural to define a graph property, which states that a graph has a given γ -regular partition, that is, an equipartition which is γ -regular and such that the densities between the sets V_i satisfy some predefined set of densities.

Definition 2.4 (Regularity-Instance) A regularity-instance is given by an error-parameter $0 < \gamma \leq 1$, an integer k , a set of $\binom{k}{2}$ densities $0 \leq \eta_{ij} \leq 1$ indexed by $1 \leq i < j \leq k$, and a set \bar{R} of pairs (i, j) of size at most $\gamma \binom{k}{2}$. A graph is said to satisfy the regularity-instance if it has an equipartition $\{V_i \mid 1 \leq i \leq k\}$ such that for all $(i, j) \notin \bar{R}$ the pair (V_i, V_j) is γ -regular and satisfies $d(V_i, V_j) = \eta_{i,j}$. The complexity of the regularity-instances is $\max(k, 1/\gamma)$.

Note, that in the above definition the set \bar{R} corresponds to the set of pairs (i, j) for which (V_i, V_j) is not necessarily a γ -regular pair (possibly, there are at most $\gamma \binom{k}{2}$ such pairs). Also, note that the definition of a regularity-instance does not impose any restriction on the graphs spanned by any single set V_i . By Theorem 2.3, for any $0 < \gamma \leq 1$ any graph satisfies some regularity instance with an error parameter γ and with an order bounded by a function γ . The first step needed in order to obtain our characterization of the testable properties, is that the property of satisfying any given regularity-instance is testable. This is also the main technical result of this paper.

Theorem 1 For any regularity-instance R , the property of satisfying R is testable.

2.2 The characterization

Many of the recent results on testing graph properties in the dense graph model relied on Lemma 2.3. Our main result in this paper shows that this is not a coincidence. Each of the papers which applied the regularity lemma to test a graph property used different aspects of what can be inferred from certain properties of a regular partition of a graph. These results however, use the properties of the regularity partition in an *implicit* way. For example, the main observation needed in order to infer the triangle-freeness is testable, is that if the regularity partition has three sets V_i, V_j, V_k , which are connected by regular and dense bipartite graphs then the graph is far from being triangle-free. However, to *test* triangle freeness we do not need to know the regular partition, we just need to find a triangle in the graph. As Theorem 1 allows us to test for having a certain regular partition, it seems possible to try and test properties by *explicitly* checking for properties of the regularity partition of the input. Returning to the previous discussion on viewing the regularity lemma as constant complexity description of a graph, being able to explicitly test for having a given regular partition, should allow us to test more complex properties as we can obtain all the information of the regular partition and not just *consequences* of having some regular partition. The next definition tries to capture the graph properties \mathcal{P} , such that \mathcal{P} can be tested via testing a certain set of regularity instances.

Definition 2.5 (Regular-Reducible) *A graph property \mathcal{P} is regular-reducible if for any $\delta > 0$ there exists an $r = r(\delta)$ such that for any n there is a family \mathcal{R} of at most r regularity-instances each of complexity at most r , such that the following holds for every n -vertex graph G :*

1. *If G satisfies \mathcal{P} then for some $R \in \mathcal{R}$, G is δ -close to satisfying R .*
2. *If G is ϵ -far from satisfying \mathcal{P} , then for any $R \in \mathcal{R}$, G is $(\epsilon - \delta)$ -far from satisfying R .*

The reader may want to observe that in the above definition the value of δ may be arbitrarily close to 0. If we think of $\delta = 0$ then we get that a graph satisfies \mathcal{P} if and only if it satisfies one of the regularity instances of \mathcal{R} . With this interpretation in mind, in order to test \mathcal{P} one can test the property of satisfying any one of the instances of \mathcal{R} . Therefore, in some sense we “reduce” the testing of property \mathcal{P} to the testing of regularity-instances. As the main result of this paper states, the testable graph properties are precisely those for which testing them can be carried out by testing for some property of their regular partitions.

Theorem 2 (Main Result) *A graph property is testable if and only if it is regular-reducible.*

If we have to summarize the moral of our characterization in one simple sentence, then it says that a graph property \mathcal{P} is testable if and only if \mathcal{P} is such that knowing a regular partition of a graph G is sufficient for telling whether G is ϵ -far or ϵ -close to satisfying \mathcal{P} . In other words, there is a short “proof” that G is either ϵ -close or ϵ -far from satisfying \mathcal{P} . Thus, in a more “computational complexity” jargon, we could say that a graph property is testable if and only if it has the following “interactive proof”: A prover gives a verifier the description of a regularity-instance R , which the input G is (supposedly) close to satisfying. The verifier, using Theorem 1, then verifies if G is indeed close to satisfying R . The way to turn this interactive proof into a testing algorithm is to apply the constant-complexity properties of the regularity lemma that we have previously discussed; as the order of the regular partition is bounded by a function of ϵ , then there are only *finitely* many

regularity-instances that the prover may potentially send to the verifier. Therefore, the verifier does not need to get an alleged regular-instances, it can simply try them all! Theorem 2 thus states that in some sense testing regularity-instances is the “hardest” property to test, because by Theorem 2 any testing algorithm can be turned into a testing algorithm for regularity-instances. However, we stress that this is true only on the *qualitative* level, because using Theorem 2 in order to turn a tester into an equivalent tester, which tests for regularity-instances, may significantly increase its query complexity. The main reason is that the proofs of Theorems 1 and 2 apply Lemma 2.3 and thus only give weak upper bounds. Having said that, it should also be clear that one cannot prove general results on testing graph properties which guarantee good upper bounds (say, $\text{poly}(1/\epsilon)$) on the query complexity as it was proved in [6] that there are graph properties (even monotone ones) that are testable and yet may require arbitrarily large query complexity. We also note that the terminology of regular-reducible is not far from being a standard reduction because in order to prove one of the directions of Theorem 2 we indeed test a property \mathcal{P} , which is regular-reducible to a set \mathcal{R} , by testing the regularity-instances of \mathcal{R} . Theorem 2 also gives further convincing evidence as to the “combinatorial” nature of property testing in the dense graph model as was recently advocated by Goldreich [23].

As is evident from Definition 2.5, the characterization given in Theorem 2 is not a “quick recipe” for inferring whether a given property is testable. Still, we can use Theorem 2 in order to obtain unified proofs for several previous results. As we have alluded to before, these results can be inferred by showing that it is possible (or impossible) to test if a graph satisfies certain regularity-instances. We believe that these proofs give some (non-explicit) structural explanation as to what makes a graph property testable. See Section 7 for more details. It is thus natural to ask if one can come up with more “handy” characterizations. We doubt that such a characterization exists, mainly because it should (obviously) be equivalent to Theorem 2. Of course, we cannot formally prove that no simpler sufficient condition exists. However, as we discuss in the next subsection we can at least disprove a possible simpler sufficient condition of testability.

2.3 Disproving a simpler sufficient condition

Observing previous results in property testing (not necessarily of graphs) reveals that essentially all the properties that were shown to be testable had the following *downscaling* property: If an object is close to satisfying a property then a sample from the object will not be very far from satisfying it. This is true, for example, for properties studied in the context of functions [16, 19], graphs [24, 7, 13], geometric objects [2, 11], algebraic structures [10, 20, 8] and languages [4, 31]. Therefore, a natural question is whether being downscaling is enough for being testable. To formally state this feature for graphs we introduce the following definition:

Definition 2.6 (Downscaling) *A graph property \mathcal{P} is downscaling if for every $\delta > 0$ there is a $q = q(\delta)$ with the following property: suppose that an n -vertex graph is ϵ -close to satisfying \mathcal{P} . Then, for every m such that $q \leq m \leq n$ the graph induced by a randomly chosen set of vertices of size m is $(\epsilon + \delta)$ -close to satisfying \mathcal{P} with probability at least $2/3$.*

We note that essentially the same definition as above applies to other combinatorial structures. The reader may want to note, that the family of hereditary properties, which were shown to be testable in [7] (see also [28]), are all downscaling. Also, all the partition properties, which were shown to be testable in [24], are downscaling. If downscaling was indeed sufficient for being testable,

this would immediately give simple and uniform proofs for many testability results. As it turns out however, this is not the case.

Theorem 3 *There is a downscaling non-testable graph property.*

2.4 Organization and overview of the paper

The first main technical step of the proof of Theorem 2 is taken in Section 3. In this section we prove that if the densities of pairs of subsets of vertices of a bipartite graph are close to the density of the bipartite graph itself, then the bipartite graph can be turned into a regular-pair using relatively few edge modifications. Rephrasing this gives that we can increase the regularity measure of a bipartite pair by making relatively few edge modifications. The second main step is taken in Section 5. In this section we show that sampling a constant number of vertices, guarantees that the sample and the graph will have (roughly) the *same* set of regular partitions. We believe that this result may be of independent interest. By applying the results of Sections 3 and 5 we prove Theorem 1 in Section 6. In this section we also prove one of the directions of Theorem 2, asserting that if a graph property is regular-reducible then it is testable. Along with Theorem 1, a second tool that we need in order to prove this direction is the main result of [17]. We apply this result in order to infer that for any regularity-instance R , one cannot only test for the property of satisfying R , but can also estimate how far is a given graph from satisfying R . This *estimation* of the distance to satisfying regularity-instances is key to *testing* a property via a regularity-reduction. The proof of the second direction of Theorem 2 appears in Section 4. To prove this direction we first show that knowing that a graph G satisfies a regularity instance enables us to estimate the number of copies of certain graphs in G . We then apply the main result of [25] about canonical testers along with the main result of Section 3 in order to “pick” those regularity-instances that can constitute the family \mathcal{R} in Definition 2.5. In Section 7 we use Theorem 2 in order to reprove some previously known results in property-testing. The main interest of these proofs is that they apply Theorem 2 in order to prove in a unified manner results that had distinct proofs. Due to space limitations the proof of Theorem 3 will appear in the full version of the paper. We briefly mention that in order to prove this theorem we use a subtle variation of the graph isomorphism problem, which was known to be non-testable but is far from being downscaling. Section 8 contains some concluding remarks. Due to space limitations most proofs appear in the appendix of this extended abstract.

3 Enhancing Regularity with Few Edge Modifications

The definition of γ -regular pair of density η requires a pair of sets of vertices to satisfy several density requirements. Our main goal in this section is to show that if a pair of vertex sets are close (in an appropriate sense) to satisfying these requirements, then it is indeed close to a γ -regular pair of density η . For example, consider the property of being a 0.1-regular pair with edge density 0.5. Intuitively, it seems that if the edge density of a bipartite graph G on vertex sets A and B of size m each, is close to 0.5 and the density of any pair $A' \subseteq A$ and $B' \subseteq B$ of sizes $0.1m$ is close to 0.5 ± 0.1 , then G should be close to satisfying the property. However, note that in this case it may be the case that there are pairs (A', B') whose density is smaller than 0.4 and other pairs, whose density is larger than 0.6. Thus, only removing or only adding edges (even randomly) will most likely not turn G into a 0.1-regular pair of density 0.5. In order to show that G is indeed close to satisfying the property,

we take a convex combination of G with a random graph, whose density is $1/2$. The intuition is that the random graph will not change the density of G much, but, because a random graph is highly regular, this will increase the regularity of G . The main result of this section is formalized in the following lemma, which is an important ingredient in the proofs of both directions of Theorem 2.

In this lemma, as well as throughout the rest of the paper, when we write $x = a \pm b$ we mean $a - b \leq x \leq a + b$.

Lemma 3.1 *The following holds for any $0 < \delta \leq \gamma \leq 1$: Suppose that (A, B) is a $(\gamma + \delta)$ -regular pair with density $\eta \pm \delta$, where $|A| = |B| = m \geq m_{3.1}(\eta, \delta)$. Then, it is possible to make at most $50 \frac{\delta}{\gamma^2} m^2$ edge modifications and turn (A, B) into a γ -regular pair with density precisely η .*

The proof of Lemma 3.1 has two main steps, which are captured in Lemmas 3.2 and 3.3 below. The proofs appear in Appendix 9. The first step is given in the following lemma that enables us to make relatively few edge modifications and thus make sure that the density of a pair is exactly what it should be, while at the same time not decreasing its regularity by much.

Lemma 3.2 *Suppose that (A, B) is a $(\gamma + \delta)$ -regular pair satisfying $d(A, B) = \eta \pm \delta$, where $|A| = |B| = m \geq m_{3.2}(\eta, \delta)$. Then, it is possible to make at most $2\delta m^2$ modifications, and thus turn (A, B) into a $(\gamma + 2\delta)$ -regular pair with density precisely η .*

The second and main step, which implements the main idea presented at the beginning of this section, takes a bipartite graph, whose density is precisely η , and returns a bipartite graph, whose density is still η but with a better regularity measure.

Lemma 3.3 *The following holds for any $0 < \delta \leq \gamma \leq 1$. Let A and B be two vertex sets of size $m \geq m_{3.3}(\delta, \gamma)$, satisfying $d(A, B) = \eta$. Suppose further that for any pair of subsets $A' \subseteq A$ and $B' \subseteq B$ of size γm we have $d(A', B') = \eta \pm (\gamma + \delta)$. Then, it is possible to make at most $\frac{3\delta}{\gamma} m^2$ edge modifications and thus turn (A, B) into a γ -regular pair with density precisely η .*

4 Any Testable Property is Regular-Reducible

In this section we prove the first direction of Theorem 2, namely,

Lemma 4.1 *If a graph property is testable then it is regular-reducible.*

Our starting point in the proof of Lemma 4.1 is the following result of [25] (extending a result of [3]) about canonical testers:

Lemma 4.2 ([25],[3]) *If graph property \mathcal{P} can be tested on n -vertex graphs with $q = q(\epsilon, n)$ edge queries, then it can also be tested by a tester, which makes its queries by uniformly and randomly choosing a set of $2q$ vertices, querying all their pairs and then accepting or rejecting (deterministically) according to the graph induced by the sample, the value of ϵ and the value of n .*

Restating the above, by (at most) squaring the query complexity, we can assume without loss of generality that a property-tester works by sampling a set of vertices of size $q(\epsilon, n)$ and accepting or rejecting according to some graph property of the sample. As noted in [25] the graph property that the algorithm may search for in the sample, may be different from the property, which is tested. In fact, the property the algorithm checks for in the sample may depend on ϵ and on the size of

the input graph. Our main usage of Theorem 4.2 is that it allows to pick the graphs of size q that cause a tester for \mathcal{P} to accept. The first technical step we take toward proving Lemma 4.1 is proving some technical results about induced copies of graphs spanned by graphs satisfying a given regularity-instance. These results enable us to deduce from the fact that a graph satisfies some regularity-instance the probability that a given tester accepts the graph. We then use these results along with Theorem 4.2 and some additional arguments in order to prove that any testable property is regular reducible. The full details appear in Appendix 10.

5 Sampling Regular Partitions

The main result of this section (roughly) asserts that for every fixed γ , if we sample a constant number of vertices from a graph G then with high probability, the graph induced by the sample and the graph G will have the same set of γ -regular partitions. The proofs of this section appear in Appendix 11. To formally state this result we introduce the following definition:

Definition 5.1 (δ -similar regular-partition) *An equipartition $\mathcal{U} = \{U_i \mid 1 \leq i \leq k\}$ is δ -similar to a γ -regular equipartition $\mathcal{V} = \{V_i \mid 1 \leq i \leq k\}$, of the same order k (where $0 < \gamma \leq 1$), if: (1) $d(U_i, U_j) = d(V_i, V_j) \pm \delta$ for all $i < j$. (2) Whenever (V_i, V_j) is γ -regular, (U_i, U_j) is $(\gamma + \delta)$ -regular.*

Observe, that in the above definition the two equipartitions \mathcal{V} and \mathcal{U} may be equipartitions of different graphs. In what follows, if $G = (V, E)$ is a graph and $Q \subseteq V(G)$, then $G[Q]$ denotes the graph induced by G on Q . Our main result in this section is the following:

Lemma 5.2 *For every k, δ there is $q = q_{5.2}(k, \delta)$ such that a sample Q , of q vertices from a graph G , satisfies the following with probability at least $2/3$: If G (resp. $G[Q]$) has a γ -regular equipartition \mathcal{V} of order at most k , then $G[Q]$ (resp. G) has an equipartition \mathcal{U} , which is δ -similar to \mathcal{V} .*

The proof of Lemma 5.2 has two main stages. For the first we need a weaker result, which says that a sample of vertices has a regular partition, but with a *weaker* regularity measure.

Lemma 5.3 ([13]) *For every k and γ there is $q = q_{5.3}(k, \gamma)$ such that if a graph G has a γ -regular equipartition $\mathcal{V} = \{V_1, \dots, V_k\}$ of order k , then with probability at least $2/3$, a sample of q vertices will have an equipartition $\mathcal{U} = \{U_1, \dots, U_k\}$ satisfying: (1) $d(U_i, U_j) = d(V_i, V_j) \pm \delta$ for all $i < j$. (2) Whenever (V_i, V_j) is γ -regular (U_i, U_j) is $50\gamma^{1/5}$ -regular.*

For our purposes however, we can not allow a weaker regularity as in the above lemma. Our main tool in the proof of Lemma 5.2 is Lemma 5.5 below, which establishes that if two graphs share *one* γ -regular equipartition, then they share *all* the γ' -regular-partitions where γ' is slightly larger than γ . This will allow us to strengthen Lemma 5.3 and thus obtain Lemma 5.2. For the statement of this lemma we need the following definition:

Definition 5.4 ((δ, γ) -similar regular-partitions) *Two equipartitions $\mathcal{V} = \{V_i \mid 1 \leq i \leq k\}$ and $\mathcal{U} = \{U_i \mid 1 \leq i \leq k\}$ of the same order k , are said to be (δ, γ) -similar if: (1) $d(U_i, U_j) = d(V_i, V_j) \pm \delta$ for all $i < j$. (2) For all but at most $\gamma \binom{k}{2}$ of the pairs $i < j$, both (V_i, V_j) and (U_i, U_j) are γ -regular.*

Lemma 5.5 *For every k and δ there is $\zeta = \zeta_{5.5}(k, \delta)$ with the following property: suppose two graphs $G = (V, E)$ and $\bar{G} = (\bar{V}, \bar{E})$ have (ζ, ζ) -similar regular-equipartitions $\mathcal{V} = \{V_1, \dots, V_\ell\}$ and $\bar{\mathcal{V}} = \{\bar{V}_1, \dots, \bar{V}_\ell\}$ with $\ell \geq 1/\zeta$. Then, if \bar{G} has a γ -regular equipartition $\bar{\mathcal{A}} = \{\bar{A}_1, \dots, \bar{A}_k\}$ then G has an equipartition $\mathcal{A} = \{A_1, \dots, A_k\}$, which is δ -similar to $\bar{\mathcal{A}}$.*

6 Testing Regular Partitions and Proof of the Main Result

In this section we apply the results of Sections 3 and 5 in order to prove Theorem 2. In Appendix 12 we prove Theorem 1, thus establishing that any regularity instance is testable. Having established the testability of any given regularity-instance we can prove Theorem 2. The last tool we need for the proof is the main result of [17] about estimating graph properties.

Theorem 4 ([17]) *Suppose graph property \mathcal{P} is testable. Then for every $0 \leq \epsilon_1 < \epsilon_2 \leq 1$ there is a randomized algorithm for distinguishing between graphs that are ϵ_1 -close to satisfying \mathcal{P} and those that are ϵ_2 -far from satisfying it. Furthermore, the query complexity of the algorithm can be upper bounded by a function of $\epsilon_2 - \epsilon_1$.*

Proof (of Theorem 2): The first direction is given in Lemma 4.1. For the other direction, suppose that a graph property \mathcal{P} satisfies Definition 2.5. Let us fix n and ϵ . Put $r = r(\frac{1}{4}\epsilon)$ and let \mathcal{R} be the corresponding set of regularity instances for $\delta = \frac{1}{4}\epsilon$ as in Definition 2.5. Recall that Definition 2.5 guarantees that the number and the complexity of the regularity-instances of \mathcal{R} are bounded by a function of $\delta = \frac{1}{4}\epsilon$. By Theorem 1 for any regularity-instance $R \in \mathcal{R}$, the property of satisfying R is testable. Thus, by Theorem 4 for any such R , we can distinguish between graphs that are $\frac{1}{4}\epsilon$ -close to satisfying R from those that are $\frac{3}{4}\epsilon$ -far from satisfying it, while making a number of queries, which is upper bounded by a function of ϵ . In particular, by repeating the algorithm of Theorem 4 an appropriate number of times (that depends only on r), and taking majority, we get an algorithm for distinguishing between the above two cases, whose query complexity is a function of ϵ and r , which succeeds with probability at least $1 - \frac{1}{3^r}$. As r itself is bounded by a function of ϵ , the number of queries of this algorithm can be bounded by a function of ϵ only.

We are now ready to describe our tester for \mathcal{P} : Given a graph G of size n and $\epsilon > 0$ the algorithm uses, for every $R \in \mathcal{R}$, the version of Theorem 4 described in the previous paragraph, which succeeds with probability at least $1 - \frac{1}{3^r}$ in deciding, whether G is $\frac{1}{4}\epsilon$ -close to satisfying R or $\frac{3}{4}\epsilon$ -far from satisfying it. If it finds that G is $\frac{1}{4}\epsilon$ -close to satisfying some R , the algorithm accepts, otherwise it rejects. Observe, that as there are at most r regularity-instances in \mathcal{R} , we get by the union-bound, that with probability at least $2/3$ the subroutine for estimating how far is G from satisfying some $R \in \mathcal{R}$ never errs. Let's prove that the above algorithm is indeed a tester for \mathcal{P} . Suppose first that G satisfies \mathcal{P} . As we set $\delta = \frac{1}{4}\epsilon$ and \mathcal{P} is regular-reducible to \mathcal{R} , the graph G must be $\frac{1}{4}\epsilon$ -close to satisfying some regularity-instance $R' \in \mathcal{R}$. Suppose now that G is ϵ -far from satisfying \mathcal{P} . Again, as we assume that \mathcal{P} is regular-reducible to \mathcal{R} , we conclude that G must be $\frac{3}{4}\epsilon$ -far from satisfying all of the regularity-instances $R \in \mathcal{R}$. As with probability at least $2/3$ the algorithm correctly decides for any $R \in \mathcal{R}$ if G is $\frac{1}{4}\epsilon$ -close to satisfying R or $\frac{3}{4}\epsilon$ -far from satisfying it, we get that if G satisfies \mathcal{P} then with probability at least $2/3$ it will find that G is $\frac{1}{4}\epsilon$ -close to satisfying some $R \in \mathcal{R}$, while if G is ϵ -far from satisfying \mathcal{P} it will find that G is $\frac{3}{4}\epsilon$ -far from all $R \in \mathcal{R}$. By the definition of the algorithm, we get that with probability at least $2/3$ it distinguishes between graphs satisfying \mathcal{P} from those that are ϵ -far from satisfying it. This means that the algorithm is indeed a tester for \mathcal{P} . ■

7 Applications of the Main Result

In this section we show that Theorem 2 can be used in order to derive some positive and negative results on testing graph properties. Due to space limitations the proofs of the results appear in

Appendix 13. We would like to stress that all these proofs implicitly apply the main intuition behind our characterization, which was explained after the statement of Theorem 2, that a graph property is testable iff knowing the regularity partition of the graph is sufficient for inferring if a graph is far from satisfying the property. Our first application of Theorem 2 concerns testing for H -freeness; A graph is said to be H -free if it contains no (not necessarily induced) copy of H . It was implicitly proved in [1] that for any H , the property of being H -free is testable. The main idea of the proof in [1] is that if G is ϵ -far from being H -free then a large enough sample of vertices will contain a copy of H with high probability. Here we derive this result from Theorem 2 by giving an alternative proof, which checks if the input satisfies some regularity-instance. For simplicity, we only consider testing triangle-freeness. We briefly mention that an argument similar to the one we use in order to test triangle-freeness can be used to test any monotone graph property. However, to carry out the proof one needs one additional non-trivial argument, which was proved in [6], thus we refrain from including the proof.

Our second applications of Theorem 2 is concerned with testing k -colorability. This property was first implicitly proved to be testable in [32]. Much better upper bounds were obtained in [24], and further improved by [5]. As in the case of H -freeness the main ideas of the proofs in [32, 24, 5] is that if G is ϵ -far from being k -colorable then a large enough sample of vertices will not be k -colorable with high probability. Here we derive this result by applying Theorem 2. Though we derive here only the testability of k -colorability, simple variants of the argument can be used to show that all the partition-problems studied in [24] are testable¹.

The above mentioned apply Theorem 2 in order to obtain positive results. Our third application of 2 derives negative results. The main focus of [15] is testing for isomorphism to a given fixed graph. It shows that the query complexity of testing for isomorphism grows with a certain parameter, which measures the “complexity” of the graph. Without going into too much detail we just mention that under this measure random graphs are complex. Here we prove that testing for being isomorphic to a graph generated by $G(n, 0.5)$ requires a super-constant number of queries.

8 Concluding Remarks and Open Problems

The main result of this paper gives a combinatorial characterization of the graph properties, which can be tested with a constant number of edge queries in the dense graph model, possibly with two sided error. Together with the (near) characterization of [7] of the graph properties that can be tested with *one-sided* error, and the result of [17] showing that any testable property is also estimable, we thus get a more or less complete answer to all the *qualitative* questions on testing graph properties in the dense model. While property testing in the dense model is relatively well understood, there are no general positive/negative results on testing graph properties in the bounded-degree model [26] or the general density model [30]. In these models the query complexity of the tester usually depends on the size of the input. It seems interesting and challenging to obtain general results in these models. One interesting problem is which of the partition problems, which were studied in [24] can be tested using a sublinear number of queries. It will also be very interesting to give general positive and negative results concerning testing of boolean functions.

¹An alert reader may note that our proof of Theorem 2 applies the result of [17], which relies on the result of [24]. Thus, in the strict sense it is wrong to say that we infer the result of [24] from ours. However, it is not difficult to see that the result of [24] also follows from our (self-contained) proof of Lemma 5.2.

Acknowledgements: The authors would like to thank the organizers of the Dagstuhl Workshop on Sublinear Algorithms, where this research has emerged. A.S. would also like to thank his hosts at the Theory Group of Microsoft Research, where parts of this research were conducted.

References

- [1] N. Alon, R. A. Duke, H. Lefmann, V. Rödl and R. Yuster, The algorithmic aspects of the Regularity Lemma, Proc. of 33rd IEEE FOCS, Pittsburgh, IEEE (1992), 473-481. Also: J. of Algorithms 16 (1994), 80-109.
- [2] N. Alon, S. Dar, M. Parnas and D. Ron, Testing of clustering, Proc. 41st IEEE FOCS, IEEE (2000), 240-250.
- [3] N. Alon, E. Fischer, M. Krivelevich and M. Szegedy, Efficient testing of large graphs, Proc. of 40th FOCS, New York, NY, IEEE (1999), 656-666. Also: Combinatorica 20 (2000), 451-476.
- [4] N. Alon, M. Krivelevich, I. Newman and M. Szegedy, Regular languages are testable with a constant number of queries, Proc. 40th FOCS, New York, NY, IEEE (1999), 645-655. Also: SIAM J. on Computing 30 (2001), 1842-1862.
- [5] N. Alon and M. Krivelevich, Testing k-colorability, SIAM J. Discrete Math., 15 (2002), 211-227.
- [6] N. Alon and A. Shapira, Every monotone graph property is testable, Proc. of STOC 2005, 128-137.
- [7] N. Alon and A. Shapira, A characterization of the (natural) graph properties testable with one-sided error, Proc. of FOCS 2005.
- [8] M. Ben-Or, D. Coppersmith, M. Luby and R. Rubinfeld, Non-abelian homomorphism testing, and distributions close to their self-convolutions, Proc. of APPROX-RANDOM (2004), 273-285.
- [9] E. Ben-Sasson, P. Harsha and S. Raskhodnikova, Some 3-CNF properties are hard to test Proc. of STOC 2003, 345-354.
- [10] M. Blum, M. Luby and R. Rubinfeld, Self-testing/correcting with applications to numerical problems, JCSS 47 (1993), 549-595.
- [11] A. Czumaj, C. Sohler and M. Ziegler, Property testing in computational geometry, Proc. of ESA (2000), 155-166.
- [12] A. Czumaj and C. Sohler, Abstract combinatorial programs and efficient property testers, SIAM Journal on Computing 34 (2005), 580-615.
- [13] E. Fischer, Testing graphs for colorability properties, Proc. of the 12th SODA (2001), 873-882.
- [14] E. Fischer, The art of uninformed decisions: A primer to property testing, The Computational Complexity Column of The Bulletin of the European Association for Theoretical Computer Science 75 (2001), 97-126.

- [15] E. Fischer, The difficulty of testing for isomorphism against a graph that is given in advance, *SIAM Journal on Computing*, 34, 1147-1158.
- [16] E. Fischer, G. Kindler, D. Ron, S. Safra, and A. Samorodnitsky, Testing juntas, *JCSS* 68 (2004), 753-787. Also, *Proc. of FOCS* (2002), 103-112.
- [17] E. Fischer and I. Newman, Testing versus estimation of graph properties, *Proc. of 37th STOC*, 2005, 138-146.
- [18] E. Fischer, I. Newman and J. Sgall, Functions that have read-twice constant width branching programs are not necessarily testable, *Random Structures and Algorithms*, in press.
- [19] E. Fischer, E. Lehman, I. Newman, S. Raskhodnikova, R. Rubinfeld and A. Samorodnitsky, Monotonicity testing over general poset domains, *Proc. of STOC* (2002), 474-483.
- [20] K. Friedl, G. Ivanyos and M. Santha, Efficient testing of groups, *Proc. of STOC* (2005), 157-166.
- [21] A. Frieze and R. Kannan, Quick approximation to matrices and applications, *Combinatorica* 19 (1999), 175-220.
- [22] O. Goldreich, Combinatorial property testing - a survey, In: *Randomization Methods in Algorithm Design* (P. Pardalos, S. Rajasekaran and J. Rolim eds.), AMS-DIMACS (1998), 45-60.
- [23] O. Goldreich, Contemplations on testing graph properties, manuscript, 2005.
- [24] O. Goldreich, S. Goldwasser and D. Ron, Property testing and its connection to learning and approximation, *Proc. of 37th Annual IEEE FOCS*, (1996), 339-348. Also, *JACM* 45(4): 653-750 (1998).
- [25] O. Goldreich and L. Trevisan, Three theorems regarding testing graph properties, *Proc. 42nd IEEE FOCS*, IEEE (2001), 460-469. Also, *Random Structures and Algorithms*, 23(1):23-57, 2003.
- [26] O. Goldreich and D. Ron, Property Testing in Bounded-Degree Graphs, *Proc. of STOC 1997*, 406-415.
- [27] J. Komlós and M. Simonovits, Szemerédi's Regularity Lemma and its applications in graph theory. In: *Combinatorics, Paul Erdős is Eighty*, Vol II (D. Miklós, V. T. Sós, T. Szönyi eds.), János Bolyai Math. Soc., Budapest (1996), 295-352.
- [28] L. Lovász and B. Szegedy, Graph limits and testing hereditary graph properties, manuscript, 2005.
- [29] I. Newman, Testing of functions that have small width branching programs, *Proc. of 41th FOCS* (2000), 251-258.
- [30] M. Parnas and D. Ron, Testing the diameter of graphs, *Random structures and algorithms*, 20 (2002), 165-183.
- [31] M. Parnas, D. Ron and R. Rubinfeld, Testing membership in parenthesis languages, *Random Struct. and Algorithms* 22 (2003), 98-138.

- [32] V. Rödl and R. Duke, On graphs with small subgraphs of large chromatic number, *Graphs and Combinatorics* 1 (1985), 91–96.
- [33] D. Ron, Property testing, in: P. M. Pardalos, S. Rajasekaran, J. Reif and J. D. P. Rolim, editors, *Handbook of Randomized Computing*, Vol. II, Kluwer Academic Publishers, 2001, 597–649.
- [34] R. Rubinfeld and M. Sudan, Robust characterization of polynomials with applications to program testing, *SIAM J. on Computing* 25 (1996), 252–271.
- [35] E. Szemerédi, Regular partitions of graphs, In: *Proc. Colloque Inter. CNRS* (J. C. Bermond, J. C. Fournier, M. Las Vergnas and D. Sotteau, eds.), 1978, 399–401.

9 Proofs from Section 3

For the proofs of this section we need the following large deviation inequality.

Lemma 9.1 *Suppose X_1, \dots, X_n are n independent Boolean random variables, where $\text{Prob}[X_i = 1] = p_i$. Let $E = \sum_{i=1}^n p_i$. Then, $\text{Prob}[|\sum_{i=1}^n X_i - E| \geq \delta n] \leq 2e^{-2\delta^2 n}$.*

Proof (of Lemma 3.2): Suppose that $d(A, B) = \eta + p$, where $|p| \leq \delta$, and assume for now that $p \geq 0$. Suppose first that $p \leq \delta(\gamma + 2\delta)^2$. In this case we just remove any $pm^2 (\leq \delta m^2)$ edges and thus make sure that $d(A, B) = \eta$. Furthermore, as for any pair (A', B') of size $(\gamma + 2\delta)m$ we initially had $d(A', B') = \eta + p \pm (\gamma + \delta)$, it is easy to see that because we remove $pm^2 \leq \delta(\gamma + 2\delta)^2 m^2$ edges, we now have $\eta - \gamma - 2\delta \leq d(A', B') \leq \eta + \gamma + \delta$, which satisfies $d(A', B') = \eta \pm (\gamma + 2\delta)$. Thus, in this case we turned (A, B) into a $(\gamma + 2\delta)$ -regular pair of density η .

Suppose now that $p \geq \delta(\gamma + 2\delta)^2$. Our way for turning (A, B) into a $(\gamma + 2\delta)$ -regular pair with density η will consist of two stages. In the first we will randomly remove some of the edges connecting A and B . We will then deterministically make some additional modifications. To get that after these two stages (A, B) has the required properties we show that with probability $3/4$ the pair (A, B) is $(\gamma + 2\delta)$ -regular and with the same probability $d(A, B) = \eta$. By the union bound we will get that with probability at least $1/2$ the pair (A, B) has the required two properties.

In the first (random) step, we remove each of the edges connecting A and B randomly and independently with probability $\frac{p}{\eta+p}$. Then, the expected number of edges removed is $\frac{p}{\eta+p}(\eta+p)|A||B| = p|A||B| \leq \delta|A||B|$, and the expected value of $d(A, B)$ is η . As we assumed that $p \geq \delta(\gamma + 2\delta)^2$ we have $d(A, B) \geq \delta(\gamma + 2\delta)^2$. Therefore, the number of edges we may randomly remove is at least $\delta(\gamma + 2\delta)^2 m^2$. Therefore, by Lemma 9.1, for large enough $m \geq m_{3.2}(\delta, \gamma)$, the probability that $d(A, B)$ deviates from η by more than $m^{-0.5}$ is at most $3/4$. In particular, the number of edge modifications made is at most $\frac{3}{2}\delta m^2$ with probability at least $3/4$. Now (this is the second, deterministic step) we can add or remove at most $m^{1.5}$ edges arbitrarily and thus make sure that $d(A, B) = \eta$. The total number of edge modifications is also at most $\frac{3}{2}\delta m^2 + m^{1.5} \leq 2\delta m^2$, for large enough $m \geq m_{3.2}(\delta, \gamma)$. Note that we have thus established that with probability at least $3/4$ after the above two stages $d(A, B) = \eta$.

As (A, B) was assumed to be $(\gamma + \delta)$ -regular, we initially had $d(A', B') = \eta + p \pm (\gamma + \delta)$ for any pair of subsets $A' \subseteq A$ and $B' \subseteq B$ of size $(\gamma + 2\delta)$. As in the first step we removed each edge with probability $\frac{p}{\eta+p}$, the expected value of $d(A'B')$ after the first step is between

$$(\eta + p + \gamma + \delta)\left(1 - \frac{p}{\eta + p}\right) \leq \eta + \gamma + \delta$$

and

$$(\eta + p - \gamma - \delta)\left(1 - \frac{p}{\eta + p}\right) \geq \eta - \gamma - \delta.$$

Recall that we have already established that with probability at least $3/4$ we have $d(A, B) = \eta$ and that for any pair (A', B') of size $(\gamma + 2\delta)m$ the expected value of $d(A', B')$ is $\eta \pm (\gamma + \delta)$. Hence, to show that after the two steps (A, B) is a $(\gamma + 2\delta)$ -regular pair with probability at least $1/2$, it suffices to show that with probability at least $3/4$, the densities of all pairs (A', B') do not deviate from their expectation by more than δ .

Suppose first that $d(A', B')$ was originally at most $\frac{1}{2}\delta$. This means that when we randomly remove edges from (A, B) we can change $d(A', B')$ by at most $\frac{1}{2}\delta$. Thus in this case $d(A', B')$ can deviate from its expectation by at most $\frac{1}{2}\delta$. Also, when adding or removing $m^{1.5}$ edges to (A, B) in the second step we can change $d(A', B')$ by at most $m^{-0.5}/(\gamma + 2\delta)^2 \leq \frac{1}{2}\delta$ for large enough $m \geq m_{3.2}(\delta, \gamma)$. Thus, for such pairs we are guaranteed that $d(A', B') = \eta \pm (\gamma + 2\delta)$.

Suppose now that $d(A', B')$ was at least $\frac{1}{2}\delta$. Thus the number of edges, which were considered for removal between A' and B' in the first step was at least $\frac{1}{2}\delta(\gamma + 2\delta)^2 m^2$. Hence, by Lemma 9.1 the probability that $d(A', B')$ deviates from its expectation by more than $\frac{1}{2}\delta$ is at most $2e^{-2(\frac{1}{2}\delta)^2 \frac{1}{2}\delta(\gamma + 2\delta)^2 m^2}$. Thus, as there are at most 2^{2m} pairs of such sets (A', B') , we conclude by the union-bound that for large enough $m \geq m_{3.2}(\delta, \gamma)$, with probability at least $3/4$ all sets (A', B') of size $(\gamma + 2\delta)m$ satisfy $d(A', B') = \eta \pm (\gamma + \frac{3}{2}\delta)$. As in the previous paragraph, adding or removing $m^{1.5}$ edges in the second step can change $d(A', B')$ by at most $\frac{1}{2}\delta$, so in this case we also have $d(A', B') = \eta \pm (\gamma + \frac{3}{2}\delta)$.

Finally, in the case that p above is negative we can use essentially the same argument. The only modification is that we add edges instead of remove them. \blacksquare

Proof (of Lemma 3.3): For any vertex $a \in A$ and $b \in B$ we do the following: we flip a coin with bias $\frac{2\delta}{(\delta + \gamma)}$. If the coin comes up heads we make no modification between the vertices a and b . If the coin comes up tails then we disregard the adjacency relation between a and b and do the following: we flip another coin with bias η . If the coin comes up heads then we connect a and b , and otherwise we leave them disconnected. In what follows we call the coins flipped in the first step the *first* coins, and those flipped in the second step the *second* coins.

Claim: With probability at least $3/4$, we make at most $\frac{3\delta}{\gamma}m^2$ edge modifications.

Proof. Note that the number of edge modifications is at most the number of first coins that came up heads. The distribution of these m^2 coins is given by the Binomial distribution $B(m^2, \frac{2\delta}{(\delta + \gamma)})$, whose expectation is $\frac{2\delta}{(\delta + \gamma)}m^2$, and by Lemma 9.1 the probability of deviating by more than $\frac{1}{2}\delta m^2$ from this expectation is at most $2e^{-2(\delta/2)^2 m^2}$. For large enough $m \geq m_{3.3}(\delta, \gamma)$ we get that with probability at least $3/4$ we make at most $\frac{2\delta}{(\delta + \gamma)}m^2 + \frac{1}{2}\delta m^2 \leq \frac{2.5\delta}{\gamma}m^2$ modifications. \blacksquare

The following observation will be useful for the next two claims: Fix a pair of connected vertices $a \in A$ and $b \in B$. For them to become disconnected both coins must come up tails, thus the probability of them staying connected is $(1 - \frac{2\delta}{(\delta + \gamma)} + \frac{2\eta\delta}{(\delta + \gamma)})$. Now, fix a pair of disconnected vertices $a \in A$ and $b \in B$. For them to become connected the first coin must come up tails and the second must come up heads, so the probability of them becoming connected is $\frac{2\eta\delta}{(\delta + \gamma)}$.

Claim: With probability at least $3/4$, we have $d(A, B) = \eta \pm m^{-0.5}$.

Proof. Recall that by assumption the number of connected vertices was ηm^2 . Thus, by the above observation the expected number of connected vertices is

$$\eta m^2 \left(1 - \frac{2\delta}{(\delta + \gamma)} + \frac{2\eta\delta}{(\delta + \gamma)}\right) + (1 - \eta)m^2 \frac{2\eta\delta}{(\delta + \gamma)} = \eta m^2.$$

By Lemma 9.1 we get that for large enough $m \geq m_{3.3}(\delta, \gamma)$ the probability of deviating from this expectation by more than $m^{-0.5}$ is at most $1/4$. \blacksquare

Claim: With probability at least $3/4$, all sets $A' \subseteq A$ and $B' \subseteq B$ of size γm satisfy $d(A', B') = \eta \pm (\gamma - \frac{1}{2}\delta)$.

Proof. Fix any pair of such sets. Let e denote the number of edges originally spanned by these sets. As in the previous claim we get that the expected number of edges spanned by (A', B') is

$$e\left(1 - \frac{2\delta}{(\delta + \gamma)} + \frac{2\eta\delta}{(\delta + \gamma)}\right) + (|A'||B'| - e)\frac{2\eta\delta}{(\delta + \gamma)} = e\left(1 - \frac{2\delta}{(\delta + \gamma)}\right) + |A'||B'|\frac{2\eta\delta}{(\delta + \gamma)}.$$

Recall that by assumption $e = |A'||B'|(\eta \pm (\gamma + \delta))$. Thus, the expected number of edges spanned by (A', B') is at most

$$\begin{aligned} |A'||B'|(\eta + \gamma + \delta)\left(1 - \frac{2\delta}{(\delta + \gamma)}\right) + |A'||B'|\frac{2\eta\delta}{(\delta + \gamma)} &= |A'||B'|(\eta + \gamma + \delta - \frac{2\delta\gamma}{\delta + \gamma} - \frac{2\delta^2}{\delta + \gamma}) = \\ &= |A'||B'|(\eta + \gamma - \delta), \end{aligned}$$

Similarly, we infer that the expected number of edges spanned by (A', B') is at least

$$\begin{aligned} |A'||B'|(\eta - \gamma - \delta)\left(1 - \frac{2\delta}{(\delta + \gamma)}\right) + |A'||B'|\frac{2\eta\delta}{(\delta + \gamma)} &= |A'||B'|(\eta - \gamma - \delta + \frac{2\delta\gamma}{\delta + \gamma} + \frac{2\delta^2}{\delta + \gamma}) = \\ &= |A'||B'|(\eta - \gamma + \delta). \end{aligned}$$

By Lemma 9.1 the probability that the number of edges between A' and B' will deviate from its expectation by more than $\frac{1}{2}\delta|A'||B'|$ is at most $2e^{-2(\delta/2)^2|A'||B'|} = 2e^{-2(\delta/2)^2(\gamma m)^2}$. As the number of pairs (A', B') is at most 2^{2m} we get by the union bound, provided that $m \geq m_{3.3}(\delta, \gamma)$ is large enough, that with probability at least $3/4$ all the pairs (A', B') of size γm satisfy this property. Thus for all pairs (A', B') of size γm we have $d(A', B') = \eta \pm (\gamma - \frac{1}{2}\delta)$. ■

Combining the above three claims we get that with constant probability we make at most $\frac{2.5\delta}{\gamma}m^2$ modifications and thus make sure that $d(A, B) = \eta \pm m^{-0.5}$ and furthermore that for any pair of sets (A', B') of size γm we have $d(A', B') = \eta \pm (\gamma - \frac{1}{2}\delta)$. Now we can add or remove at most $m^{1.5}$ edges to make sure that $d(A, B) = \eta$. For any pair of sets (A', B') of size γm this will change $d(A', B')$ by at most $m^{-0.5}/\gamma^2 \leq \frac{1}{2}\delta$ for large enough m . This means that we will have $d(A', B') = \eta \pm \gamma$, implying that (A, B) is γ -regular with density η , completing the proof of the lemma. ■

Proof (of Lemma 3.1): By Lemma 3.2 we can make at most $2\delta m^2$ edge modifications and thus turn (A, B) into a $(\gamma + 2\delta)$ -regular pair with density η . Thus, every pair of subsets $A'' \subseteq A$ and $B'' \subseteq B$ of size γm has density at most

$$(\eta + \gamma + 2\delta)(\gamma + 2\delta)^2 m^2 / \gamma^2 m^2 \leq (\eta + \gamma + 2\delta)(1 + 8\delta/\gamma) \leq \eta + \gamma + 14\delta/\gamma.$$

Similarly, the density of such a pair is at least $\eta - \gamma - 14\delta/\gamma$. We thus conclude that (A, B) has density precisely η , and every pair of subsets (A'', B'') of size γm has density $\eta \pm (\gamma + 14\delta/\gamma)$. Now we can use Lemma 3.3 to make at most $3\frac{14\delta/\gamma}{\gamma}m^2 = 42\frac{\delta}{\gamma^2}m^2$ additional edge modifications and thus turn (A', B') into γ -regular pair with density precisely η . The total number of modifications is $42\frac{\delta}{\gamma^2}m^2 + 2\delta m^2 \leq 50\frac{\delta}{\gamma^2}m^2$ as needed. ■

The following application of Lemma 3.1 will be useful later in the paper.

Lemma 9.2 *Let R be a regularity-instance of order k , error-parameter γ , $\binom{k}{2}$ edge densities $\eta_{i,j}$ and set of non-regular pairs \bar{R} . If a graph G has an equipartition $\mathcal{V} = \{V_1, \dots, V_k\}$ of order k such that*

1. $d(V_i, V_j) = \eta_{i,j} \pm \frac{\gamma^2 \epsilon}{50}$ for all $i < j$.
2. Whenever $(i, j) \notin \bar{R}$, the pair (V_i, V_j) is $(\gamma + \frac{\gamma^2 \epsilon}{50})$ -regular.

Then G is ϵ -close to satisfying R .

Proof (of Lemma 9.2): For any $(i, j) \notin \bar{R}$ we can use Lemma 3.1 and make at most $50 \frac{\gamma^2 \epsilon / 50}{\gamma^2} (n/k)^2 \leq \epsilon n^2 / k^2$ edge modifications to turn (V_i, V_j) into a γ -regular pair with density $\eta_{i,j}$. As there are at most $\binom{k}{2}$ pairs this is a total of at most ϵn^2 modifications. We have thus turned G into a graph satisfying R by making at most ϵn^2 edge modifications, as needed. ■

10 Proofs from Section 4

Definition 10.1 *Let H be a graph on h vertices, let W be a weighted complete graph on h vertices, where the weight of edge (i, j) is $\eta_{i,j}$. For a permutation $\sigma : [h] \rightarrow [h]$ define*

$$IC(H, W, \sigma) = \prod_{(i,j) \in E(H)} \eta_{\sigma(i), \sigma(j)} \prod_{(i,j) \notin E(H)} (1 - \eta_{\sigma(i), \sigma(j)})$$

Suppose V_1, \dots, V_k are k vertex sets, each of size m , and suppose the bipartite graph spanned by V_i and V_j is a bipartite random graph with edge density $\eta_{i,j}$. Let H be a graph of size k , and let $\sigma : [k] \rightarrow [k]$ be some permutation. What is the expected number of k -tuples of vertices $v_1 \in V_1, \dots, v_k \in V_k$, which span an induced copy of H with v_i playing the role of $\sigma(i)$? It is easy to see that the answer is $IC(H, W, \sigma)m^k$, where W is the weighted complete graph with weights $\eta_{i,j}$. The following claim shows that this is approximately the case when instead of random bipartite graphs we take regular enough bipartite graphs. The proof is a standard application of the definition of a regular pair and is thus omitted from this extended abstract. See Lemma 4.2 in [15] for a version of the proof.

Claim 10.2 *For any δ and h , there exists a $\gamma = \gamma_{10.2}(\delta, h)$ such that the following holds: Suppose V_1, \dots, V_h are h sets of vertices of size m each, and that all the pairs (V_i, V_j) are γ -regular. Define W to be the weighted complete graph on h vertices, whose weights are $\eta_{i,j} = d(V_i, V_j)$. Then, for any graph H on h vertices and for any $\sigma : [h] \rightarrow [h]$, the number of h -tuples $v_1 \in V_1, \dots, v_h \in V_h$, which span an induced copy of H with v_i playing the role of vertex $\sigma(i)$ is*

$$(IC(H, W, \sigma) \pm \delta)m^h$$

We would now want to consider the total number of induced copies of some graph.

Definition 10.3 *Let H be a graph on h vertices, let W be a weighted complete graph on h vertices, where the weight of edge (i, j) is $\eta_{i,j}$. Let $Aut(H)$ denote the number of automorphisms of H . Define*

$$IC(H, W) = \frac{1}{Aut(H)} \sum_{\sigma} IC(H, W, \sigma).$$

Continuing the discussion before Claim 10.2, it is easy to see that in this case the expected number of induced copies of H having one vertex in each of the sets V_i is $IC(H, W)$. Again, we can show that the same is approximately true when we replace random bipartite graphs with regular enough bipartite graphs.

Claim 10.4 *For any δ and k , there exists a $\gamma = \gamma_{10.4}(\delta, k)$ such that the following holds: Suppose that V_1, \dots, V_k are sets of vertices of size m each, and that all the pairs (V_i, V_j) are γ -regular. Define K to be the weighted complete graph on k vertices, whose weights are $\eta_{i,j} = d(V_i, V_j)$. Then, for any graph H of size k , the number of induced copies of H , which have precisely one vertex in each of the sets V_1, \dots, V_k is*

$$(IC(H, W) \pm \delta)m^k$$

Proof. Set $\gamma_{10.4}(\delta, k) = \gamma_{10.2}(\delta/k!, k)$. Suppose V_1, \dots, V_k are as in the statement of the claim and let H be any graph on k vertices. By Claim 10.2 for every permutation $\sigma : [k] \rightarrow [k]$, the number of induced copies of H which have precisely one vertex v_i in each set V_i such that v_i plays the role of vertex $\sigma(i)$ is $IC(H, W, \sigma) \pm \delta m^k / k!$. If we sum over all permutations $\sigma : [k] \rightarrow [k]$ we get $\sum_{\sigma} (IC(H, W, \sigma) \pm \delta / k!) m^k$. This summation, however, counts copies of H several times. More precisely, each copy is thus counted $Aut(H)$ times. Thus, dividing by $Aut(H)$ gives that the number of such induced copies is

$$\frac{1}{Aut(H)} \left(\sum_{\sigma} (IC(H, W, \sigma) \pm \delta / k!) m^k \right) = \left(\frac{1}{Aut(H)} \sum_{\sigma} IC(H, W, \sigma) \pm \delta \right) m^k = (IC(H, W) \pm \delta) m^k.$$

■

We would now want to consider the number of induced copies of a graph H , when the number of sets V_i is larger than the size of H .

Definition 10.5 *Let H be a graph on h vertices, let R be a weighted complete graph of size at least h where the weight of edge (i, j) is $\eta_{i,j}$, and let \mathcal{W} denote all the subsets of $V(W)$ of size h . Define*

$$IC(H, R) = \sum_{W \in \mathcal{W}} IC(H, W).$$

The following lemma shows that knowing that a graph satisfies some regularity-instance R , enables us to estimate the number of induced copies spanned by any graph, which satisfies R .

Lemma 10.6 *For any δ and q , there are $k = k_{10.6}(\delta, q)$ and $\gamma = \gamma_{10.6}(\delta, q)$ with the following properties: For any regularity-instance R of order at least k and with error parameter at most γ , and for every graph H of size $h \leq q$, the number of induced copies of H in any n -vertex graph satisfying R is*

$$(IC(H, R) \pm \delta) \binom{n}{h}$$

Proof. Put $k = k_{10.6}(\delta, q) = \frac{\delta}{10q^2}$ and $\gamma = \gamma_{10.6}(\delta, q) = \min\{\frac{\delta}{3q^2}, \gamma_{10.4}(\frac{1}{3}\delta, q)\}$. Let R be any regularity instance as in the statement, let G be any graph satisfying R , and let H be any graph of size $h \leq q$. Let V_1, \dots, V_ℓ be an equipartition of G satisfying R . For the proof of the lemma it will

be simpler to consider an equivalent statement of the lemma, stating that if one samples an h -tuple of vertices from G , then the probability that it spans an induced copy of H is $IC(H, R) \pm \delta$.

First, note that by our choice of k we get from a simple birthday-paradox argument, that the probability that the h -tuple of vertices has more than one vertex in one of the sets V_i is at most $\frac{1}{3}\delta$. Second, observe that as the equipartition of R is γ -regular and $\gamma \leq \delta$ we get that the probability that the h -tuple of vertices contains a pair $v_i \in V_i$ and $v_j \in V_j$ such that (V_i, V_j) is not γ -regular is at most $\binom{h}{2}\gamma \leq \binom{q}{2}\gamma \leq \frac{1}{3}\delta$. Thus, it is enough to show that conditioning on the events: (i) the h vertices v_1, \dots, v_h belong to distinct sets V_i , (ii) if $v_i \in V_i$ and $v_j \in V_j$ then (V_i, V_j) is γ -regular, then the probability that they span an induced copy of H is $IC(H, R) \pm \frac{1}{3}\delta$. Assuming events (i) and (ii) hold let us compute the probability that the h -tuple of vertices spans an induced copy of H , while conditioning on the h sets from V_1, \dots, V_ℓ which contain the h vertices. For every possible set W of h sets V_i we get from the choice of γ via Claim 10.4 that the probability that they span an induced copy of H is $IC(H, W) \pm \frac{1}{3}\delta$. This means that the probability that the h -tuple of vertices span an induced copy of H is $IC(H, R) \pm \frac{1}{3}\delta$, as needed. \blacksquare

Proof (of Lemma 4.1): Suppose \mathcal{P} is testable by a tester T , and assume without loss of generality that T is canonical. This assumption is possible by Lemma 4.2. Let $q(\epsilon)$ be the upper bound guarantee for the query complexity of T . Fix any n and δ and assume that $\delta < 1/12$ (otherwise, replace δ with $1/13$). Let $q = q(\delta, n) \leq q(\delta)$ be the query complexity, which is sufficient for T to distinguish between n -vertex graphs satisfying \mathcal{P} from those that are δ -far from satisfying it, with success probability at least $2/3$. As T is canonical, if it samples a set of vertices and gets a graph of size q , it either rejects or accepts deterministically. Hence, we can define a set \mathcal{A} , of all the graphs Q of size q , such that if the sample of vertices spans a graph isomorphic to Q , then T accepts the input. We finally put $k = k_{10.6}(\delta/2^{\binom{q}{2}}, q)$, $\gamma = \gamma_{10.6}(\delta/2^{\binom{q}{2}}, q)$ and $T = T_{2.3}(k, \gamma)$. For any $k \leq t \leq T$ consider all the (finitely many) regularity-instances of order t , where for the edge densities $\eta_{i,j}$ we choose a real from the set $\{0, \frac{\delta\gamma^2}{50q^2}, 2\frac{\delta\gamma^2}{50q^2}, 3\frac{\delta\gamma^2}{50q^2}, \dots, 1\}$. Let \mathcal{I} be the union of all these regularity-instances. Note, that all the above constants, as well as the size of \mathcal{I} and the complexity of the regularity-instances in \mathcal{I} are determined as a function of δ only (and the property \mathcal{P}).

We claim that we can take \mathcal{R} in Definition 2.5 to be

$$\mathcal{R} = \{R \in \mathcal{I} : \sum_{H \in \mathcal{A}} IC(R, H) \geq 1/2\}.$$

To see this, first note that the expression $\sum_{H \in \mathcal{A}} IC(R, H)$ is an estimation of the fraction of induced copies of graphs from \mathcal{A} in a graph satisfying R . Combining the facts that the graphs in \mathcal{A} all have size q and we use Lemma 10.6 with $\delta/2^{\binom{q}{2}}$ we infer that the expression $\sum_{H \in \mathcal{A}} IC(R, H)$ is an estimate of the number of induced copies of graphs from \mathcal{A} in a graph satisfying R , up to an additive error of at most $\delta \binom{n}{q}$.

Suppose a graph G satisfies \mathcal{P} . This means that T accepts G with probability at least $2/3$. In other words, this means that at least $\frac{2}{3} \binom{n}{q}$ of the subsets of q vertices of G span a graph isomorphic to one of the members of \mathcal{A} . By Lemma 2.3 G has some γ -regular partition of size at least k and at most T . As the densities in the regularity-instances in \mathcal{A} differ by $\frac{\delta\gamma^2}{50q^2}$ we get that the densities of the regular partition of G differ by at most $\frac{\delta\gamma^2}{50q^2}$ from the densities of one of the regularity-instances $R \in \mathcal{I}$. Lemma 9.2 implies that G is δ/q^2 -close to satisfying one of the regularity-instances of \mathcal{I} . Note that adding-removing an edge can decrease the number of induced copies of members of \mathcal{A} in G

by at most $\binom{n-2}{q-2}$. Thus adding/removing $\delta n^2/q^2$ edges can decrease the number of induced copies of members of \mathcal{A} in G by at most $\delta \frac{n^2}{q^2} \binom{n-2}{q-2} \leq \delta \binom{n}{q}$. Thus, after these at most $\delta n^2/q^2$ edge modifications we get a graph that satisfies one of the regularity-instances $R \in \mathcal{I}$ and at least $(\frac{2}{3} - \delta) \binom{n}{q} > (\frac{1}{2} + \delta) \binom{n}{q}$ of the subsets of q vertices of the new graph span a member of \mathcal{A} (here we use the assumption that $\delta < 1/12$). As explained in the previous paragraph, by our choice of k and γ via Lemma 10.6, this means that $\sum_{H \in \mathcal{A}} IC(R, H) \geq 1/2$. By definition of \mathcal{R} this means that $R \in \mathcal{R}$, thus G is indeed δ -close to satisfying one of the regularity-instances of \mathcal{R} .

Suppose now that a graph G is ϵ -far from satisfying \mathcal{P} . If $\delta \geq \epsilon$ then there is nothing to prove, so assume that $\delta < \epsilon$. If G is $(\epsilon - \delta)$ -close to satisfying a regularity-instance $R \in \mathcal{R}$, then by definition of \mathcal{R} and our choice of k and γ via Lemma 10.6 it is $(\epsilon - \delta)$ -close to a graph G' , such that at least $(\frac{1}{2} - \delta) \binom{n}{q} > (\frac{1}{3} + \delta) \binom{n}{q}$ of the subsets of q vertices of G' span an induced copy of a graph from \mathcal{A} . In other words, this means that T accepts G' with probability at least $\frac{1}{3} + \delta$. This means that G' cannot be δ -far from satisfying \mathcal{P} as we assume that q is enough for T to reject with probability at least $2/3$ graphs that are δ -far from satisfying \mathcal{P} . However, as G is ϵ -far from satisfying \mathcal{P} any graph that is $(\epsilon - \delta)$ -close to G must be δ -far from satisfying \mathcal{P} . ■

11 Proofs from Section 5

Proof (of Lemma 5.5): Let $\bar{A}_1, \dots, \bar{A}_k$ be any equipartition of \bar{G} . Recall that ℓ denotes the order of the equipartition $\bar{\mathcal{V}}$, which is also the order of \mathcal{V} . For every $1 \leq p \leq \ell$ and $1 \leq q \leq k$ set $\bar{A}\bar{V}_{p,q} = \bar{V}_p \cap \bar{A}_q$ and $\alpha_{p,q} = |\bar{A}\bar{V}_{p,q}|/|\bar{V}_p|$. For every $1 \leq p \leq \ell$ and $1 \leq q \leq k$ let $AV_{p,q}$ be **any** subset of V_p of size $\alpha_{p,q}|V_p|$. Finally for every $1 \leq q \leq k$ define $A_q = \bigcup_{p=1}^{\ell} AV_{p,q}$. Instead of stating what $\zeta_{5.5}(k, \delta)$ should be, we state along the way different upper bound on $\zeta_{5.5}(k, \delta)$ that will depend only on k and δ . One can then take the minimum of all these values as $\zeta_{5.5}(k, \delta)$

Claim 1: If $(\bar{A}_q, \bar{A}_{q'})$ is γ -regular then $(A_q, A_{q'})$ is $(\gamma + \delta)$ -regular.

Proof. To simplify the notation we assume that (\bar{A}_1, \bar{A}_2) is γ -regular and prove that (A_1, A_2) is $(\gamma + 2\delta)$ -regular. Set $\eta = d(\bar{A}_1, \bar{A}_2)$. As Claim 2 below asserts $d(A_1, A_2) = \eta \pm \delta$. Thus we need to show that $d(A'_1, A'_2) = \eta \pm (\gamma + \delta)$ for every $A'_1 \subseteq A_1$ and $A'_2 \subseteq A_2$ of sizes $(\gamma + \delta)|A_1|$ and $(\gamma + \delta)|A_2|$, respectively. For simplicity we show that $d(A'_1, A'_2) \leq \eta + \gamma + \delta$, as showing that $d(A'_1, A'_2) \geq \eta - \gamma - \delta$ is similar. Recall that each set A_q is the union of ℓ sets $AV_{1,q}, \dots, AV_{\ell,q}$. For every $1 \leq i, j \leq \ell$ put $AV'_{i,1} = AV_{i,1} \cap A'_1$ and $AV'_{j,2} = AV_{j,2} \cap A'_2$. We can rephrase our goal in terms of the number of edges as follows

$$\sum_{1 \leq i, j \leq \ell} e(AV'_{i,1}, AV'_{j,2}) \leq (\eta + \gamma + \delta)|A'_1||A'_2| = (\eta + \gamma + \delta)(\gamma + \delta)^2|A_1||A_2|. \quad (1)$$

Let n denote the number of vertices of G . To prove (1) we turn to bound the contribution to the LHS (= Left Hand Side) of (1) of three types of pairs of (i, j) :

- **Pairs (i, j) for which $i = j$:** Observe that the maximum possible number of edges connecting all pairs $(AV_{i,1}, AV_{j,2})$ for which $i = j$ is given by $\sum_i \alpha_{i,1} \alpha_{i,2} |A_1||A_2|$. Furthermore, for any $1 \leq i \leq \ell$ we have $0 \leq \alpha_{i,1}, \alpha_{i,2} \leq k/\ell$ (this is because $|V_1| = \dots = |V_\ell| = n/\ell$ and $|A_1| = \dots = |A_k| = n/k$). By Claim 11.1 we get that $\sum_i \alpha_{i,1} \alpha_{i,2} |A_1||A_2| \leq \frac{k}{\ell} |A_1||A_2|$ and if we choose a

ζ satisfying $\ell \geq 1/\zeta \geq 6k/\delta^3 \geq 6k/\delta(\gamma + \delta)^2$ we can infer that the contribution of the pairs (i, i) to the LHS of (1) is at most $\frac{1}{6}\delta(\gamma + \delta)^2|A_1||A_2|$ (note that $\ell \geq 1/\zeta$ is guaranteed by the statement of the lemma).

- **Pairs (i, j) for which either $|AV'_{i,1}| < \zeta|V_i|$ or $|AV'_{j,2}| < \zeta|V_j|$:** Consider the $1 \leq i \leq \ell$ in (1) for which $|AV'_{i,1}| < \zeta|V_i| = \zeta n/\ell$. The total number of vertices of G that belong to such sets is clearly at most ζn , therefore the total number of such vertices in A_1 is at most $k\zeta|A_1|$. Similarly, the total number of vertices of A_2 which belong to sets $|AV'_{j,2}|$ for which $|AV'_{j,2}| < \zeta|V_j|$ is at most $k\zeta|A_2|$. Therefore the contribution of pairs (i, j) to the LHS of (1) for which either $|AV'_{i,1}| < \zeta|V_i|$ or $|AV'_{j,2}| < \zeta|V_j|$ is at most $2k\zeta|A_1||A_2|$. If we choose ζ so that it satisfies $\zeta \leq \frac{\delta^3}{12k} \leq \frac{\delta(\gamma+\delta)^2}{12k}$, such pairs (i, j) can contribute to the LHS of (1) a total of at most $\frac{1}{6}\delta(\gamma + \delta)^2|A_1||A_2|$.

For a later step of the proof it will be important to note that by the above reasoning, the number of vertices of A'_1 that belong to sets $AV'_{i,1}$ of size smaller than $\zeta|V_i|$ is at most $\delta|A_1|$. Similarly the number of vertices of A'_2 that belong to sets $AV'_{j,2}$ of size smaller than $\zeta|V_j|$ is at most $\delta|A_2|$.

- **Pairs (i, j) for which (V_i, V_j) is not ζ -regular:** Recall, that \mathcal{V} is a ζ -regular equipartition therefore at most ζn^2 edges of G connect pairs of clusters (V_i, V_j) that are not ζ -regular. As $|A_1| = |A_2| = n/k$ this means that the number of edges connecting A_1 and A_2 that belong to pairs (V_i, V_j) that are not ζ -regular is at most $k^2\zeta(n/k)^2 = k^2\zeta|A_1||A_2|$. If we choose ζ so that $\zeta \leq \frac{1}{6}\delta^3/k^2 \leq \frac{1}{6}\delta(\gamma + \delta)^2/k^2$, such pairs can contribute at most $\frac{1}{6}\delta(\gamma + \delta)^2|A_1||A_2|$ to the sum in (1).

We have thus accounted for all pairs (i, j) in (1) for which either $i = j$, (V_i, V_j) is not ζ -regular, $|AV'_{i,1}| < \zeta|V_i|$ or $|AV'_{j,2}| < \zeta|V_j|$. Specifically, we have shown that they can contribute at most $\frac{1}{2}\delta(\gamma + \delta)^2|A_1||A_2| = \frac{1}{2}\delta|A'_1||A'_2|$ to the LHS of (1). Therefore, we can now reduce proving (1) to showing that

$$\sum_{i \in I, j \in J, i \neq j} e(AV'_{i,1}, AV'_{j,2}) = \sum_{i \in I, j \in J, i \neq j} d(AV'_{i,1}, AV'_{j,2})|AV'_{i,1}||AV'_{j,2}| \leq (\eta + \gamma + \frac{1}{2}\delta)|A'_1||A'_2|, \quad (2)$$

while assuming that all $i \in I$ and $j \in J$ in the above sum satisfy $|AV'_{i,1}| \geq \zeta|V_i|$ and $|AV'_{j,2}| \geq \zeta|V_j|$. Note, that the lemma assumes that if (V_i, V_j) is ζ -regular then so is (\bar{V}_i, \bar{V}_j) . Therefore we can assume that for any $i \in I, j \in J, i \neq j$

$$d(AV'_{i,1}, AV'_{j,2}) = d(V_i, V_j) \pm \zeta. \quad (3)$$

and

$$d(\bar{AV}'_{i,1}, \bar{AV}'_{j,2}) = d(\bar{V}_i, \bar{V}_j) \pm \zeta. \quad (4)$$

The reason is that if $i < j$ is such that (V_i, V_j) and (\bar{V}_i, \bar{V}_j) are ζ -regular and furthermore $|AV'_{i,1}| \geq \zeta|V_i|$ and $|AV'_{j,2}| \geq \zeta|V_j|$ then the above follows from the definition of a ζ -regular pair. If one of these conditions does not hold then we will possibly recount some of the edges, which we have already accounted for before. If we choose ζ so that $\zeta \leq \frac{1}{6}\delta$ we can use (3) to reduce (2) to showing

$$\sum_{i \in I, j \in J, i \neq j} d(V_i, V_j)|AV'_{i,1}||AV'_{j,2}| \leq (\eta + \gamma + \frac{2}{3}\delta)|A'_1||A'_2| \quad (5)$$

As we assume that \mathcal{V} and $\bar{\mathcal{V}}$ are (ζ, ζ) -similar we have $d(V_i, V_j) = d(\bar{V}_i, \bar{V}_j) \pm \zeta$ for every $i < j$. If we choose ζ so that $\zeta \leq \frac{1}{6}\delta$, we can reduce (5) to showing that

$$\sum_{i \in I, j \in J, i \neq j} d(\bar{V}_i, \bar{V}_j) |AV'_{i,1}| |AV'_{j,2}| \leq (\eta + \gamma + \frac{1}{3}\delta) |A'_1| |A'_2| \quad (6)$$

By (4) we can reduce (6) to showing that

$$\sum_{i \in I, j \in J, i \neq j} d(\overline{AV'_{i,1}}, \overline{AV'_{j,2}}) |AV'_{i,1}| |AV'_{j,2}| \leq (\eta + \gamma) |A'_1| |A'_2|. \quad (7)$$

Let $A''_1 = \bigcup_{i \in I} AV'_{i,1}$ and $A''_2 = \bigcup_{j \in J} AV'_{j,2}$. Clearly $|A''_1| \leq |A'_1|$ and $|A''_2| \leq |A'_2|$, thus we can prove (7) by deriving the following stronger assertion:

$$\sum_{i \in I, j \in J, i \neq j} d(\overline{AV'_{i,1}}, \overline{AV'_{j,2}}) |AV'_{i,1}| |AV'_{j,2}| \leq (\eta + \gamma) |A''_1| |A''_2|. \quad (8)$$

Note, that as we have already mentioned, by our choice of ζ at most $\delta|A_1|$ vertices of A'_1 belong to sets $AV'_{i,1}$ for which $AV'_{i,1} < \zeta|V_1|$. Therefore, we have $|A''_1| \geq |A'_1| - \delta|A_1| \geq \gamma|A_1|$. Similarly, $|A''_2| \geq \gamma|A_2|$. Put $\beta_{i,1} = |AV'_{i,1}|/|A''_1|$ and $\beta_{j,2} = |AV'_{j,2}|/|A''_2|$. For every $i \in I$ let $\overline{AV'_{i,1}}$ be any subset of $\overline{AV'_{i,1}}$ of size $\beta_{i,1}|\overline{AV'_{i,1}}|$. Similarly, for every $j \in J$ let $\overline{AV'_{j,2}}$ be any subset of $\overline{AV'_{j,2}}$ of size $\beta_{j,2}|\overline{AV'_{j,2}}|$. Put $\bar{A}''_1 = \bigcup_{i \in I} \overline{AV'_{i,1}}$ and $\bar{A}''_2 = \bigcup_{j \in J} \overline{AV'_{j,2}}$ and note that just as $|A''_1| \geq \gamma|A_1|$ and $|A''_2| \geq \gamma|A_2|$ we also have $|\bar{A}''_1| \geq \gamma|\bar{A}_1|$ and $|\bar{A}''_2| \geq \gamma|\bar{A}_2|$. Dividing by $|A''_1||A''_2|$ we can restate (8) as

$$\sum_{i \in I, j \in J, i \neq j} d(\overline{AV'_{i,1}}, \overline{AV'_{j,2}}) \beta_{i,1} \beta_{j,2} \leq \eta + \gamma.$$

Finally, note that the above holds because

$$\sum_{i \in I, j \in J, i \neq j} d(\overline{AV'_{i,1}}, \overline{AV'_{j,2}}) \beta_{i,1} \beta_{j,2} \leq \sum_{1 \leq i, j \leq \ell} d(\overline{AV'_{i,1}}, \overline{AV'_{j,2}}) \beta_{i,1} \beta_{j,2} = d(\bar{A}''_1, \bar{A}''_2) \leq \eta + \gamma$$

due to the fact that (\bar{A}_1, \bar{A}_2) is by assumption γ -regular, $d(\bar{A}_1, \bar{A}_2) = \eta$, $|\bar{A}''_1| \geq \gamma|\bar{A}_1|$ and $|\bar{A}''_2| \geq \gamma|\bar{A}_2|$. This completes the proof of the claim. \blacksquare

Claim 2: For all $q < q'$ $d(A_q, A_{q'}) = d(\bar{A}_q, \bar{A}_{q'}) \pm \delta$

Proof. The proof is identical to the above proof. \blacksquare

The proof of the lemma follows from the above two claims. \blacksquare

Claim 11.1 Let a_1, \dots, a_ℓ and b_1, \dots, b_ℓ satisfy $\sum_{1 \leq i \leq \ell} a_i = \sum_{1 \leq i \leq \ell} b_i = 1$ and $0 \leq a_i, b_i \leq k/\ell$, where $k \leq \ell$. Then $\sum_{1 \leq i \leq \ell} a_i b_i \leq k/\ell$.

Proof. Observe that $\sum_{1 \leq i \leq \ell} a_i b_i \leq \max_{1 \leq i \leq \ell} \{a_i\} \sum_{1 \leq i \leq \ell} b_i \leq k/\ell$. ■

Proof (of Lemma 5.2): Set $\zeta = (\zeta_{5.5}(k, \delta)/50)^5$ and $\zeta' = 50\zeta^{1/5}$ and note that $\zeta, \zeta' \leq \zeta_{5.5}(k, \delta)$. Let $\mathcal{V} = \{V_1, \dots, V_\ell\}$ be a ζ -regular partition of G of order at least $1/\zeta$. Such an equipartition of order at most $T_{2.3}(1/\zeta, \zeta)$ exists by Lemma 2.3. By Lemma 5.3 we get that if we sample a set Q of at least $q_{5.3}(\ell, \zeta')$ vertices from G then with probability at least $2/3$ the graph induced on Q , which we denote by $G[Q]$ will have an equipartition $\mathcal{U} = \{U_1, \dots, U_\ell\}$, such that $d(V_i, V_j) = d(U_i, U_j) \pm \zeta'$ and such that if (V_i, V_j) is ζ -regular then (U_i, U_j) is ζ' -regular. This means that with probability at least $2/3$, the graph $G[Q]$ is such that G and $G[Q]$ have equipartitions, which are $(\zeta_{5.5}(k, \delta), \zeta_{5.5}(k, \delta))$ -similar. As these equipartition we can take \mathcal{V} and \mathcal{U} , because as $\zeta' \leq \zeta_{5.5}(k, \delta)$ then $d(V_i, V_j) = d(U_i, U_j) \pm \zeta_{5.5}(k, \delta)$. Also, as $\zeta \leq \zeta' \leq \zeta_{5.5}(k, \delta)$, then for all but at most $\zeta_{5.5}(k, \delta) \binom{k}{2}$ of the pairs $i < j$, both (V_i, V_j) and (U_i, U_j) are $\zeta_{5.5}(k, \delta)$ -regular. Thus, Lemma 5.5 implies that for any γ -regular partition in G (respectively $G[Q]$) $G[Q]$ (respectively G) has an equipartition that is δ -similar to it. We can thus take $q_{5.2}(k, \delta) = q_{5.3}(\ell, \zeta')$ in the statement of the lemma because ℓ and ζ' depend on k and δ . ■

12 Proofs from Section 6

Proof (of Theorem 1): Suppose the regularity-instance R has error parameter γ , $\binom{k}{2}$ edge densities $\eta_{i,j}$, and a set of non-regular pairs \bar{R} . Given $G = (V, E)$ the algorithm for testing the property of having R , samples a set of vertices Q , of size q , where q will be chosen later, and accepts G if and only if the graph spanned by Q is $\frac{\gamma^4 \epsilon}{200k^2}$ -close to satisfying R . In what follows we denote by $G[Q]$ the graph spanned by Q .

Claim 1: If G satisfies R , and $q \geq q_1(\epsilon, k, \gamma)$ then $G[Q]$ is $\frac{\gamma^4 \epsilon}{200k^2}$ -close to satisfying R with probability at least $2/3$.

Proof. If $G = (V, E)$ satisfies R , then V has an equipartition into V_1, \dots, V_k such that for all $(i, j) \notin \bar{R}$ the pair (V_i, V_j) is γ -regular. If we take $q_2(\epsilon, k, \gamma) = q_{5.2}(k, \frac{\gamma^6 \epsilon}{10000k^2})$, then by Lemma 5.2, with probability at least $2/3$ the graph $G[Q]$ will have an equipartition into k sets A_1, \dots, A_k , such that $d(A_i, A_j) = \eta_{i,j} \pm \frac{\gamma^6 \epsilon}{10000k^2}$ for all $i < j$, and if (V_i, V_j) is γ -regular then (A_i, A_j) is $(\gamma + \frac{\gamma^6 \epsilon}{10000k^2})$ -regular. By Lemma 9.2, this means that $G[Q]$ is $\frac{\gamma^4 \epsilon}{200k^2}$ -close to satisfying R . ■

Claim 2: If G is ϵ -far from satisfying R , and $q \geq q_2(\epsilon, k, \gamma)$ then $G[Q]$ is $\frac{\gamma^4 \epsilon}{200k^2}$ -far from satisfying R with probability at least $2/3$.

Proof. We take $q_2(\epsilon, k, \delta) = q_{5.2}(k, \frac{\gamma^4 \epsilon}{200k^2})$. By Lemma 5.2 we get that with probability at least $2/3$ the graph $G[Q]$ is such that if it has a γ' -regular equipartition of order k , then G has an equipartition which is $\frac{\gamma^4 \epsilon}{200k^2}$ -similar to it. We claim that if this event occurs then $G[Q]$ is $\frac{\gamma^4 \epsilon}{200k^2}$ -far from satisfying R , which is what we want to show. Suppose $G[Q]$ satisfies the above property and assume that none the less it is $\frac{\gamma^4 \epsilon}{200k^2}$ -close to satisfying R . Consider the $\frac{\gamma^4 \epsilon}{200k^2} q^2$ edge modifications that make $G[Q]$ satisfy R and consider an equipartition $\mathcal{U} = \{U_1, \dots, U_k\}$ of $G[Q]$, which satisfies R after performing these modifications. As we made at most $\frac{\gamma^4 \epsilon}{200k^2} q^2$ edge modifications, we initially had

$d(U_i, U_j) = \eta_{i,j} \pm \frac{\gamma^4 \epsilon}{200}$. Consider now any $(i, j) \notin \bar{R}$. After these modifications (U_i, U_j) must be γ -regular with density $\eta_{i,j}$. Therefore, after these modifications every pair $U'_i \subseteq U_i, U'_j \subseteq U_j$ satisfying $|U'_i| \geq \gamma|U_i|$ and $|U'_j| \geq \gamma|U_j|$ satisfies $d(U'_i, U'_j) = \eta_{i,j} \pm \gamma$. Hence, before the modifications every such pair satisfied $d(U'_i, U'_j) = \eta_{i,j} \pm (\gamma + \frac{\gamma^2 \epsilon}{200})$. Note that this means that every such pair was originally $(\gamma + \frac{\gamma^2 \epsilon}{100})$ -regular. By our assumption on $G[Q]$ this means that G has an equipartition in V_1, \dots, V_k such that $d(V_i, V_j) = \eta_{i,j} \pm \frac{\gamma^2 \epsilon}{50}$ holds for all $i < j$ and for all $(i, j) \notin \bar{R}$ the pair (V_i, V_j) is $(\epsilon + \frac{\gamma^2 \epsilon}{50})$ -regular. By Lemma 9.2, this means that G is ϵ -close to satisfying R contradicting our assumption. ■

Combining the above two claims we infer that if $q \geq \max(q_1(\epsilon, k, \gamma), q_2(\epsilon, k, \gamma))$ then with probability at least $2/3$ the algorithm distinguishes between the required two cases. Furthermore, the number of queries performed by the algorithm depends only on ϵ, k and γ , and is thus upper bounded by a function of ϵ and r . This completes the proof of the theorem. ■

13 Proofs from Section 7

Corollary 13.1 *Triangle-freeness is testable.*

Proof (of Corollary 13.1): By Theorem 2 it is enough to show that triangle-freeness is regular-reducible. Fix any $\delta > 0$ and set $\gamma' = \gamma_{10.4}(\delta, 3)$. Define $\gamma = \min\{\gamma', \delta\}$. We define \mathcal{R} to be all the regularity-instances R satisfying the following: (i) They have regularity parameter γ (ii) They have order at least $1/\gamma$ and at most $T_{2.3}(1/\gamma, \gamma)$ (iii) Their densities $\eta_{i,j}$ are taken from $\{0, \gamma, 2\gamma, \dots, 1\}$. (iv) They do not contain three clusters V_i, V_j, V_k such that $\eta_{i,j}, \eta_{j,k}, \eta_{i,k}$ are all positive.

To show that this is a valid reduction, assume first that G is ϵ -far from being triangle-free. Assume G is $(\epsilon - \delta)$ -close to satisfying a regularity instance $R \in \mathcal{R}$. We can thus make $(\epsilon - \delta)n^2$ edge modifications and get a graph satisfying R . We also remove all edges inside the sets V_i . As by item (ii) each set has size at most $\gamma n \leq \delta n$ we remove less than δn^2 edges. The total number of edges removed is thus less than ϵn^2 . By property (iv) of the regularity instances of \mathcal{R} this means that the new graph is triangle-free, which is impossible because we made less than ϵn^2 edge modifications and G was assumed to be ϵ -far from being triangle-free. Assume now that G is triangle-free. By Lemma 2.3 G has a γ -regular equipartition V_1, \dots, V_k of order $1/\gamma \leq k \leq T_{2.3}(1/\gamma, \gamma)$. Note that by our choice of γ' via Claim 10.4, and because $\gamma \leq \gamma'$, there are no i, j, k such that $(V_i, V_j), (V_j, V_k), (V_i, V_k)$ are γ -regular and $d(V_i, V_j), d(V_j, V_k), d(V_i, V_k) \geq \delta$ because such sets span at least one triangle (in fact, many). As by item (iii) the densities of the instances in \mathcal{R} are taken from $\{0, \gamma, 2\gamma, \dots, 1\}$ we can make at most $\gamma n^2 \leq \delta n^2$ and thus “round down” the densities between the sets into a multiple of γ , while maintaining the regularity of the regular-pairs. This means that the new graph satisfies a regularity-instance $R \in \mathcal{R}$, which means that G was δ -close to satisfying R . ■

Corollary 13.2 *k -colorability is testable.*

Proof (of Corollary 13.2): By Theorem 2 it is enough to show that k -colorability is regular-reducible. Fix any $\delta > 0$ and define \mathcal{R} to be all the regularity-instances R satisfying the following: (i) They have regularity measure δ (ii) They have order at least $1/\delta$ and at most $T_{2.3}(2/\delta, \delta)$ (iii)

Their densities $\eta_{i,j}$ are taken from $\{0, \delta, 2\delta, \dots, 1\}$. (iv) The following graph $T = T(R)$ is k -colorable: if R has order t then T has t vertices, and $(i, j) \in E(T)$ iff $\eta_{i,j} > 0$.

To show that this is a valid reduction, assume first that G is ϵ -far from being k -colorable. Assume G is $(\epsilon - \delta)$ -close to satisfying a regularity instance $R \in \mathcal{R}$. We can thus make $(\epsilon - \delta)n^2$ edge modifications and get a graph satisfying R . We also remove all edges inside the sets V_i . As by item (ii) each set has size at most δn we remove less than δn^2 edges. The total number of edges removed is thus less than ϵn^2 . By property (iv) of the regularity instances of \mathcal{R} this means that the new graph is k -colorable, which is impossible because we made less than ϵn^2 edge modifications and G was assumed to be ϵ -far from being k -colorable. Assume now that G is k -colorable and let V_1, \dots, V_k be the partition of $V(G)$, which is determined by a legal k -coloring of G . Break every set V_i into sets $U_{i,1}, \dots, U_{i,2/\delta k}$ of size $\frac{1}{2}\delta n$. Put all the leftovers from each set in another set L of size $\frac{1}{2}\delta n$. By Lemma 2.3, starting from this equipartition we can get a δ -regular equipartition of G of order at most $T_{2.3}(2/\delta, \delta)$. Note that disregarding the refinement of L the new equipartition must satisfy item (v) in the definition of \mathcal{R} . As by item (iii) the densities of the instances in \mathcal{R} are taken from $\{0, \delta, 2\delta, \dots, 1\}$ we can make at most δn^2 edge modifications and thus “round down” the densities between the sets into a multiple of δ , while maintaining the regularity of the regular-pairs. This means that the new graph satisfies a regularity-instance $R \in \mathcal{R}$, which means that G was δ -close to satisfying R . ■

Corollary 13.3 *Let I be a graph generated by $G(n, 0.5)$. Then, with probability $1 - o(1)$ the property of being isomorphic to I is not testable.*

Proof (of Corollary 13.3): By Theorem 2 it is enough to show that with probability $1 - o(1)$ the property of being isomorphic to I is not regular-reducible. Note, that now there is only one value of n to consider in Definition 2.5 because the property we consider is a property of n -vertex graphs. Consider a graph generated by $G(n, 0.5)$. Clearly, by Lemma 9.1 the bipartite graph on any pair of sets of vertices of size \sqrt{n} has density ≈ 0.5 . We claim that if I satisfies this property then it is not regular-reducible. Suppose it is regular-reducible and consider a small δ , say $\delta = 0.01$. Let \mathcal{R} be the set of regularity-instances, which corresponds to this value of δ . Let G be a graph isomorphic to I . By Definition 2.5 it must be the case that G is δ -close to satisfying some $R \in \mathcal{R}$. By the properties of I this means that most densities of R must be close to 0.5. Let k denote the order of R and let $\eta_{i,j}$ denote its densities. Consider a random k -partite graph on sets of vertices V_1, \dots, V_k each of size n/k , where the bipartite graph connecting V_i and V_j is a random bipartite graph with edge density $\eta_{i,j}$. Clearly this graph is δ -close to satisfying R . On the other hand, it is not difficult to see that as most of the densities $\eta_{i,j}$ should be close to 0.5, then with high probability such a graph must be α -far from being isomorphic to I , for some fixed $\alpha > 0$, say $\alpha = 0.03$. This means that we have a graph that is 0.03-far from satisfying the property and is yet 0.01-close to satisfying one of the regularity-instances of \mathcal{R} . As we chose $\delta = 0.01$, this violates the second condition of Definition 2.5. ■